Erratum to: Differences in monoterpene emission characteristics after ozone exposure between three clones representing major gene pools of *Cryptomeria japonica*

Takafumi MIYAMA^{a,†}, Hiroyuki TOBITA^a, Kentaro UCHIYAMA^a, Kenichi YAZAKI^a, Saneyoshi UENO^a,
Takami SAITO^a, Asako MATSUMOTO^a, Mitsutoshi KITAO^b and Takeshi IZUTA^c

^a Forestry and Forest Products Research Institute, 1 Matsunosato, Tsukuba, Ibaraki 305-8687, Japan

^b Hokkaido Research Center, Forestry and Forest Products Research Institute,
7 Hitsujigaoka, Toyohira, Sapporo, Hokkaido 062-8516, Japan

^c Institute of Agriculture, Tokyo University of Agriculture and Technology, 3-5-8 Saiwaicho, Fuchu, Tokyo 183-8509, Japan

The Editorial Office of Journal of Agricultural Meteorology wishes to correct the errors in this paper (Miyama *et al.* 2018). Figure 3 and the caption should be as followed.

Fig. 3. Comparison of mean basal emission rate (E_b) between plots. Error bars indicate standard error ($n = 6$). Asterisks indicate statistical differences ($p < 0.05$) analyzed by ANOVA. Enriched ozone plots controlled the concentration to twice-ambient. There were significant differences ($p < 0.05$) between Yakushima clone in enriched ozone plots and another.

References

Miyama T, Tobita H, Uchiyama K, Yazaki K, Ueno S, Saito T, Matsumoto A, Kitao M, Izuta T, 2018: Differences in monoterpene emission characteristics after ozone exposure between three clones representing major gene pools of *Cryptomeria japonica*. *Journal of Agricultural Meteorology* **74**(3), 102–108.