351 (1960). 10) Megawa, A., Osaka Daigaku Igaku Zasshi, 12, 771 (1960). 11) Sakai, A., T. Momose, T. Ban and T. Kurotsu, Med. J. Osaka Univ., 10, 327 (1960). 12) Sakai, A., T. Momose, T. Ban and T. Kurotsu, Ibid., 11, 69 (1960). 13) Sakai, A., T. Ban and T. Kurotsu, Ibid., 11, 401 (1961). 14) Yokoyama, S., T. Ban and T. Kurotsu, Ibid., 11, 191 (1960). 15) Hashimoto, P. H., T. Maeda, K. Torii and N. Shimizu, Ibid., 12, 425 (1962). 16) Gillilan, L. A., J. Comp. Neur., 78, 213 (1943). 17) Hashimoto, P. H., T. Abe, Y. Yamada and N. Shimizu, Acta. Anat. Nippon., 37, 67 (1962).

Discussion

- Dr. Uono : Which do you think MAO of the present study, noradrenalin-MAO or serotonin-MAO? It is desirable to differentiate it by means of fluorescein method. And it should be also clarify the character of ChE which localizes in other than nervous tissues, i. e. true or pseudo ChE.
- Dr. Hashimoto : I think MAO of the present study represents both noradrenelin-, and serotonin-MAO. The differentiation of MAO is now investigating biochemically.

Studies on the Neurohumoral Theory-Continued.

The Characteristic on Histochemical Distributions of Monoamine Oxidase and Cholinesterase Activity Concerning the Innervation of Esophagus, and a Consideration on the Etiology of Cardiospasm.

Shigeo OKINAKA, Masaki YOSHIKAWA, Masanori UONO,
Takao MURO, Akihiro IGATA, Hitoshi TANABE,
Satoshi UEDA and Masanori TOMONAGA

*The Third Department of Internal Medicine (Director : Prof. S. Okinaka, M.D.)
Faculty of Medicine, University of Tokyo, Tokyo.*

The distribution of monoamine oxidase (MAO) activity in digestive canal of dogs had previously been investigated histochemically after Takamatsu's potassium tellurite method¹⁰⁾, on the standpoint of the neurohumoral theory, comparing with the distribution of cholinesterase (ChE) activity after Koelle's thiocholine method.^{4,5,6,7,12)}

In muscle layer, media of artery and gland cells, MAO activity was moderately positive and ChE activity was almost negative, while nerve fibers around those tissues showed intense activity of ChE and extremely poor activity of MAO. In Meissner's plexus, MAO activity was poor or almost negative generally in digestive canal. Auerbach's plexus revealed, on the contrary, positive activity in various degrees. Auerbach's plexus of esophagus, especially in the lower part, showed high activity of MAO and contained many "nuclear positive type" nerve cells, which demonstrated not only high activity in the cytoplasm but marked positive reaction outwardly in the nuclei, and which was characteristic

of sympathetic ganglia. In Auerbach's plexus of the stomach, moderate activity of MAO was demonstrated, but "nuclear positive type" cells could hardly be found except in a few groups of nerve cells in the upper part of the cardia. In Auerbach's plexus of the small intestine, moderate activity was observed as was shown in the stomach, but "nuclear positive type" cells were not yet demonstrated. Marked activity of ChE was found generally in nerve cells and fibers both of Meissner's and Auerbach's plexus.

Based on these results,^{9a,14)} this time, distributions of both enzymes in ganglia and nerve fibers concerning the innervation of esophagus, in nerve plexus and effector tissues of esophageal wall were investigated in detail.

Further study was also presented, on the histochemical changes in five cases of cardiospasm surgically resected, and a consideration was given about etiology of the disease.

I Characteristic on histochemical distribution of MAO and ChE activity concerning the innervation of esophagus

Materials and Methods

Potassium tellurite method¹⁰⁾ for MAO and thiocholine method for ChE⁶⁾ were used as usual. The experiments were carried out with human autopsied cases, dogs and also goats as an example of ruminant.

Results

1) Cervical sympathetic ganglia (Ggl. cervicale superius, medius et Ggl. stellatum) and ganglia of vagus nerve (Ggl. nodosum et jugulare) were in relation to the innervation of esophagus, and each ganglion showed both MAO and ChE activity in several degrees. In cervical sympathetic ganglia, high activity of MAO was observed in nerve cells, and most of them were "nuclear positive type" as in other sympathetic ganglia. In Ggl. nodosum, "nuclear positive type" were also partly mixed. These results were almost common among human, dog and goat. Ggl. jugulare, however, showed relatively weak activity of MAO and "nuclear positive type" was not yet detected.

ChE activity was also high in nerve cells and fibers both of sympathetic and vagus ganglia. Most nerve cells of Ggl. nodosum and Ggl. jugulare showed intense activity with sharp contour, but in sympathetic ganglia relatively obscure contour of the nerve cells were demonstrated.

2) Internal and external nerve fibers of esophagus contained generally moderate or intense activity of ChE and extremely poor activity of MAO, which was, however, moderately positive near the ganglia.

3) Striate muscle fiber in the upper part of esophagus, smooth muscle layers and media of arteries showed moderate activity of MAO, and especially marked in small arteries. MAO activity was also relatively intense in the gland cells and duct of esophageal gland (Fig. 1). However, muscle fiber or gland cell itself exhibited almost negative activity of ChE, but nerve tissues surrounding them showed markedly positive activity like a network.

4) End plates in the striate muscle of esophagus, previously demonstrated

Fig. 1 MAO activity in esophagus (esophageal glands, media of arteries and muscle layer). 40× (dog)

Fig. 2 ChE activity in end plates and nerve fibers in esophageal striate muscle. 210× (dog)

by Shiba after Bielschowsk's staining to be quite alike to those of motor nerve in skeletal muscle, contained intense activity of ChE, and in the nerve fibers connected them ChE activity could be observed towards the terminal" (Fig. 2).

5) Elaborate research for MAO activity in the nerve cells of Auerbach's plexus throughout the esophagus of dogs, disclosed that many groups of nerve cells containing "nuclear positive type" were found most frequently in the lower portion of esophagus. In Meissner's plexus, MAO activity was extremely poor, ChE activity in both plexus equally intense. Such a tendency was also same in human autopsied case and goat.

Comments

Those results should be carefully estimated, because MAO in the digestive canal is responsible for the metabolism of serotonin and other amines besides adrenergic transmission, and because "true MAO" (playing an important role directly in the chemical transmission) and "pseudo-MAO" are not yet discriminated still now.^{1,3,8,9b,11,13} Thus, it is required to investigate the distribution of noradrenaline-like fluorescing substance (NAd-FI) histochemically after Eränkö's fluorescence method²⁾ in support of the specificity of MAO.

However, these results were interesting, concerning the innervation of esophagus in following points.

1) The characteristic of the distribution of the enzymes was generally common to human, dog and goat.

2) Distributive pattern of MAO and ChE activity was specific between sympathetic ganglia and those of vagus nerve.

3) In glands, arteries and muscle bundles, localization of MAO (in gland cell or muscle fiber itself) and ChE (in surrounding nerve tissue) could be compared to "the casting and the mould".

4) Meissner's plexus, showing poor activity of MAO and high activity of ChE, was considered to be dominant in cholinergic mechanism. On the other hand, in Auerbach's plexus revealing marked ChE and MAO activity and NAD-Fl, adrenergic transmission as well as cholinergic, is presumed to play an important part, especially in the lower portion of esophagus.

II Histochemical change in Cardiospasm (Idiopathische Oesophagusdilatation)

Materials and Methods

The same method as described above for the histochemistry of MAO and ChE. Five cases of cardiospasm (4, 32 and 42 years old males; 8 and 30 years old females) was selected. Lower part of each esophagus, surgically resected, was investigated histochemically. As the control, human autopsied esophagus without pathological change was used.

Results

MAO activity in the nerve cells of Auerbach's plexus was extremely poor in all materials given. "Nuclear positive type" cells was also hardly found, except a few remained in only one case (Fig. 3). In the control group, MAO activity was observed as usual (Fig. 4). Histological research is now going on in detail.

Changes in Meissner's plexus was not enough to be appreciated.

ChE activity in both plexus seemed to show remarkable changes till now, but more cases were required for further study.

Comments

As for the etiology of this disease, several hypotheses had previously been advanced. Dysfunction of vagus or sympathetic nerve was speculated chiefly on experimental base, but manifest organic change had not yet demonstrated in any clinical case.

In this aspect, reduction of MAO activity and disappearance of "nuclear positive type" cells in Auerbach's plexus of these materials were considered to be a specific finding in the case of cardiospasm.

Clinically, application of both adrenergic agents (epinephrine, ephedrine etc.) and cholinergic agents (pilocarpine, mecholyl, prostigmine etc.) was more effective for the remission of symptoms than single use of cholinergic drugs.

So, based on our results, it may be considered that sympathetic or adrenergic innervation in Auerbach's plexus of lower portion of esophagus related, at least partly, to the etiology of cardiospasm.

Fig. 3 Poor activity of MAO (lacking in "nuclear positive type" cells) in Auerbach's plexus of lower esophagus in a case of cardiospasm. 400 $\times$ (human)

Fig. 4 MAO activity (with "nuclear positive type" cells) in Auerbach's plexus of lower esophagus, control. 400 $\times$ (human)

Acknowledgment

Authors' acknowledgment is due to 1st Department of Surgery, Faculty of Medicine, University of Chiba, and Department of Veterinary Medicine, Nihon University, for the kind offer of valuable materials.

References

- 1) Davison, A. N. : *Physiol. Rev.*, **38** : 729, 1958. 2) Eränkő, O. : *Endocrinol.*, **57** : 363, 1955. 3) Garven, J. D. : *Brit. J. Pharmacol.*, **11** : 66, 1956. 4) Gerebtzoff, M. A. : *Cholinesterase, a histochemical contribution to the solution of some functional problems.* Pergamon Press, 1959. 5) Igata, A. : *Jap. J. Med. Prog.*, **46** : 668, 1959. 6) Koelle, G. B. : *J. Pharmacol. Exp. Therap.*, **103** : 153, 1951. 7) Muro, T. : *Jap. J. Med. Prog.*, **47** : 812, 1960. 8) Okinaka, S. : *The Saishin-igaku*, **15** : 2493, 1960. 9 a) Okinaka, S. et al. : *Lecture in the 47th General Meeting of the Gastroenterological Society of Japan, Tokyo, 1961.* b) Okinaka, S. et al. : *Proc. Jap. Histochem. Ass.* **1** : 60, 1960. 10) Takamatsu, H. : *Tr. Soc. Path. Jap.* **45** : 307, 1956. 11) Tanabe, H. : *Jap. J. Med. Prog.*, **47** : 833, 1960 : **48** : 20 ; **48** : 105, 1961. 12) Toyoda, M. : *Jap. J. Med. Prog.*, **40** : 559, 1953. 13) Uono, M. & Tanabe, H. : *The Saishin-igaku*, **15** : 2298, 1960. 14) Uono, M., Tanabe, H. & Tomonaga, M. : *The Saishin-igaku*, **16** : 946, 1961.

Discussion

Dr. Otori :

1. Are negative the reactions by Takamatsu's method when it was tested anerobically ?

2. Are confirmed identically "nuclear positive type" in serial sections both by Takamatsu's methods ?
3. Potassium tellurite has much lower oxidation-reduction potential than that of Nitro BT. It is questionable to use tellurite for MAO reaction concerning specificity of reactions.

Dr. Tanabe :

1. We have not yet observed the systematic experiment about anaerobiasis and serial sections,
2. In regard to the specificity, we have confirmed by various examinations.

Double Staining Method for Monoamine Oxidase and Cholinesterase

Masanori UONO, Hitoshi TANABE,
Satoshi UEDA and Masanori TOMONAGA

*The Third Department of Internal Medicine (Director: Prof. S. Okinaka, M. D.),
Faculty of Medicine, University of Tokyo, Tokyo.*

For the investigation on neurohumoral theory, we have studied histochemically the distribution of monoamine oxidase (MAO) and cholinesterase (ChE) in various animals. Our hypothesis thereby was that the neuron with ChE activity is cholinergic and the neuron with MAO activity and noradrenaline-like fluorescing is adrenergic. In the course of our study, however, some nerve cells with high activity of MAO were found to show marked reaction not only in the cytoplasm, but also seemingly in the nucleus. Sympathetic ganglia consist mainly cells of this "nuclear-positive" type and this type was supposed to be the characteristic feature of the adrenergic cell.¹⁾ The question arised about the nature of another "nuclear-negative" type of MAO-positive cells and about the relationship of these two types with ChE-positive cells. On the other hand, both MAO and ChE activity were found to be present simultaneously in some ganglia and nuclei of the central nervous system.^{1,2,3,4)} The problem was whether these two enzymes coexisted to the same cell or they were contained in different cells which were mixed up one with the other.

To meet the technical demands in solving these problems a "double staining method" was investigated which would allow to demonstrate the activity of both enzymes on the same tissue section. This method was found to be valuable also for the observation of minute interrelationship between the localization on MAO and ChE.

Preliminary Experiments

Takamatsu's potassium tellurite method for MAO,⁵⁾ and modified Koelle's