- Beitr., 75, 486-588, 1926. 3) Lillie, R.D., Daft, F.S., and Sebrell, W.H.: *Pub. Health Rep.*, 56, 1255-1258, 1941. 4) Masson, K.E., and Emmel, A.F.: *Anat. Rec.*, 92, 33-59, 1945. 5) Stowell, R.E., and Lee, C.S.: *Arch. Parth.*, 50, 519, 1950. 6) Tadaka, R.: *Tr. Soc. Path. Jap.*, 49, 1035-1066, 1960. 7) Teilum, G.: *Ann. Rheumat. Dis.*, 11, 119-136, 1952.

Explanations of figures

The localization of a ceroid-like pigment and a lipofuscin is shown in figures 1-6, using formalin-fixed, paraffin sections of the liver of adult mice, divided into three experimental groups.

- Fig. 1. Sections of group II which was subjected to subcutaneous administration of sodium silicate plus prednisolone. A coarsely globular yellow pigment, occurring in hypertrophied Kupffer cells or phagocytes. An arrow shows a ceroid-like pigment. Hematoxylin and eosin.
- Fig. 2. Sections of group II. A ceroid-like pigment, being Sudan III-positive on formalin-fixed, paraffin sections. Sudan III stain.
- Fig. 3. Sections of group III which was subjected to subcutaneous administration of sodium nucleate plus prednisolone. A ceroid-like pigment, being acid fast fuchsin-positive. Acid fast fuchsin stain.
- Fig. 4. Sections of adult mice, fed with vitamin E deficient diet for 134 days. The ceroid pigment which occurs in the Kupffer cell and liver cell, being shown to be acid fast fuchsin positive. Acid fast fuchsin stain.
- Fig. 5. Sections of group III. A ceroid-like pigment, being chrome alum hematoxylin negative. An arrow shows the pigment. Chrome alum hematoxylin stain.
- Fig. 6. Sections of adult rats in fasting for seven days. A number of fine granular lipofuscin which is chrome alum hematoxylin positive, being scattered in the cytoplasm of liver cells. An arrow shows a lipofuscin. Chrome alum hematoxylin stain.

Distribution of Vitamin A in the Liver

Kenjiro WAKE

Department of Anatomy, Osaka City University Medical School, Osaka

It has been generally accepted that, if large amounts of vitamin A are given to animals, the excess of vitamin A is stored in the Kupffer cells (reticulo-endothelial system) of the liver^{1,4,5,8)}.

In this laboratory, during the course of histological, histochemical and electron microscopic studies on rat liver in hypervitaminosis A, it is clearly demonstrated that vitamin A is not stored in the Kupffer cells, but in the transmittal cells⁶⁻⁹⁾ (interstitial cells) of the autonomic nervous periphery.

Albino rats (weight 150g) and rabbits (weight 2.5kg) were used in this study. Animals were injected subcutaneously with an excess of vitamin A acetate (Chocla A, EISAI, Tokyo). At periods of 1,2,3, and 4 weeks following injections animals were sacrificed by exsanguination.

Fig.

Histochemical demonstration of vitamin A was performed by an A.O. Spencer fluorescence microscope. The Suzuki's silver impregnation method⁶⁹ and the v. Kupffer's gold-chloride method⁷⁰ were used to demonstrate the

Fig.

Fig. 1,2,4, and 5: Liver of a rat 3 weeks after injections with 75,000 I.U. of vitamin A. Fig. 1: The transmittal cells show bright fluorescence which is quickly fading. $\times 200$. Fig. 2: The transmittal cells are intensely impregnated with gold. v. Kupffer's method. $\times 1,200$. Fig. 3: A transmittal cell(t) of rabbit liver 2 weeks after injections with 150,000 I. V. of vitamin A. Note the mesh-like cytoplasm including autonomic axons(a). K=a Kupffer cell with carbon particles. s=hepatic sinusoid. Suzuki's silver method. $\times 2,300$. Fig. 4: Note a swollen transmittal cell(t) whose cytoplasmic processes are withdrawn. k=Kupffer cell. e=endothelial cell. s=hepatic sinusoid. Suzuki's silver method. $\times 2,300$. Fig. 5: An electron micrograph showing a part of a transmittal cell with large fat droplets(f). The cell body is separated from the hepatic sinusoid (s) with a perforated endothelial cell(en). m=mitochondrion. er=erythrocyte. $\times 37,000$.

transmittal cells. For electron microscopy, pieces of rat liver were fixed in osmium tetroxide buffered to a pH of 7.2 and embedded in Epon 812 by the method of Luft²⁹.

The liver of rats injected with a large dose of vitamin A shows numerous scattered cells containing fat droplets. Under a fluorescence microscope these droplets impart bright but quickly fading fluorescence, which is typical of vitamin A destroyed by ultraviolet light. The present study have indicated that the cells which show the vitamin A uptake are transmittal cells of the autonomic endformation in the liver. These cells occur within hepatic cell cords and are set between the hepatic cells. They are found to envelope axons when impregnated with silver. The author, however, is particularly impressed by a remarkable change in structure of the transmittal cells in hypervitamin-

osis A. The cell body swells, the processes are withdrawn and the cytoplasm is reduced to a coarse mesh. The nucleus is compressed by the surrounding fat droplets. By the gold method, the transmittal cells which have taken up a large amount of vitamin A are more intensely impregnated with gold than in the normal condition. This finding suggests the mechanism of the gold method with the aid of *in vitro* test for the reaction of gold chloride to vitamin A ester. In the hepatic cells vitamin A fluorescence is also demonstrable in fat droplets arranged on the cell border adjacent to the sinusoids. Whereas, the Kupffer cells of the reticulo-endothelial system fail to show any vitamin A uptake. When a rabbit is injected subcutaneously with vitamin A and intravenously with Indian ink, the former is taken up by the transmittal cells, while the latter is deposited as carbon particles in the Kupffer cells. Electron microscopically, the striking finding is that the cell filled with masses of lipid are situated outside of the lining of the hepatic sinusoids and are definitely different from the Kupffer cells.

The present findings may shed another light to the possible significance of vitamin A for the autonomic nervous periphery.

References

- 1) Hirt, A. u. K. Wimmer (1939) *Klin. Wochschr.*, 18:733. 2) Kupffer, C. (1876) *Arch. f. mikr. Anat.*, 12: 352. 3) Luft, J.H. (1961) *J. Biochem. Biophysic. Cytol.* 9:409. 4) Popper, H. (1944) *Physiol. Rev.* 24:205. 5) Popper, H. and F. Schaffner (1957) *Liver: Structure and function.* McGraw-Hill, New York. 6) Suzuki, K. (1958) *The experimental therapy.* Vol. 310-320. 7) Suzuki, K. (1963) *Proc. of the 16th general assembly of the Japan medical congress.* April, 1963, Osaka. IV : 13. 8) Tomita, Y. (1955) *J. Kyoto Pref. Med. Univ.* 57 : 489. 9) Watanabe, S. (1960) *J. Osaka City Med. Center* 9 (Suppl. 8) : 99.

A Histochemical Study of Proliferated Bile Ductules in Viral Hepatitis and Posthepatitis Cirrhosis

Kiyowo KOSAKA, Yasuyuki OHTA, Hiroshi ENDO,
Shunji KINUGASA, and Akio ORIME

The First Department of Internal Medicine, Okayama University Medical School, Okayama

Ductural cell proliferation is thought to be a reaction of the liver against injury. It is one of the parenchymal cell reactions appeared in various pathologic conditions in combination with the mesenchymal cell reactions such as Kupffer cell activation. Recently, rather selective proliferation of the cholangiole cell itself was produced by feeding the rat with α -naphthyl isothiocyanate¹⁾. However, primary hepatic cell injury caused by usual hepatotoxic agents and