PURIFICATION AND IMMUNOHISTOCHEMISTRY OF FETAL ACID α -GLUCOSIDASE

TERUO IWAMASA, SEIJI FUKUDA, TETSUO HAMADA
AND NOBUYUKI NINOMIYA

Department of Pathology, Kumamoto University Medical School, Kumamoto 860

Received for publication October 30, 1981

Fetal rat liver acid α -glucosidase was purified more than 20,000 fold from the fetal rat liver 2 days before delivery. The disc gel electrophoresis of this preparation showed a single band of protein. The Michaelis constants of this enzyme were 3.3 mM with maltose and 3.1 mM with 4-methylumbelliferyl α -glucoside as substrates. Molecular weight measured by SDS electrophoresis and TOYOPEARL gel filtration was about 100,000. The pH optima was 4.0 with 4-methylumbelliferyl α -glucoside as substrate and 3.7 with maltose as substrate.

Antibody was obtained by injection of this enzyme into rabbits with Freund's complete adjuvant. The anti-fetal acid α -glucosidase IgG showed the cross-reaction to the adult rat liver acid α -glucosidase. F(ab')₂ was separated by the filtration of Sephacryl S-200 column chromatography after digestion of anti acid α -glucosidase IgG pepsin and then F(ab')₂ was treated with 2-mercaptoethanol. Coupling of F(ab')₂ obtained by use of Sephacryl S-200 column chromatography after treatment of F(ab')₂ with 2-mercaptoethanol, to horseradish peroxidase was performed according to the method of Wilson and Nakane (14).

Light microscopical observation of the immunohistochemical localization of this enzyme in fetal rat and newborn rat liver showed small granular deposits of DAB reaction products in the cytoplasm.

The purification and characterization of adult liver acid α -glucosidase was reported by many authors (1, 3, 4, 5, 10, 11). This enzyme has attracted much attention in connection with Pompe's disease. Hers and his co-workers noted the important role of this enzyme in glycogen hydrolysis in the lysosome system (2, 7). The lysosome system of the liver and muscle in Pompe's disease revealed glycogen autophagy and was termed glycogenosome. The formation of the glycogenosome was considered to be caused by a deficiency of this enzyme in Pompe's disease. On the other hand, glycogenosome were also observed in the newborn rat hepatocytes without deficiency of acid α -glucosidase (8, 9). Further biochemical and histochemical examinations of this enzyme in relation to the formation mechanism of glycogenosome needs investigation.

However, histochemical demonstration of this enzyme has not been reported as yet. Only neutral α -glucosidase has been demonstrated histochemically by Rutenburg (13), Lojda (12) and Gossrau (6) using synthetic substrates. Further-

more, purification and characterization of fetal acid α -glucosidase has not yet been reported. The results reported here show that the purified fetal acid α -glucosidase has a little different K_m value from adult enzyme and immunohistochemically its localization in hepatocytes cytoplasm can be demonstrated in granular appearances indicating lysosomal locations.

MATERIALS AND METHODS

Reagents: Glucose oxidase type II, peroxidase grade VI, maltose grade I and DAB (3,3'-diaminobenzidine tetrahydrochloride grade II) were obtained from Sigma Co., and 4-methylumbelliferyl α -D-glucoside was from Koch-Light Lab. Ltd. Sephadex G-25, Sephadex G-100 and Sephacryl S-200 were purchased from Pharmacia Fine Chemicals. Freund's adjuvant from Difco Lab. and pepsine from Boehringer Co. TOYOPEARL HW-55 was obtained from TOYO SODA Co. Animals; Fetal rat liver was obtained from pregnant Donryu rats killed 2 days before delivery. New-born rat livers were from rats 6 hr after birth.

Assay system: Determination of α -glucosidase activity; The formation of glucose from maltose or glycogen was measured by the method of Hers. The concentration of glucose in the medium was determined using glucose oxidase. Employing 4-methylumbelliferyl α -D-glucoside as a substrate, the reaction was carried out in the presence of 1 mM substrate and stopped by addition of 3 ml of 0.1 M glycine-NaOH buffer, pH 10.7. The fluorescence was determined with an Aminco-Bowman Spectrofluorometer at an excitation wavelength of 365 nm and an emission wavelength of

FIG. 1. Filtration of fetal rat liver acid α -glucosidase on a Sephadex G-100 column.
 ●—● Acid α -glucosidase activity expressed in terms of 4-methylumbelliferone (4-MU) nmoles per hr.
 ○—○ Proteins indicated by measurement of the optical density (OD) at 280 nm.
 A 2.5×70 cm column was used. The flow rate was 4 ml per hr.

450 nm. Determination of protein concentration; Measurement of absorption of 280 nm using Hitachi spectrometer Model 100-20 and Lowry's method for determination of protein concentration were performed.

Other assay methods; Disc gel electrophoresis, SDS electrophoresis and immunodiffusion methods were employed according to standard techniques.

Purification of fetal acid α -glucosidase: Fetal rat livers were homogenated very gently in a Teflon homogenizer with 9 volumes of ice-cold 0.25 M sucrose/1 mM EDTA solution. The homogenate pH was adjusted to 5.0 by the addition of 1.0 N HCl

TABLE 1. *Purification of acid α -glucosidase from fetal rat liver*

Step	Activity (liberated 4-MU,* mM per mg protein/hr)	Purification (fold)	Recovery (%)
Homogenate	0.000731	1	100
10,000 rpm supernatant	0.00181	2.5	72
(NH ₄)SO ₄ precipitate	0.35453	48.5	63
Sephadex G-100 eluate	3.35529	4590	15.7
TOYOPEARL HW-55 eluate	14.69310	20100	2.4

*: 4-methylumbelliferyl α -glucoside.

FIG. 2. Acrylamide disc gel electrophoresis of purified fetal acid α -glucosidase. 100 μ g of the purified enzyme was applied to the gel. The run was carried out at pH 7.0, 2 mA per tube, for 2.5 hr at 4°C.

and cold centrifuged at 3,000 rpm for 30 min. The sediment was resuspended in 1 mM acetate buffer, pH 5.0, and left overnight at 4°C. It was then rehomogenized in a Teflon homogenizer and cold centrifuged at 10,000 rpm for 30 min. To each one liter of this supernatant, 370 g solid ammonium sulfate was added under continuous stirring. After the ammonium sulfate had completely dissolved, the mixture was left at 4°C overnight. The precipitate, collected by centrifugation at 10,000 rpm for 30 min, was dissolved in 25 mM NaCl/1 mM EDTA solution, pH 5.0, and dialysed against the same solution. The some precipitate appearing during the dialysis was centrifuged off. The resultant solution was applied to a Sephadex G-100 column (2.5 $\times$ 70 cm). The Sephadex G-100 was equilibrated with 25 mM NaCl/1 mM EDTA solution, pH 5.0, and the elution was also performed with same solution. The flow rate was 4 ml per hr, and 4 ml fractions were collected. The fetal acid α -glucosidase was adsorbed by the Sephadex G-100 and therefore eluted after majority of protein as described for adult cattle liver by Bruni *et al.* (1, 3, 4). The fractions with no measurable protein were kept and concentrated by ultrafiltration in an Amicon ultrafiltration cell with a Diaflo YM-10 membrane.

Further purification of the fetal enzyme was undertaken using TOYOPEARL HW-55 gel chromatography. The concentrated enzyme solution was loaded onto a TOYOPEARL HW-55 column (1.5 $\times$ 80 cm). The TOYOPEARL HW-55 was equilibrated with 25 mM NaCl/1 mM EDTA solution, pH 5.0, and eluted with the same NaCl-EDTA solution. The eluates between 80 ml and 110 ml were kept and concentrated using a Diaflow YM-10 membrane. Acrylamide disc gel electro-

FIG. 3. A calibration graph for the estimation of molecular weight (M. W.) by SDS-gel electrophoresis. All proteins were electrophoresed under standard conditions.

BSA; Bovin serum albumin, M. W.: 6.7×10^4 .

OA; Ovalbumin, M. W.: 4.3×10^4 .

Ribo A; Ribonuclease A, M. W.: 1.37×10^4 .

The arrow indicates the mobility of purified fetal α -glucosidase.

phoresis of the purified enzyme was carried out at pH 7.0. The effect of pH and antibody on the enzyme activity for maltose, glycogen and 4-methylumbelliferyl α -glucoside were tested using 0.1 M cacodylate buffer. The influence of the antibody on the enzyme activity was measured, after centrifuging off the anti-enzyme IgG-Enzyme complex at 40,000 rpm for one hr at 2°C. The molecular weight, as estimated by TOYOPEARL HW-55 gel filtration and SDS acrylamide electrophoresis, was carried out.

The Michaelis constants of this enzyme were estimated using maltose and

Fig.4a

Fig.4b

FIGS. 4a and 4b.: Double reciprocal plot of acid α -glucosidase activity of purified enzyme, using maltose (4a) and 4-methylumbelliferyl α -glucoside (4b) as substrates. Velocity (v) is expressed as nmoles of glucose (4a) and 4-methylumbelliferone (4b) liberated from the substrates per min per mg of protein.

4-methylumbelliferyl α -glucoside as substrate.

Preparation of antibody and peroxidase-labeled Fab': Purified acid α -glucosidase (2 mg/0.5 ml) was mixed with an equal volume of Freund's complete adjuvant and injected into the foot pad of male New Zealand white rabbits, weighing from 2.5 to 3.0 kg. On each following week, 1 mg of the enzyme was injected into the rabbits with Freund's adjuvant. The antibody produced was identified by the agar diffusion test. To this antibody, purified adult rat liver acid α -glucosidase was also tested on the agar plate. This immune serum and normal rabbit serum as a control were collected and treated with ammonium sulfate (50% saturation). The precipitates were applied to a Sephadex G-25 column for desalting of the ammonium sulfate. They were then applied to a Sephacryl S-200 column (2.5 $\times$ 90 cm) equilibrated with 0.01 M phosphate buffer, pH 8.0, for separation of IgG. The fractions obtained were pooled and concentrated with a Diaflo YM-10 membrane and digested with pepsine (0.3 mg per 10 mg IgG) at 37°C for 20 hr. F(ab')₂ fraction was treated with 2-mercaptoethanol. The Fab' fraction was also separated by Sephacryl S-200 gel filtration. Coupling of Fab' to horseradish peroxidase was carried out according to the method of Wilson and Nakane (14). Peroxidase-labeled Fab' was separated using a Sephacryl S-200 column (1.5 $\times$ 55 cm) which was equilibrated with 0.01 M phosphate buffer, pH 8.0, and eluted with the same buffer.

Section preparation and staining procedures for light microscope: For the light microscopic immunohistochemistry, 1.0 $\times$ 1.0 $\times$ 0.5 cm of the fetal and newborn liver tissues were fixed in ice-cold 10% formaline buffered with phosphate buffer, pH 7.0, for 5 hr. After fixation, they were washed 5 times with ice-cold phosphate buffered saline.

Fig. 5. Influence of pH and antibody on the enzyme activity.

- Enzyme activity measured with maltose as substrate, in absence of antibody.
 - With maltose as substrate in presence of antibody.
 - With glycogen as substrate in absence of antibody.
 - ★—★ With 4-methylumbelliferyl α -glucosidase as substrate in absence of antibody.
- Glu; Glucose liberated by the enzyme activity. 4-MU; 4-methylumbelliferone liberated by the enzyme activity.

with 0.3% H_2O_2 in methanol for 10 min, and rinsed in ice-cold phosphate buffered saline. They were then incubated with peroxidase-labeled Fab' for 30 min at 37°C in a moist chamber. The sections were washed in phosphate-buffered saline for one hr, followed by peroxidase staining (incubation in 100 mg DAB and 0.03% H_2O_2 in 0.05 M Tris-HCl buffer, pH 7.4). After peroxidase staining, the nuclei were stained with Mayer's hematoxylin.

RESULTS

As shown in Fig. 1, the fetal acid α -glucosidase was eluted after the majority of the protein.

The degree and recovery at each step are listed in Table 1. The enzyme was highly purified more than 20,000 fold.

This enzyme obtained was homogeneous and showed a single band on acrylamide gel electrophoresis (Fig. 2).

The molecular weight of this enzyme, determined by the SDS electrophoresis and TOYOPEARL HW-55 gel filtration was found to be about 100,000 (Fig. 3).

FIG. 6. Diffusion analysis on an agar plate of the anti-fetal α -glucosidase IgG.

A; Anti-fetal acid α -glucosidase IgG 25 μ l (3 mg/ml). 1; Purified fetal acid α -glucosidase 25 μ l (3 mg/ml). 2; Purified adult rat liver acid α -glucosidase 25 μ l (1.5 mg/ml). 3; Purified adult rat liver acid α -glucosidase 25 μ l (0.75 mg/ml). 4; Purified adult rat liver acid α -glucosidase 25 ml (0.2 mg/ml).

FIG. 7a. Immunohistochemistry of acid α -glucosidase of the fetal rat hepatocytes, obtained from fetal rats 2 days before delivery. The liver structure was still immature and lobular architecture was not easily singled out. Numerous numbers of the extramedullary hematopoietic cells were observed. $\times 200$.

FIG. 7b. Immunohistochemistry of the acid α -glucosidase of the fetal rat hepatocytes, obtained from fetal rats 2 days before delivery. Granular localization of the acid α -glucosidase was found in the cytoplasm of the hepatocytes. No positive reaction was demonstrated in the hematopoietic cells. After immunohistochemical staining the sections were stained with hematoxylin. $\times 200$

FIG. 7c. Immunohistochemistry of the acid α -glucosidase of the newborn rat hepatocytes 6 hr after birth. Large numbers of extramedullary hematopoietic cells were found too. The immunohistochemical reaction products were also demonstrated in the cytoplasm as granular appearances. $\times 200$

The Michaelis constants of the enzyme, estimated with maltose and 4-methylumbelliferyl α -glucoside as substrates were illustrated in Figs. 4a and 4b. The constants for maltose and 4-methylumbelliferyl α -glucoside were 3.3 mM and 3.1 mM respectively. The effect of pH and antibody on the enzyme activity for maltose, glycogen and 4-methylumbelliferyl α -glucoside appear in Fig. 5. The pH optima for maltose, glycogen and 4-methylumbelliferyl α -glucoside were 3.7, 4.0 and 4.0. The antibody strongly inhibited the enzyme activity. When maltose was used as substrate the enzyme activity was reduced to 30% by use of the antibody.

On the agar plate the antibody showed a positive reaction to the fetal enzyme and also to the adult acid α -glucosidase, prepared separately by our laboratory (Fig. 6).

Light microscopic demonstration of fetal enzyme on the fetal rat liver, obtained 2 days before delivery, and newborn rat liver, 6 hr after birth is shown in Figs. 7a, 7b and 7c by use of immunohistochemical method.

Small granular reaction products were observed on the hepatocytes. At these stages the rat liver showed an immature structure and the hepatic lobules were not yet completely developed. Some hepatic cells were binucleated. Extramedullary hematopoietic cells were prominently observed in the sinusoidal areas. Immunohistochemically, reaction products appeared as small granular depositions in the hepatic cells. However, no reaction product was found in the hematopoietic cells. Hepatocytes in all areas showed similar positive reactions.

In the hepatic cell cytoplasm, reaction products were distributed diffusely in granular appearances.

Newborn rats also showed the same localization of the enzyme in the hepatocytes obtained from the rats 6 hr after delivery.

DISCUSSION

The fetal acid α -glucosidase was slightly different from adult enzyme in relation to the Michaelis constant. The K_m value of the fetal enzyme was a little larger than that of adult enzyme. However, the molecular weight and influence of the pH on the enzyme activity were very similar to the adult enzyme. Furthermore, the anti-fetal acid α -glucosidase IgG revealed a cross-reaction to the adult enzyme. This indicated that the fetal enzyme protein had the same similarity to the adult enzyme in respect to the immunological site. The immunohistochemistry of the enzyme showed granular appearances which were considered to indicate the lysosomal localization of this enzyme. In the hepatic lobules the enzyme was not unevenly distributed. All of the hepatocytes revealed a similar localization of the enzyme.

In the newborn rat hepatocytes a similar lysosomal localization of the enzyme was demonstrated.

Previously we reported that the glycogenosomes were frequently observed in the newborn rat hepatocytes, about 6 hr after delivery, without deficiency of the acid α -glucosidase (8, 9).

Furthermore about 6 hr after birth, the acid α -glucosidase activity was rather high. Hereafter it is necessary to clarify whether the localization of the acid α -glucosidase is in the glycogenosome or not.

In this paper, it was demonstrated that numerous granular localization of the

enzyme was found in the hepatocytes. The lysosomal localization was indicated. However, electron microscopical immunohistochemistry must be done in relation to this problem.

REFERENCES

1. Auricchio, F., Bruni, C. B. and Sica, V.: Further purification and characterization of acid α -glucosidase. *Biochem. J.* 108; 161, 1968.
2. Baudhuin P., Hers, H. G. and Loeb, H.: An electron microscopic and biochemical study of type II glycogenosis. *Lab. Invest.* 13; 1139, 1964.
3. Bruni, C. B., Auricchio, F. and Covelli, I.: Acid α -D-glucoside glucohydrolase from cattle liver. Isolation and properties. *J. Biol. Chem.* 244; 4735, 1969.
4. Bruni, C. B., Sica, V., Auricchio, F. and Covelli, I.: Further kinetic and structural characterization of the lysosomal α -D-glucoside glucohydrolase from cattle liver. *Biochim. Biophys. Acta* 212; 470, 1970.
5. De Barsy, J., Jacquemin, P., Devos, P. and Hers, H. G.: Rodent and human acid α -glucosidase purification, properties and inhibition by antibodies. Investigation in type II glycogenosis. *Eur. J. Biochem.* 31; 156, 1972.
6. Gossrau, R.: Localization of glycosidase with naphthyl substrates. *Histochem. J.* 8; 271, 1976.
7. Hars, H. G.: α -Glucosidase deficiency in generalized glycogen storage disease (Pompe's disease). *Biochem. J.* 86; 11, 1963.
8. Iwamasa, T., Tsuru, T., Hamada, T. and Takeuchi, T.: Physicochemical and ultrastructural studies on glycogenosomes in newborn rat hepatocytes. *Path. Res. Pract.* 167; 363, 1980.
9. Iwamasa, T. and Takeuchi, T.: The mechanism of the formation of glycogenosome in the newborn rat hepatocytes. *J. Histochem. Cytochem.* 27; 714, 1979.
10. Jeffrey, P. L., Brown, D. H. and Illingworth-Brown, B.: Studies of lysosomal α -glucosidase. I. Purification and properties of the liver enzyme. *Biochemistry* 9; 1403, 1970.
11. Jeffrey, P. L., Brown, B. H. and Illingworth-Brown, B.: Studies of lysosomal α -glucosidase. II. Kinetics of action of the rat liver enzyme. *Biochemistry* 9; 1416, 1970.
12. Lojda, Z.: Some remarks concerning the histochemical demonstration of disaccharidases and glucosidases. *Histochem.* 5; 339, 1965.
13. Rutenburg, A. M., Goldberg, J. A. Rutenberg, S. H. and Lang, R. T.: The histochemical demonstration of α -glucosidase in mammalian tissues. *J. Histochem. Cytochem.* 8; 268, 1960.
14. Wilson, M. B. and Nakane, P. K.: Recent developments in the peroxidase method of conjugating horseradish peroxidase (HRPO) to antibody. In *Immunofluorescence and Related Staining Techniques*, ed. by W. Knapp, K. Holuber and G. Wick, Elsevier/North-Holland Biochemical Press, Amsterdam, 1978, p. 215-224.