HISTOCHEMICAL IDENTIFICATION OF CALCIUM OXALATE

TAKAO YASUE

Department of Pathology, Gifu University School of Medicine, Gifu

(Director: Prof. AKITSUGU OJIMA)

Received for Publication August 2, 1969

Histochemistry of calcium oxalate has not been definitive. The present author has critically investigated various histochemical methods for calcium salts and finally found a new method advantageous to those hitherto reported.

Celloidin models of calcium salts, rat kidneys of oxalosis induced with ethylene glycol and human materials were used for study.

New method: Firstly calcium phosphate and carbonate are removed by immersing sections into 5% acetic acid for 30 minutes. Secondly the sections still containing calcium oxalate are made to react with 5% aqueous silver nitrate for 15 minutes, and thirdly the reaction product appears dark brown to black with the use of rubeanic acid (saturated rubeanic acid in 70% alcohol with 2 drops of strong ammonium for 1 minute) that is very sensitive to silver without losing the definite localization of calcium oxalate. This method has a merit; it does not depend on weather or illumination and takes a shorter time than the Kóssa method.

A confident histochemical method for identifying calcium oxalate was established by combining this method with other characteristics of calcium oxalate such as the unstainability with hematoxylin, birefringency, solubility after incineration and gypsum formation.

In the present study calcium oxalate was also found to react with lead besides silver and to appear brownish-black when one uses lead nitrate and ammonium polysulphide.

Deposition of calcium oxalate crystals in the kidneys has been recently reported not only in oxalosis but also in cases of uremia, renal transplantation and artificial dialysis, and its significance presented a new problem. However identifying the crystals has depended on an uncertain chemical analysis or on an x-ray diffraction method which is not favorable practically and economically.

The present author found numerous crystals of possible calcium oxalate in the kidneys of an autopsy case of Alport's syndrome (41) and tried to prove them histochemically, and realized that no histochemical method for calcium oxalate has been established yet. It is the motive of this study.

Though various histochemical methods for calcium are reported, it seems to be difficult to demonstrate calcium oxalate histochemically due to its chemical stability. McGee-Russell (1958) (25) examined reactivity of various staining dyes with calcium oxalate and found that only quinalizarin and silver nitrate were useful. Von Kóssa (19) method, though it is used as the most commonly for calcium, does not demonstrate calcium cation itself but the anion composing the calcium salts. Pizzolato (1964) (30) in his review pointed out that stainability of calcium oxalate with the von Kóssa method was varied depending on reporters and about half of them obtained positive but uncertain results. Since Johnson *et al.* (1962) (17) and

Wolman *et al.* (1962) (40) failed to demonstrate calcium oxalate with the von Kóssa method, they presented a technique to stain calcium oxalate after converting it into carbonate by incineration, but a question about specificity and localization remains in the method.

calcium already reported are critically investigated again with special reference to calcium oxalate and finally a new method of histochemical demonstration for calcium oxalate established by the author is described.

MATERIALS

1) Celloidin models of calcium salts

Calcium phosphate tribasic ($\text{Ca}_3(\text{PO}_4)_2$), calcium carbonate (CaCO_3) and calcium oxalate monohydrate ($\text{CaC}_2\text{O}_4 \cdot \text{H}_2\text{O}$) of superior quality were embedded into celloidin and cut with microtome. Sections obtained were used for study as routine tissue sections. (In this paper the names of these models are shown by the chemical sign such as $\text{Ca}_3(\text{PO}_4)_2$, CaCO_3 and CaC_2O_4).

2) Rat renal tissues of experimental oxalosis

Kidneys of experimental oxalosis were obtained from Wistar strain male rats weighing 150 g after being administrated 5% ethylene glycol orally for 2–5 days (30).

3) Human renal tissues containing possible calcium oxalate

The kidneys of an autopsy case of an Alport's syndrome (autopsy No. 782, Gifu University School of Medicine) (41), and the kidneys of a renal transplantation case (autopsy No. 1153 *ibid.*). These kidneys contain numerous colorless or faintly yellow, translucent, birefringent, and radiated crystals which are not stained with hematoxylin and eosin (Figs. 1 and 2). These crystals are suggested to be calcium oxalate.

4) Renal tissues containing calcium phosphate and possible calcium carbonate

- a. The renal tissue of a myeloma case (autopsy No. 1155 *ibid.*).
- b. The tumor tissue of Gardner's transplantable osteogenic sarcomas of mice which form osteoid or calcification tissue (27).

METHODS

A) Prescribed methods

1. Solubility test:

The solubility of crystals was tested by immersing materials in 5% acetic acid solution or 2% hydrochloric acid solution for 30 minutes.

2. Microincineration method:

Johnson and Pani's method (17). In addition to the original method, incinerated sections were tested for solubility by 5% acetic acid and 2% hydrochloric acid.

Fig. 1. Colorless or faintly yellow, translucent, and radiated crystals of calcium oxalate which are not stained with hematoxylin and eosin. The kidney of a patient received renal transplantation. Hematoxylin and eosin. $\times 200$.

Fig. 2. Birefringent crystals. Partially crossed polars by a polarizing microscope. The same portion with Fig. 1. $\times 200$.

3. *Gypsum formation reaction:*
Schujeninoff's method (34).
4. *Various staining methods:*
 - 1) Silver nitrate substitution.
 - a) Von Kóssa's method (19). b) Pizzolato's method (30).

- 2) Cobalt substitution. (McGee-Russell's method (25)).
- 3) Iron substitution. (McGee-Russell's method (25)).
- 4) Copper substitution. (McGee-Russell's method (25)).
- 5) Lead substitution. (Roehl's method using lead acetate (32)).
- 6) Reaction with pyrogallol. (Von Kóssa's method (19)).
- 7) Staining with hematoxylin. (McGee-Russell's method (25)).
- 8) Staining with alizarin red S. (McGee-Russell's method (25)).
- 9) Staining with nuclear fast red. (McGee-Russell's method (25)).
- 10) Staining with quinalizarin. (McGee-Russell's method (25)).
- 11) Staining with sodium rhodizonate. (McGee-Russell's method (25)).

B) New methods developed by the present author

1. *Silver nitrate - rubeanic acid method:*

- 1) Immerse slide in 5% aqueous AgNO_3 for 10–20 minutes.
- 2) Rinse thoroughly with running distilled water.
- 3) Immerse for about 1 minute in saturated rubeanic acid in 70% alcohol, adding 2 drops of strong ammoniac water per 100 ml. Rinse in 50% alcohol and water.
- 4) Counterstain, if required, with methyl green or safranin O.
- 5) Dehydrate, clear and mount in balsam.
- 6) Calcium deposits are colored dark brown or black.

2. *Lead nitrate - ammonium polysulphide method:*

- 1) Bring slide to distilled water.
- 2) Immerse 30 minutes in 10% aqueous $\text{Pb}(\text{NO}_3)_2$.
- 3) Rinse thoroughly with running distilled water for 1 minute.
- 4) Immerse in dilute ammonium polysulphide solution for 1 minute.
- 5) Rinse in water, counterstain, if required, with methyl green or safranin O.
- 6) Dehydrate, clear and mount in balsam.
- 7) Calcium deposits are colored brownish-black.

3. *Nickel nitrate - dimethyl glyoxime method:*

- 1) Bring slide to distilled water.
- 2) Immerse 20 minutes in 5% aqueous $\text{Ni}(\text{NO}_3)_2$.
- 3) Rinse thoroughly with running distilled water.
- 4) Immerse 1 minute in a mixture of the following solution:
Saturated dimethyl glyoxime in 70% alcohol.....100 ml.
Strong ammoniac water1 ml.
- 5) Rinse with a stream of water.
- 6) Dehydrate, clear and mount in balsam.
- 7) Calcium deposits are colored pink.

RESULTS

1. Solubility test, microincineration method and gypsum formation reaction

The results are shown in Table 1. CaCO_3 produced gas bubbles and rapidly dissolved with acetic acid and hydrochloric acid, and $\text{Ca}_3(\text{PO}_4)_2$, calcium deposition in the myeloma kidney and in the osteosarcoma dissolved without gas bubbles using

TABLE 1
Solubility and gypsum formation reaction

Materials Tests	Celloidin models			Rat Kidney with oxalosis	Kidney of Alport syn- drome	Kidney of renal trans- planta- tion	Mye- loma kidney	Osteo- sarcoma of mice
	$\text{Ca}_3(\text{PO}_4)_2$	CaCO_3	CaC_2O_4					
5% acetic acid	S	S (G)	—	—	—	—	S	S
5% acetic acid after incineration			S (G)	S (G)	S (G)	S (G)	S	S
2% hydrochloric acid	S	S (G)	S	S	S	S	S	S
2% hydrochloric acid after incineration			S (G)	S (G)	S (G)	S (G)	S	S
Gypsum formation	+	+(G)	+	+	+	+	+	+

S: soluble, —: insoluble, G: gas formation

TABLE 2
Stainability of calcium salts

Materials Reagents	Celloidin models			Rat kidney with oxalosis	Kidney of Alport syn- drome	Kidney of renal trans- planta- tion	Mye- loma kidney	Osteo- sarcoma of mice
	$\text{Ca}_3(\text{PO}_4)_2$	CaCO_3	CaC_2O_4					
Hematoxylin	+	+	—	—	—	—	+	+
Alizarin S	+	+	—	—	—	—	+	+
Nuclear fast red	+	+	—	÷	÷	÷	+	+
Quinalizarin	+	—	—	—	—	—	+	+
Sodium rhodizonate	+	+	—	—	—	—	+	+
Pyrogallol	+	—	—	—	÷	—	+	+
$\text{Co}(\text{NO}_3)_2$	+	—	—	—	—	—	+	+
FeCl_3	+	÷	—	—	—	—	+	+
CuSO_4	+	÷	+	*	*	*	+	+
$\text{Pb}(\text{CH}_3\text{COO})_2$	+	÷	—	—	—	—	+	+
AgNO_3	++	÷	+	+	+	+	++	++
$\text{AgNO}_3 + \text{H}_2\text{O}_2$	++	+	+	+	+	+	++	++
$\text{Ni}(\text{NO}_3)_2$	+	—	—	*	*	*	+	+
$\text{Pb}(\text{NO}_3)_2$	+	+	+	+	+	+	+	+
$\text{AgNO}_3 + \text{Rubeanic acid}$	++	++	++	++	++	++	++	++
After treatment with 5% acetic acid	$\text{Pb}(\text{NO}_3)_2$	—	—	+	+	+	—	—
	AgNO_3 + Rubeanic acid	—	—	++	++	++	—	—

++: marked stained, +: moderately stained, ÷: slightly stained, —: not stained

*: disappeared through staining

TABLE 3
Summarized results of histochemical identification for calcium
oxalate from other calcium salts.

Materials Tests	Celloidin models			Rat kidney with oxalosis	Kidney of Alport syn- drome	Kidney of renal trans- plant- ation	Mye- loma kidney	Osteo- sarcoma of mice
	$\text{Ca}_3(\text{PO}_4)_2$	CaCO_3	CaC_2O_4					
Gypsum formation	+	+	(G)	+	+	+	+	+
5% acetic acid	S	S	(G)	—	—	—	S	S
5% acetic acid after incineration			+	(G)	+	(G)	+	+
$\text{Pb}(\text{NO}_3)_2$	+	+	+	+	+	+	+	+
AgNO_3	++	+	+	+	+	+	++	++
After treatment with 5% acetic acid	$\text{Pb}(\text{NO}_3)_2$	—	—	+	+	+	—	—
	AgNO_3 + Rubeanic acid	—	—	++	++	++	—	—

The marks indicate the same meaning as in the previous tables.

the same acids. CaC_2O_4 , crystals in the kidneys of the rat with oxalosis, and the kidneys from Alport's syndrome and renal transplantation were all dissolved with hydrochloric acid, but not with acetic acid. However, after incineration these were also rapidly dissolved with gas bubbles by acetic acid.

Gypsum formation reaction which forms specific needle shaped crystals of $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ was positive in all sections. But this reaction showed a defect which is that the reaction products lost their localization.

2. Stainability of the former methods

The results are shown in Table 2. Celloidin models of CaCO_3 , $\text{Ca}_3(\text{PO}_4)_2$, calcium in the kidneys of myeloma and osteosarcoma were stained, with lake pigments or chelate reagents such as hematoxylin, alizarin S, nuclear fast red, quinalizarin and sodium rhodizonate, but only CaCO_3 was not stained with quinalizarin. CaC_2O_4 was not stained with any of them. Crystals in the kidney of the rat with oxalosis, Alport's syndrome and renal transplantation were faintly stained by nuclear fast red only, but not by others.

In the reaction with pyrogallol, CaCO_3 , CaC_2O_4 , crystals in the kidneys of the rat with oxalosis and of renal transplantation were not stained, and crystals in the kidneys of Alport's syndrome were faintly stained, but all the rest of the materials were stained.

In the metal substitution methods using metal salts such as cobalt, iron, copper, lead, and silver, $\text{Ca}_3(\text{PO}_4)_2$ in the kidneys of myeloma and osteosarcoma were all stained. But CaCO_3 did not react to cobalt phosphate, and faintly react to ferric chloride, copper sulfate, lead acetate and silver nitrate. In the Pizzolato's method CaCO_3 was stained yellowish-brown.

However, CaC_2O_4 , and crystals in the kidneys of the rat with oxalosis, Alport's

Fig. 3. Celloidin model of pure calcium oxalate monohydrate crystals stained with the modified von Kóssa method. $\times 100$.

Fig. 4. Celloidin model of pure calcium oxalate monohydrate crystals stained with the lead nitrate-ammonium sulfide method. The fine crystals have a tendency to dissolve during the reaction procedure. $\times 100$.

Fig. 5. Calcium oxalate crystals stained with the modified von Kóssa method. The kidney of a case of an Alport's syndrome. $\times 200$.

syndrome and renal transplantation were not stained by metal salts except for copper sulfate and silver nitrate. CaC_2O_4 was stained with copper sulfate, but crystals in the tissue sections were dissolved and disappeared during treatment, so that judgement was impossible.

In the von Kóssa's method which still has some problems, its stainability was not enough when the sections were immersed into sodium thiosulphate after staining for 30 minutes following the original method. It was strongly influenced by weather conditions or artificial light. Calcium oxalate crystals were not stained under the

Fig. 6. Calcium oxalate crystals stained with the silver nitrate-rubeanic acid method after immersing in 5% acetic acid. The kidney of the rat with oxalosis. $\times 100$.

Fig. 7. Calcium oxalate crystals stained with the silver nitrate-rubeanic acid method after immersing in 5% acetic acid. The transplanted kidney of a case with renal transplantation. $\times 100$.

insufficient illumination conditions. However when the specimens were exposed to daylight for more than 30 to 60 minutes before immersing in sodium thiosulphate after staining for 30 minutes under the best condition, crystals in the celloidin models

and the tissue sections were able to be stained clearly (Figs. 3 and 5). This procedure is almost the same as the modification by McGee-Russell (25).

The stainability of Pizzolato's method was certainly stronger than the von Kóssa's method, but under the insufficient conditions crystals were stained a yellowish-brown color and then blackend after a certain amount of time passed.

3. The stainability with the new methods

In the cases using nickel nitrate, $\text{Ca}_3(\text{PO}_4)_2$, calcium deposit in the kidneys of myeloma and the osteosarcoma were stained, but CaCO_3 and CaC_2O_4 were not stained, and crystals from the kidneys of others experimented on disappeared while staining.

In the cases using lead nitrate, crystals in both the tissue sections and celloidin models were stained and their localization was accurate, but it was observed that the fine crystals had a tendency to dissolve during the reaction (Fig. 4).

In the cases using the silver nitrate-rubeanic acid method, the preparations with very clear localization of stained crystals were obtained within a shorter time than the von Kóssa method including all materials (Figs. 6 and 7). Since this reaction was very sensitive and not influenced at all by illumination, this staining method was easily undertaken independent of weather, day or night and illuminating apparatus, and the result was excellent.

When the silver nitrate-rubeanic acid method (or the lead nitrate-ammonium polysulphide method) was used after dissolving calcium phosphate and calcium carbonate from tissue sections by acetic acid, CaC_2O_4 and crystals in the kidneys of the rat oxalosis, Alport's syndrome and renal transplantation were clearly stained, but $\text{Ca}_3(\text{PO}_4)_2$, CaCO_3 , and calcium in the kidneys of myeloma and osteosarcoma, which had been already dissolved, were not demonstrated.

DISCUSSION

Calcium phosphate is the most common form (85–90%) among calcium deposits in human tissues and small percentage of them is calcium carbonate (10–15%). Accordingly the majority of calcium histochemistry hitherto studied was on calcium phosphate and carbonate, and so various methods for calcium investigated in this paper were all effective for demonstration of calcium phosphate. However, as for calcium oxalate, it is no exaggeration to say that a specific histochemical method for it has not been established, because the crystals were not stained with histochemical methods for calcium hitherto employed due to their very low ability to react.

Solubility test: The results of the solubility tests of calcium oxalate by many reporters (1, 2, 5, 8, 11, 16, 23, 31, 43) nearly coincided, that is, it is soluble in such acids as hydrochloric acid, sulfuric acid, nitric acid, and periodic acid, however, it does not dissolve in acetic acid, sodium hydrochloride, lithium carbonate, ether, alcohol, xylene etc.

Gypsum reaction: This is one of the admirable specific reactions to identify calcium qualitatively, but is not good enough for histochemistry because the reaction products show transition and then their definite localization is lost.

Microincineration method : Johnson *et al.* (17) noticed a unique nature of

calcium oxalate in that it changes to calcium carbonate by incineration and that it can be useful for histochemistry. They described that oxalate of strontium and barium and also succinate, malonate, citrate and maleate of calcium should be distinguished from calcium oxalate, however, that the incineration method is effective to prove calcium oxalate because of no existence of other salts in the tissues. Results of the present investigation on the solubility test, gypsum formation test and microincineration coincided with those of previous researchers, but these methods play only an insignificant role and not a specific one for demonstration of calcium oxalate.

Among such staining reagents for calcium as lake pigments, chelate reagents, pyrogallol and naphthalo-hydroxamic acid (38), only alizarin S, quinalizarin and naphthalo-hydroxamic acid show some affinity to calcium oxalate. Macaluso (23) reported that birefringency of the crystals in the sections increased with the use of naphthalo-hydroxamic acid. However, the author supposes that this phenomenon is not favorable to differentiate crystals, because the crystals show birefringence by nature. McGee-Russell (25) found that only quinalizarin among lake pigments was able to stain calcium oxalate contained in the cockroach. In spite of a very careful examination using quinalizarin the author failed to demonstrate calcium oxalate in both celloidin models and tissues. Hematoxylin is commonly used for calcium staining, but most workers (1, 2, 3, 9, 12, 16, 23, 30, 31, 33, 42, 43) reported its negative stainability on calcium oxalate, however, some (7, 11, 13) indicated a very weak reaction on it. Regarding the latter the author thinks that the matrix of the crystals or small amount of other ion such as phosphate contained in the crystals may be faintly stained.

The lake pigments, chelate reagents and staining with pyrogallol indicated in Table 2 were not able to demonstrate not only pure calcium oxalate of celloidin models but also calcium oxalate in the kidneys of experimental oxalosis, Alport's syndrome and of renal transplantation. Nuclear fast red and pyrogallol faintly stained the surface of the crystals of calcium oxalate in the present study, however, it is highly possible that the matrix or contaminations of the crystals were stained.

Metal salts such as copper sulphate (25, 32, 37), ferric chloride (25, 32, 35, 37, 39), ferrous sulfate (25, 39), cobalt nitrate (15, 37), lead acetate (32, 37), ammonium molybdate (32), and silver nitrate (4, 14, 19, 20, 22, 25, 30, 35) are used for metal substitution methods of calcium histochemistry. Strictly speaking these metal substitution methods depend on a reaction to substitute calcium, which is bound to anion, with other heavy metals, and therefore it does not detect calcium cation directly but such anion as carbonate, phosphate, sulfate and oxalate. Among these metal salts only silver nitrate was reported to react to calcium oxalate.

The famous method of von Kóssa (1901) which uses silver nitrate had been already used as a method to demonstrate calcium by Flesh (1885) and Lemberg (1892) (4). Since then many modification methods were devised and the method has been recognized as the most useful one for calcium histochemistry (4, 14, 18, 22, 25, 30, 35, 37). However, as was mentioned previously, this method is not specific for calcium cation but for phosphate, carbonate, oxalate, sulfate, sulphocyanide (22), chloride (35) and oleate (4) of calcium, for phosphate of strontium, barium and iron (ferric), for phosphate and carbonate of copper (4), for magnesium phosphate (4), and for albuminate (19). As for calcium oxalate, results from the

von Kóssa method varied: Cameron (4), Scowen *et al.* (36), Hughes (16), Fanger *et al.* (10), McGee-Russell (25), Richter *et al.* (31), Cogan *et al.* (5), Glynn (12), Neustein *et al.* (26), Edwards (8), Dunn (7), reported positive results, and Shepard and co-workers (33) found that the results were variable. McGee-Russell (25) observed that silver particles deposited on the surface of calcium oxalate in cockroaches and Cameron (4) also reported a positive result using gelatin models of calcium oxalate with the von Kóssa method. On the other hand, negative results were reported by Lund *et al.* (21), Davis *et al.* (6), Benington *et al.* (1), Enger *et al.* (9), Pizzolato (30), Ying Chou *et al.* (42), Wolman *et al.* (40), Burke *et al.* (3), Marshall *et al.* (24), Zimmerman (43), Gershoff *et al.* (11), Johnson (17) and Bennett *et al.* (2), and the number of negative reports was a few more than the positive ones. No definite view has been established about stainability of calcium oxalate with the von Kóssa method.

The author confirmed that CaC_2O_4 as well as the crystals in the tissue were able to be stained with the modified Kóssa method as shown in Fig. 3 by the present investigation. Calcium oxalate was not stained or its stainability was inconsistent when treated following the original method. Negative results due to this reason may be included among many negative results hitherto reported.

Pizzolato (30) reported that unstainable crystals of calcium oxalate became to be stainable by pretreatment of hydrogen peroxide and explained that this change depended on the conversion of calcium oxalate to carbonate *in situ*, however, the author could not agree with his opinion. It is presumed that the weak ability to react between calcium oxalate and silver nitrate was only strengthened with hydrogen peroxide.

In order to develop a new method the author paid attention to the next points: 1) Nickel nitrate and lead nitrate were used because these satisfy the necessary and sufficient conditions that new metal salts produced after substitution reaction were more insoluble than previously existing calcium salts. 2) Rubeanic acid (dithio-oxamide) was used as a reagent which was able to detect clearly the least amount of silver reacted (29).

Nickel nitrate was effective in staining salts composed of calcium phosphate, but not calcium oxalate.

The lead nitrate substitution method was correct theoretically and the method was easy, but the crystals escaped from the sections during the procedure (Fig. 4), so that it was not favorable for histochemistry. On the contrary the silver nitrate-rubeanic acid method proved to have excellent result.

Rubeanic acid's nature is to react with copper, silver, cobalt, nickel and mercury, and to form colored double salts, and Okamoto *et al.* (28, 29) found a method to demonstrate copper and silver using this nature. Brownish-black insoluble double salt is produced with a reaction of rubeanic acid and silver, and the reaction is very sensitive and specific if there is no other salt. Slightly deposited silver particles on the surface of crystals of calcium oxalate with a low ability to react are assumed to be blackened with rubeanic acid, and its sensitivity as well as localization are good enough for demonstration of calcium oxalate (Figs. 6 and 7). Besides, this method has other merits: an accurate result can be obtained briefly in spite of its easy procedure and it does not depend upon weather, time and illumination conditions compared with the von Kóssa or Pizzolato methods which depend

on those factors. If acetic acid treatment to dissolve calcium phosphate and carbonate was undertaken before application of this method, only calcium oxalate was distinctly stained, and possible calcium phosphate and carbonate were not stained completely. This reaction can not be a specific one for calcium oxalate, however, calcium oxalate crystals are certainly demonstrated histochemically when optical characteristics (birefringence), hematoxylin and eosin staining, solubility test, microincineration method and gypsum formation reaction are combined to result in this reaction (Table 3).

Finally the present author regards the following method to be the best one to prove calcium oxalate histochemically.

Histochemical identification of calcium oxalate:

1. Crystals not stained with hematoxylin and eosin.
2. Radiative crystals with birefringence.
3. Positive gypsum formation reaction.
4. Soluble in 2% hydrochloric acid, but not soluble in 5% acetic acid.
Crystals dissolve releasing gas bubbles when microincinerated before immersing in these acids.
5. Among the calcium salts only calcium oxalate is stained black with the silver nitrate-rubeanic acid method after eliminating calcium phosphate and carbonate with 5% acetic acid.

REFERENCES

1. Benington, J. L., Haber, S. L., Smith, J. V. and Warner, N. E.: Crystals of calcium oxalate in the human kidney. *Am. J. Clin. Path.* 41; 8, 1964.
2. Bennett, B., and Rosenblum, C.: Identification of calcium oxalate crystals in the myocardium in patients with uremia. *Lab. Invest.* 10; 947, 1961.
3. Burke, E. C., Baggenstoss, A. H., Owen, C. A., Power, M. H. and Lohr, O. W.: Oxalosis, *Pediatrics.* 15; 383, 1955.
4. Cameron, G. R.: The staining of calcium. *J. Path. Bact.* 33; 929, 1930.
5. Cogan, D. G., Kuwabara, T., Silbert, J., Kern, H., McMurray, V. and Hurlbut, C.: Calcium oxalate and calcium phosphate crystals in detached retinas. *Arch. Ophth.* 60; 366, 1958.
6. Davis, J. S., Klingberg, W. G. and Stowell, R. E.: Nephrolithiasis and nephrocalcinosis with calcium oxalate crystals in kidneys and bones. *J. Pediat.* 36; 323, 1950.
7. Dunn, H. G.: Oxalosis. *Am. J. Dis. Child.* 90; 58, 1955.
8. Edwards, D. L.: Idiopathic familial oxalosis. *Arch. Path.* 64; 546, 1957.
9. Enger, E., Serck-Hanssen, A. and Rokkens, T.: Oxalosis. *Acta Med. Scand.* 177; 409, 1965.
10. Fanger H. and Esparza, A.: Crystals of calcium oxalate in kidneys in uremia. *Am. J. Clin. Path.* 41; 597, 1964.
11. Gerschoff, S. N., Faragalla, F. F., Nelson, D. A. and Andrus, S. B.: Vitamin B₆ deficiency and oxalate nephrocalcinosis in the cat. *Am. J. Med.* 27; 72, 1959.
12. Glynn, L. E.: Crystalline bodies in the tunica media of a middle cerebral artery. *J. Path. Bact.* 51; 445, 1940.
13. Godwin, J. T., Fowler, M. F., Dempsey, E. F. and Henneman, P. H.: Primary hyperoxaluria and oxalosis. *New. Engl. J. Med.* 259; 1099, 1958.
14. Gomori G.: Der mikrotechnische Nachweis unlöslicher Kalksalze in den Geweben. *Virchows Arch.* 286; 682, 1932.
15. Gomori, G.: Microtechnical demonstration of phosphatase in tissue sections.: *Proc. Soc. Exper. Biol. Med.* 42; 23, 1939.

16. Hughes, D. T. D.: The clinical and pathological background of two cases of oxalosis. *J. Clin. Path.* 12; 498, 1959.
17. Johnson, F. B. and Pani, K.: Histochemical identification of calcium oxalate. *Arch. Path.* 74; 347, 1962.
18. Klotz, J. *Exper. Med.* 7; 633, 1905. Cited by Cameron (4).
19. Kóssa, J. von.: Ueber die im Organismus künstlich erzeugbaren Verkalkungen. *Beit. Path. Anat.* 29; 163, 1901.
20. Lison, L.: *Histochemie et Cytochemie Animales*. 3e édition, Gauthier-Villars, Paris, 1960. Translated in Japanese by T. Imaizumi. Hakusui-sha. Tokyo, 1966.
21. Lund, T. and Reske-Nielsen, E.: Nephrolithiasis and nephrocalcinosis with calcium oxalate crystals in the kidneys and other organs. *Acta Path. Microbiol. Scand.* 38; 353, 1956.
22. Macallum, A. B.: On the distribution of potassium in animal and vegetable cells. *J. Physiol.* 32; 95, 1905.
23. Macaluso, M. P. and Ber, N. O.: Calcium oxalate crystals in kidneys in acute tubular nephrosis and other renal diseases with functional failure. *Acta Path. Microbiol. Scand.* 46; 197, 1959.
24. Marshall, V. F. and Horwith, M.: Oxalosis. *J. Urol.* 82; 278, 1959.
25. McGee-Russell, S. M.: Histochemical methods for calcium. *J. Histochem. Cytochem.* 6; 22, 1958.
26. Neustein, H. B., Stevenson, S. S. and Krainer, L.: Oxalosis with renal calcinosis due to calcium oxalate. *J. Pediat.* 47; 624, 1955.
27. Ojima, A.: Growth-inhibiting and differentiation-promoting activities by cortisone and host tissue environments in transplantable osteogenic sarcoma of mice. *Abstracts of 9th Intern. Cancer. Cong.* 340, 1966.
28. Okamoto, K. and Utamura, M.: Biologische Untersuchungen des Kupfers. *Acta Scholae Med. Kioto.* 20; 573, 1938.
29. Okamoto, K., Utamura, M. and Akagi, T.: Biologische Untersuchungen des Silbers. *Acta Scholae Med. Kioto.* 22; 361, 1939.
30. Pizzolato, P.: Histochemical recognition of calcium oxalate. *J. Histochem. Cytochem.* 12; 333, 1964.
31. Richter, M. N. and McCarty, K. S.: Anisotropic crystals in the human thyroid gland. *Am. J. Path.* 30; 545, 1954.
32. Roehl, W.: Über Kalkablagerung und -Ausscheidung in der Niere. *Beit. Path. Anat.* 7; 456, 1905.
33. Shepard, T. H., Creighton, S. A., Krebs, E. G., Lee, L. W. and Thuline, H. C.: Primary hyperoxaluria. *Pediatrics*, 25; 582, 1960.
34. Schujeninnoff, S.: Zur Frage der Kalkablagerung in den Quergestreiften Muskeln-Muskelverkalkung in Genachten Wunden. *Z. Heilk.* 18; 79, 1897.
35. Schuscik, O.: Über die Methoden zum mikroskopischen Nachweis von Kalk im ossifizierenden Skelett. *Z. Wiss.* 37; 215, 1920.
36. Scowen, E. F., Stansfeld, A. G. and Watts, R. W. E.: Oxalosis and primary hyperoxaluria. *J. Path. Bact.* 77; 195, 1959.
37. Stoelzner, W.: Über Metallfärbungen verkalkter Gewebeteile. *Virchows Arch.* 180; 362, 1905.
38. Voigt, G. E.: Ein neuer histotopochemischer Nachweis des Calciums (mit Naphthalhydroxamsäure). *J. Histochem. Cytochem.* 4; 122, 1957.
39. Wigglesworth, V. B.: The role of iron in histochemical staining. *Quart. J. Micr. Sci.* 93; 105, 1952.
40. Wolman, M. and Goldring, D.: Histochemical demonstration of calcium oxalate crystals. *J. Histochem. Cytochem.* 10; 505, 1962.
41. Yasue, T.: Alport's syndrome. *Jap. J. Clin. Med.* 25; 2139, 1967 (Japanese).
42. Ying Chou. and Donohue, W. L.: Oxalosis. *Pediatrics*. 10; 660, 1952.
43. Zimmerman, L. E. and Johnson, F. B.: Calcium oxalate crystals within ocular tissues. *Arch. Ophth.* 60; 372, 1958.