Crystal Structure of Phenolphthalein

Hiromasa SUGIURA*, Toshiyuki KATO*, Hitoshi SENDA*, Ko-Ki KUNIMOTO*†,
Akio KUWAE** and Kazuhiko HANAI***

*Department of Chemistry and Chemical Engineering, Faculty of Technology, Kanazawa University,
Kakuma-machi, Kanazawa 920-1192, Japan

**Institute of Natural Sciences, Nagoya City University, Nagoya 467-8501, Japan

***Gifu Pharmaceutical University, Gifu 502-8585, Japan

We have studied the molecular structures of various organic dyes by the vibrational spectroscopies.¹ Our recent research interests have been focused on the inclusion phenomena of the phthalein dyes in various organic hosts. As a part of our study, the X-ray analysis of 3,3-bis(4-hydroxyphenyl)-1(3*H*)-isobenzofuranone, better known as phenolphthalein, was undertaken. The pH dependent color change of phenolphthalein has been ascribed to a structure change from the lactone (I) to the dianionic resonating form (II), as shown in Fig. 1. Phenolphthalein shows vibrational bands characteristic of the structure (I) in the solid state. The OH stretching bands are observed at 3383, 3329 and 3291 cm⁻¹ in the IR spectrum. The intense band at 1737 cm⁻¹ with a shoulder peak at 1718 cm⁻¹ is assigned to the C=O stretching of the lactone group. The corresponding Raman bands are observed at 1737 and 1719 cm⁻¹.

Crystals suitable for X-ray analysis were grown from an aqueous ethanol solution at room temperature. A colorless prism with dimensions 0.6×0.3×0.4 mm was mounted on a glass fiber. All measurements were made on a Rigaku AFC-5R diffractometer with a graphite monochromated Mo K α radiation ($\lambda=0.71069$ Å). The detailed measurement conditions and crystal data are listed in Table 1. The intensity data were collected at 23°C using the ω -2 θ scan technique to a maximum 2 θ of value of 55.0°. A total of 4195 reflections were collected. The intensities of three representative reflections which were measured after every 150 reflections declined by 0.49%. A linear correction was applied to the data to account for this phenomenon. The linear absorption coefficient for Mo K α is 0.8 cm⁻¹. An

empirical absorption correction, based on azimuthal scans of several reflections, was applied, which resulted in transmission factors ranging from 0.98 to 1.00. The data were corrected for Lorentz and polarization effects.

The structure was solved by direct methods.² The non-hydrogen atoms were refined anisotropically. All hydrogen atoms were located from a difference Fourier map and included in the full-matrix least squares refinement. The atomic scattering factors and anomalous dispersion terms were taken from the International Tables for X-ray Crystallography, Vol. IV.³ All calculations were performed using the program TEXSAN crystallographic software package.² Selected positional parameters are listed in Table 2. The molecular structure is shown in Fig. 2, together with the atomic labeling scheme. Selected bond distances and angles are listed in Table 3.

There are two independent molecules in the asymmetric unit (Molecule 1 and Molecule 2). The molecules consist essentially of three groups: an isobenzofu-

Fig. 1 Dissociation equilibrium of phenolphthalein

Table 1 Crystal and experimental data

Formula: C ₂₀ H ₁₄ O ₄
Formula weight: 318.33
Crystal system: orthorhombic
Space group: <i>Pna</i> 2 ₁ <i>Z</i> = 8
<i>a</i> = 19.276(3) Å
<i>b</i> = 14.822(2) Å
<i>c</i> = 11.3884(9) Å
<i>V</i> = 3254(1) Å ³
<i>D</i> _{calc} = 1.299 g/cm ³
No. of reflections used = 2970 (<i>I</i> > 1.20σ(<i>I</i>))
No. of parameters = 542
<i>R</i> = 0.038
<i>R</i> _w = 0.037
Goodness-of-fit = 1.25
(Δρ) _{max} = 0.18 eÅ ⁻³
(Δρ) _{min} = -0.18 eÅ ⁻³
Measurement: Rigaku AFC-5R
Program system: TEXSAN
Structure determination: direct method
Refinement: full-matrix least-squares

† To whom correspondence should be addressed.

Table 2 Fractional coordinates and equivalent isotropic thermal parameters of non-hydrogen atoms

Atom	x	y	z	$B_{eq}/\text{\AA}^2$
O1	0.1304(1)	0.4152(2)	0.0982	4.2(1)
O2	0.4595(1)	-0.0192(2)	0.4275	4.9(1)
O3	0.3255(1)	0.0906(1)	-0.0469(2)	3.4(1)
O4	0.3494(1)	0.0339(2)	-0.2235	5.7(1)
O5	0.4450(1)	0.4361(2)	0.3963(3)	4.7(1)
O6	0.3495(1)	0.4153(2)	-0.4315(2)	4.4(1)
O7	0.2592(1)	0.6303(1)	0.0143(2)	3.22(9)
O8	0.1913(1)	0.7456(2)	0.0592(3)	5.4(1)
C1	0.3497(1)	0.1644(2)	0.0320(3)	3.0(1)
C2	0.2882(1)	0.2268(2)	0.0546(3)	2.9(1)
C3	0.2879(2)	0.2832(2)	0.1519(3)	3.6(1)
C4	0.2351(2)	0.3456(2)	0.1687(3)	3.5(1)
C5	0.1826(1)	0.3527(2)	0.0874(3)	3.1(1)
C6	0.1818(2)	0.2969(2)	-0.0094(4)	3.8(1)
C7	0.2343(2)	0.2350(2)	-0.0253(3)	3.5(1)
C8	0.3794(1)	0.1191(2)	0.1403(3)	2.7(1)
C9	0.3349(2)	0.0811(2)	0.2225(3)	3.7(2)
C10	0.3606(2)	0.0341(2)	0.3176(3)	3.9(2)
C11	0.4312(2)	0.0245(2)	0.3329(3)	3.4(1)
C12	0.4754(2)	0.0612(2)	0.2512(3)	3.9(2)
C13	0.4500(2)	0.1083(2)	0.1564(3)	3.4(1)
C14	0.4029(1)	0.2106(2)	-0.0436(3)	3.1(1)
C15	0.4432(2)	0.2868(2)	-0.0215(4)	4.2(2)
C16	0.4887(2)	0.3152(3)	-0.1065(5)	5.9(2)
C17	0.4935(3)	0.2698(4)	-0.2123(5)	6.9(3)
C18	0.4542(2)	0.1963(4)	-0.2368(4)	5.8(2)
C19	0.4088(2)	0.1669(2)	-0.1498(3)	3.7(1)
C20	0.3604(2)	0.0908(2)	-0.1487(3)	3.9(2)
C21	0.3339(1)	0.6109(2)	-0.0054(3)	2.7(1)
C22	0.3603(1)	0.5603(2)	0.1022(3)	2.6(1)
C23	0.4260(2)	0.5222(2)	0.1004(3)	3.2(1)
C24	0.4534(2)	0.4814(2)	0.1988(3)	3.6(1)
C25	0.4151(2)	0.4772(2)	0.3012(3)	3.2(1)
C26	0.3496(2)	0.5145(2)	0.3042(3)	3.6(2)
C27	0.3229(2)	0.5561(2)	0.2055(3)	3.3(1)
C28	0.3396(1)	0.5590(2)	-0.1191(3)	2.7(1)
C29	0.3254(2)	0.4664(2)	-0.1211(3)	3.1(1)
C30	0.3281(2)	0.4178(2)	-0.2231(3)	3.5(1)
C31	0.3449(2)	0.4604(2)	-0.3271(3)	3.0(1)
C32	0.3580(2)	0.5521(2)	-0.3280(3)	3.2(1)
C33	0.3550(2)	0.6005(2)	-0.2244(3)	2.9(1)
C34	0.3655(2)	0.7046(2)	-0.0070(3)	3.0(1)
C35	0.4331(2)	0.7309(2)	-0.0254(3)	3.6(1)
C36	0.4488(2)	0.8219(2)	-0.0136(4)	4.5(2)
C37	0.3977(2)	0.8838(2)	0.0158(4)	5.4(2)
C38	0.3301(2)	0.8584(2)	0.0322(4)	5.3(2)
C39	0.3147(2)	0.7667(2)	0.0202(3)	3.8(1)
C40	0.2486(2)	0.7185(2)	0.0342(3)	3.8(1)

$$B_{eq} = (4/3) \sum_i \sum_j \beta_{ij} a_i^* a_j^* (\mathbf{a}_i \cdot \mathbf{a}_j).$$

ran ring (A) and two *para*-hydroxyphenyl rings (B and C) attached to the tetrahedral carbon atom in the five-membered lactone ring. Each of the three moieties is almost planar. The two *para*-hydroxyphenyl groups lie on opposite sides of the isobenzofuran plane. The geometry differences between the two molecules are found in the orientations of the planes. The rings B and C are inclined with respect to each other at 71.40° for Molecule 1 and 74.63° for Molecule 2. The rings B and C are also oriented with respect to the isobenzofuran ring A at 76.65° and 73.63°, respectively, for Molecule 1 and 75.18° and 70.16°, respectively, for Molecule 2. The C–O bonds in the five-membered lactone rings, which cleave at alkaline pH, are 1.490(3) and 1.484(3) Å, respectively, for Molecule 1 and Molecule 2. They are longer than the normal lactone C–O single bond value (1.462(2) Å)⁴ and shorter than the value of 1.525(3) Å found for fluorescein.⁵ Fitzgerald and Gerkin⁶ have recently reported the crystal structure of phenolphthalein obtained from ethanol solution ($R=0.045$, $R_w=0.097$). Differences between the present work and the reported result were observed mainly in the molecular geometries of the lactone moieties.

Fig. 2 Molecular structure with the numbering of the atoms. Thermal ellipsoids of the non-hydrogen atoms are scaled to enclose 50% probability. The spheres of the hydrogen atoms are drawn in an arbitrary scale.

Table 3 Selected bond lengths (Å) and angles (°)

Atom	Atom	Distance	Atom	Atom	Distance
O3	C1	1.490(3)	O7	C21	1.484(3)
O3	C20	1.341(4)	O7	C40	1.343(3)
O4	C20	1.216(4)	O8	C40	1.208(4)
C19	C20	1.463(5)	C39	C40	1.469(5)

Atom	Atom	Atom	Angle	Atom	Atom	Atom	Angle
C20	O3	C1	111.3(2)	C40	O7	C21	111.3(2)
O3	C1	C14	101.7(2)	O7	C21	C34	102.4(2)
O3	C1	C8	106.5(2)	O7	C21	C28	107.4(2)
O3	C1	C2	107.7(2)	O7	C21	C22	107.4(2)
O4	C20	O3	121.1(3)	O8	C40	O7	120.2(3)
O4	C20	C19	129.7(3)	O8	C40	C39	131.0(3)
O3	C20	C19	109.2(3)	O7	C40	C39	108.8(3)

Estimated standard deviations in the least significant figure are given in parentheses.

References

1. K. Machida, H. Lee and T. Uno, *J. Raman Spectrosc.*, **8**, 172 (1979).
2. "TEXSAN", TEXRAY Structure Analysis Package, Molecular Structure Corporation, The Woodlands, TX, USA 1985.
3. "International Tables for X-ray Crystallography", Vol. IV, The Kynoch Press, Birmingham, 1974.
4. P. Murray-Rust, J. Murray-Rust and R. F. Newton, *Acta Crystallogr.*, **B35**, 1918 (1979).
5. R. S. Osborn and D. Rogers, *Acta Crystallogr.*, **B31**, 359 (1975).
6. L. J. Fitzgerald and R. E. Gerkin, *Acta Crystallogr.*, **C54**, 535 (1998).

(Received January 14, 1999)

(Accepted February 24, 1999)