ELECTRICAL POTENTIAL OSCILLATION ACROSS A LIQUID MEMBRANE OF WATER-OCTANOL-WATER SYSTEM

KENSUKE ARAI, FUMIYO KUSU and KIYOKO TAKAMURA

Tokyo College of Pharmacy,
1432-1, Horinouchi, Hachioji, Tokyo 192-03, JAPAN

Abstract - Measurement was made of electrical potential oscillation across a liquid membrane comprised of octanol between two aqueous solutions, one containing an anionic, or cationic surfactant and alcohol. Examination was also made of effects of chemical substances added to the aqueous solution.

Key words electrical potential oscillation, liquid membrane, octanol

Electrical potential oscillation was recently observed across the liquid membrane of a water-octanol-water system (octanol membrane) [1]. The octanol membrane may possibly be functionally capable of sensing certain chemical substances. Clarification should thus be made of the effects of chemical substances on oscillation for developing new types of sensors. The present study was conducted to examine the effects of chemical substances such as sodium carboxylates, tetraalkylammonium chlorides and barbiturate sodium salts, on the mode of oscillation.

EXPERIMENTAL

Fig. 1 schematically shows the apparatus for electrical potential oscillation across an octanol membrane. The aqueous solutions (1 ml each), one containing a surfactant and alcohol, were poured into an inverted U-shaped cell. 2 ml of octanol containing a supporting electrolyte were introduced onto the top of the aqueous solutions without stirring. The potential difference between the aqueous phases was measured with a potentiometer connected by a pair of silver-silver chloride electrodes.

Fig. 1 Apparatus for measuring electrical potential oscillation across an octanol membrane: (a) potentiometer, (b) silver-silver chloride electrode, (c) aqueous phase, (d) organic phase.

RESULTS

Electrical potential oscillation across an octanol membrane containing SDS or CTAB Spontaneous oscillation was observed in the presence of 8 mM sodium dodecylsulfate (SDS) and 5.0 M ethanol in the aqueous phase on the right and 5 mM tetrabutylammonium chloride (TBACl) in the organic phase (Fig. 2, A). Oscillation started at about 10 minutes after the organic solution had come into contact with the aqueous solutions. The amplitude and oscillatory period were 150 mV and 0.8 min, respectively. No oscillation could be detected in the absence of the surfactant. Similar oscillation was noted in the presence of 2 mM hexadecyl-

trimethylammonium bromide (CTAB) and 5.5 M ethanol in the aqueous phase on the left and 3 mM 2,4,6-trinitrophenol (picric acid) in the organic phase. Although SDS and CTAB were added to the aqueous phases on the right and left, respectively, potential oscillation with nearly the same wave form were observed with both the anionic and cationic surfactants. Potential oscillation may be induced by repetitive formation and destruction of the adsorption layer of the surfactant at the oil-water interface [2]. SDS was added to the bottom of the cell on the left and CTAB to that on the right. The charge of the surfactant in the aqueous phase was found to be determined only the direction of the pulse.

Effects of chemical substances

Potential oscillation across the octanol membrane containing SDS was measured in the presence of 100 mM sodium carboxylates, 20 mM tetraalkylammonium chlorides or 100 mM barbiturate sodium salts in the aqueous phase on the left (Fig. 2, B-D). Those substances caused the oscillation amplitude, and oscillatory and induction period to change. They may possibly dissolve in the adsorption layer of the surfactant at the oil-water interface, with consequent formation and destruction of the surfactant layer. The adsorption layer of surfactant may become stable due to the presence of these substances, so that destruction and formation of the adsorption layer becomes difficult. This may in turn decrease the number of surfactant molecules moving from the interface into the aqueous solution for each oscillation.

The amplitude of oscillation would consequently change. The solubility of the substances may increase with the degree of their hydrophobicity. A linear relationship between the carbon number of the alkyl chain of the carboxylate ion, that of the tetraalkylammonium ion or partition coefficient of barbiturate and amplitude of potential oscillation was found. The amplitude of oscillation across the octanol membrane may thus decrease with increase in hydrophobicity. The present data will serve as the basis for developing new types of sensors.

REFERENCES

1. K. Arai, F. Kusu and K. Takamura, *Denkikagaku*, **58**, 665(1990)
2. K. Arai, F. Kusu and K. Takamura, *Chem. Lett.*, 1715(1990)

Fig. 2 Electrical potential oscillation across octanol membrane containing 8 mM SDS and 5.0 M ethanol in the aqueous phase on the right and 5 mM tetrabutylammonium chloride in the organic phase and no substances (A), 100 mM sodium butylate (B), 100 mM tetraethylammonium chloride (C) and 100 mM pentobarbital sodium salt (D) in the aqueous phase on the left