ISOLATION OF α -AMINOACYL ARGININES IN SCREENING OF AMINOPEPTIDASE B INHIBITORS

Sir:

Enzymes, such as aminopeptidase, phosphatase and esterase are located not only in cells but also on the membranes of various kinds of mammalian cells including lymphoid cells¹⁻³.

Inhibitors of these enzymes bind to cells and are thought to effect the immune system and to be potentially useful in the analysis of functions of cell membranes. In fact, bestatin^{4,5}, amastatin⁶, forphenicine⁷ and esterastin⁸, found by the screening for inhibitory, enhanced or suppressed immune responses.

In the screening for aminopeptidase B (E.C. 3.4.11.6.) inhibitors, we have recently isolated α -aminoacyl arginines from the culture filtrate of an actinomycetes (MF931-A2).

In this paper, we report the isolation and characterization of these dipeptides and their activity.

Aminopeptidase B was prepared from rat liver as described by Hopsu *et al.*⁹ and the activity was measured as reported previously⁴, and the concentration of the inhibitor required for 50% inhibition (IC_{50}) was determined. The culture medium used for the production of the inhibitors was as follows: galactose 2.0%, potato starch 4.0%, maltose 1.0%, K_2HPO_4 0.1%, polypeptone 0.1%, $MgSO_4 \cdot 7H_2O$ 0.05%, $CaCl_2$ 0.1% and NaCl 0.3% (pH 7.4 with 2 N NaOH before sterilization). Maximum production was attained on the 5~7th day of the shaking culture of the strain MF931-A2.

Extraction and purification processes are shown in Chart 1. After 6 days shaking of the cultures at 27°C, the active material in the culture filtrate was adsorbed on a column of Amberlite IRC-50 (H^+ form, 5 cm $\times$ 20 cm). The column was washed with 1 liter of distilled water and was eluted with 2 liters of 1 N aqueous ammonia. The active eluate was concentrated under reduced pressure, yielding 5.62 g of crude powder (IC_{50} = 31 μ g/ml). This was dissolved in 100 ml of distilled water, the solution adjusted to pH 7.0 with 1 N HCl and subjected to Dowex 50W X4 column chromatography (NH_4^+ form, 4 cm $\times$ 32 cm). The column was washed with 400 ml of distilled water and eluted by a linear gradient prepared from 600 ml of water and 600 ml 1 N ammonia. The eluates was cut into 10 g fractions. The inhibitory activity was found in fractions No. 65~80. The active fractions were combined and concentrated under reduced pressure, yielding 930 mg of crude material (IC_{50} = 7.3 μ g/ml). This was chromatographed on a column of Avicel (3 cm $\times$ 30 cm) with the solvent system *n*-butylacetate, *n*-butanol, acetic acid and water (2:4:1:1). The active fractions were combined and concentrated under reduced pressure to yield 595 mg of the crude material (IC_{50} = 7.1 μ g/ml) which was chromatographed on a column of Amberlite CG-50 type II (pyridinium form, 3 cm $\times$ 45 cm) previously equilibrated with a buffer consisting of 5% pyridine and 1% acetic acid (pH 5.8). This crude material was dissolved in 10 ml of the same buffer and applied to the column. The column was developed with 600 ml of this buffer, and then with 600 ml of the buffer consisting of 10% pyridine and 2.5%

Chart 1. Purification of aminopeptidase B inhibitors.

acetic acid (pH 5.8). Two peaks of activity were detected. The active fractions, which eluted between fraction No. 50 and No. 60 (each 10 g), were combined and adsorbed on a column of Dowex 50W X4 (NH_4^+ form, 1 cm $\times$ 10 cm). After the column was washed with 20 ml of distilled water, it was eluted with 20 ml of 1 N ammonia. The eluate was concentrated under reduced pressure and dissolved in 10 ml of distilled water. The solution was adjusted to pH 7.0 with 1 N acetic acid and lyophilized to give 40 mg of a white powder, the acetate ($\text{IC}_{50}=3.1 \mu\text{g/ml}$) of Compound I.

The second active fraction, eluted between fraction No. 70 and 80, was also treated as above and 80 mg of a white powder ($\text{IC}_{50}=1.6 \mu\text{g/ml}$) was obtained. It was dissolved with 4 ml of distilled water and chromatographed on Waters' semipreparative μ -bondapak C_{18} column with 15% Na_2SO_4 using Waters' Model ALC/GPC244 HPLC system. Two active peaks were detected. The active fraction which was eluted first, was adsorbed on a column of Dowex 50W X4 (NH_4^+ form, 1 cm $\times$ 5 cm). After the column was washed with 10 ml of water, it was eluted with 10 ml of 1 N ammonia. The eluate was concentrated under reduced pressure and dissolved with 5 ml of distilled water. The solution was adjusted to pH 7.0 in 1 N acetic acid, and lyophilized; 40 mg of white powder was obtained, the acetate ($\text{IC}_{50}=1.1 \mu\text{g/ml}$) of Compound II. The second active fraction was treated similarly and 16 mg of a white powder was obtained, the acetate ($\text{IC}_{50}=45 \mu\text{g/ml}$) of Compound III.

Acid hydrolysis of Compound I with 20% HCl at 105°C for 16 hours yielded two ninhydrin positive products. Amino acid analysis of the hydrolysate showed one mole each of valine and arginine. By the column chromatography of Dowex 50W X4 (pyridinium form, 2 cm $\times$ 50 cm) using 0.2 M pyridine-acetic acid buffer (pH 4.1) and 2.0 M pyridine-acetic acid buffer (pH 5.1), only these two amino acids were detected. The one eluted first was identified as L-valine and the second as L-arginine.

In the pmr spectrum of Compound I, the α -methine proton of the valine moiety, which was observed as doublet at 4.29 ppm in pH 2.0, was shifted to 3.96 ppm in pH 9.0. The result indicated that N-terminal amino acid of Compound I was valine and thus the structure of Compound I was suggested to be L-valyl-L-

arginine. This was confirmed by the comparison with synthetic L-valyl-L-arginine, which was synthesized from carbobenzoxy-L-valine and N $^{\epsilon}$ -nitro-L-arginine benzyl ester ditosylate by the DCC method.

Amino acid analysis of the hydrolysates of Compound II and III indicated that Compound II contained one mole each of isoleucine and arginine, and Compound III contained one mole each of leucine and arginine. These amino acids were purified and identified as the L-forms.

In the pmr spectra of Compound II and III, the shifts to higher field of the α -methine protons of the isoleucine moiety of Compound II (4.44 ppm at pH 2.0 to 4.06 ppm at pH 9.0) and the leucine moiety of Compound III (4.57 ppm at pH 2.0 to 3.95 ppm at pH 9.0) indicated that Compound II should be L-isoleucyl-L-arginine and Compound III L-leucyl-L-arginine. This was confirmed by comparison with available authentic samples of L-isoleucyl-L-arginine and L-leucyl-L-arginine (Bachem). Although Compound I, II and III are simple dipeptides, they had unexpectedly strong potency to inhibit aminopeptidase B as shown in Table 1. Moreover, the activity was specific to aminopeptidase B. They did not inhibit leucine aminopeptidase and aminopeptidase A. As has been reported bestatin inhibits aminopeptidase B and leucine aminopeptidase. Furthermore intraperitoneal injection of these compounds augmented delayed-type hypersensitivity (DTH) to sheep red blood cells in footpad test¹⁰⁾ using CDF₁ mice older than 10 weeks (Table 2).

Our results indicate that aminoacylarginines represent an important type of structures which are able to inhibit aminopeptidase B and to modify immune responses.

Table 1. Kinetic studies of aminopeptidase B inhibitors.

Compound	IC_{50} ($\times 10^{-6}$ M)	K_i ($\times 10^{-6}$ M)	Type of inhibition
I (Val-Arg)	10	2.1	Competitive
II (Ile-Arg)	3.2	0.5	Competitive
III (Leu-Arg)	150	23	Competitive
Bestatin	0.16	0.05	Competitive

$K_m=1.0 \times 10^{-4}$ M (L-arginyl- β -naphthylamide)

K_i value was calculated from DIXON plot and K_m value from LINEWEAVER-BURK plot.

Table 2. Effect of aminoacylarginines on DTH*.

Compound	Dose (μ g/mouse)	DTH (% enhancement)
I (Val-Arg)	100	47
	10	18
	1	36
	0.1	37
II (Ile-Arg)	100	5
	10	14
	1	42
	0.1	45
III (Leu-Arg)	100	0
	10	38
	1	38
	0.1	38

* DTH=Delayed type hypersensitivity.

Acknowledgements

This work was partly supported by a contract from the Division of Cancer Treatment, the National Cancer Institute, NO1-CM-57009, U.S.A.

KAZUMORI YAMAMOTO
HIROYUKI SUDA
MASAAKI ISHIZUKA
TOMIO TAKEUCHI
TAKAAKI AOYAGI
HAMAO UMEZAWA

Institute of Microbial Chemistry
14-23 Kamiosaki 3-Chome, Shinagawa-ku,
Tokyo 141, Japan

(Received September 2, 1980)

References

- 1) AOYAGI, T.; H. SUDA, M. NAGAI, K. OGAWA, J. SUZUKI, T. TAKEUCHI & H. UMEZAWA: Amino-peptidase activities of the surface of mammalian cells. *Biochim. Biophys. Acta* 452: 131~143, 1976
- 2) AOYAGI, T.; M. ISHIZUKA, T. TAKEUCHI & H. UMEZAWA: Enzyme inhibitors in relation to cancer therapy. *Jap. J. Antibiotics* 30 (Suppl.): S121~S132, 1977
- 3) AOYAGI, T. & H. UMEZAWA: Hydrolytic enzymes on the cellular surface and their inhibitors found in microorganisms. *FEBS Feder, Europ. Biochem. Soc.* 61: 89~99, 1980
- 4) UMEZAWA, H.; T. AOYAGI, H. SUDA, M. HAMADA & T. TAKEUCHI: Bestatin, an inhibitor of aminopeptidase B, produced by actinomycetes. *J. Antibiotics* 29: 97~99, 1976
- 5) SUDA, H.; T. TAKITA, T. AOYAGI & H. UMEZAWA: The structure of bestatin. *J. Antibiotics* 29: 100~101, 1976
- 6) AOYAGI, T.; H. TOBE, F. KOJIMA, M. HAMADA, T. TAKEUCHI & H. UMEZAWA: Amastatin, an inhibitor of aminopeptidase A, produced by actinomycetes. *J. Antibiotics* 31: 636~638, 1978
- 7) AOYAGI, T.; T. YAMAMOTO, K. KOJIRI, F. KOJIMA, M. HAMADA, T. TAKEUCHI & H. UMEZAWA: Forphenicine, an inhibitor of alkaline phosphatase, produced by actinomycetes. *J. Antibiotics* 31: 244~246, 1978
- 8) UMEZAWA, H.; T. AOYAGI, T. HAZATO, K. UOTANI, F. KOJIMA, M. HAMADA & T. TAKEUCHI: Esterastin, an inhibitor of esterase, produced by actinomycetes. *J. Antibiotics* 31: 639~641, 1978
- 9) HOPSU, V. K.; K. K. MÄKINEN & G. G. GLENNER: Purification of a mammalian peptidase selective for N-terminal arginine and lysine residues: Aminopeptidase B. *Arch. Biochem. Biophys.* 114: 557~566, 1966
- 10) ISHIZUKA, M.; T. MASUDA, N. KANBAYASHI, S. FUKASAWA, T. TAKEUCHI, T. AOYAGI & H. UMEZAWA: Effect of bestatin on mouse immune system and experimental murine tumors. *J. Antibiotics* 33: 642~652, 1980