NOTES

**Cochlioquinones and Epi-Cochlioquinones:
Antagonists of the Human Chemokine
Receptor CCR5 from *Bipolaris brizae*
and *Stachybotrys chartarum***

K. YOGANATHAN,* LAY-KIEN YANG, CHRISTINE ROSSANT,[†]
YICUN HUANG, SIEWBEE NG, MARK S. BUTLER
and ANTONY D. BUSS

MerLion Pharmaceuticals,
59A Science Park Drive, The Fleming,
Singapore Science Park, Singapore, 118240

(Received for publication October 9, 2003)

Chemokines are small proteins (7~16 kDa) that act at G protein-coupled receptors to regulate a variety of physiological and pathophysiological processes^{1,2)}. The human immunodeficiency virus Type 1 (HIV-1) principally uses CCR5 and CXCR4 chemokine receptors as co-receptors with CD4 to gain entry into target cells^{3,4)}. Therefore, a molecule that binds to the CCR5 receptor could potentially prevent HIV-1 entry into cells, which makes CCR5 an important target for anti-HIV-1 therapy⁵⁾.

In the course of screening our extract library for inhibitors of CCR5^{6,7)}, extracts from *Bipolaris brizae* (Nisikado) Shoem and *Stachybotrys chartarum* were found to displace macrophage inflammatory protein (MIP)-1 α from the human chemokine receptor CCR5. Bioassay directed fractionation led to the isolation of three active cochlioquinones from *Bipolaris brizae* (Nisikado) Shoem, namely, 17-methoxycochlioquinone A (**1**), cochlioquinone A (**2**)^{8~10)} and isocochlioquinone A (**3**)^{8~10)}. Two active epi-cochlioquinones compounds, 11-O-methyl-epi-cochlioquinone A (**4**) and epi-cochlioquinone A (**5**)¹¹⁾, were isolated from *Stachybotrys chartarum*. Herein, we report the isolation, structure elucidation and biological activity of **1**, **2**, **3**, **4** and **5**.

Materials and Methods

CCR5 Assay

CCR5 receptor binding activity was determined in a 96-well scintillation proximity assay¹²⁾ (SPA) using a [¹²⁵I]-human MIP-1 α and membranes prepared from CHO cells over expressing the human CCR5 receptor. The samples were dissolved in 12.5% aqueous DMSO and incubated with 12 μ g membranes, 0.17 nM [¹²⁵I]-MIP-1 α , 0.25 mg Wheat Germ Agglutinin-SPA beads in assay buffer (50 mM HEPES, 1 mM CaCl₂, 1 mM MgCl₂, 1% BSA and a protease inhibitor cocktail), shaking for 5 hours at room temperature. Radioactivity (total binding) was measured after a 2 hours bead settling period. Non-specific binding was defined in the presence of 1 μ M recombinant human MIP-1 α .

Results and Discussion

Fermentation

The producing microorganism was identified as *Bipolaris brizae* (Nisikado) Shoem based on morphological characteristics. The strain has been deposited in the MerLion Pharmaceuticals culture collection as F32659. The strain was sub-cultured on malt extract agar (CM057B, Oxoid) for 7 days at 24°C. It was used to inoculate 250 ml Erlenmeyer flasks each containing 50 ml of seed medium composed of 0.4% glucose, 1% malt extract and 0.4% yeast extract. The pH of the medium was adjusted to 5.5 prior to sterilization. The seed flasks were incubated for 5 days at 24°C on a rotary shaker at 200 rpm. A volume of 5 ml of seed culture was used to inoculate 50 ml of liquid medium in a 250 ml flask. The liquid medium is composed of 0.4% yeast extract, 2% glucose and 2% oatmeal. The pH was adjusted to 7.5 and autoclaved at 121°C for 30 minutes. The fermentation was carried out for 9 days at 24°C at 200 rpm.

The second fungal strains F32923 (*Stachybotrys chartarum*) has been deposited in the MerLion

* Corresponding author: yoga@merlionpharma.com

[†] Present address: Upstate Ltd., Cowley Rd., Cambridge CB4 0DS, UK.

Table 1. Physico-chemical properties of 17-methoxycochlioquinone A (**1**) and 11-*O*-methyl-*epi*-cochlioquinone A (**4**).

	1	4
appearance	Yellow oil	Yellow solid
Molecular formula	C ₃₁ H ₄₆ O ₉	C ₃₁ H ₄₆ O ₈
ESI-MS (M+Na) ⁺	585.3042(calcd 585.3040)	569.3071 (calcd 569.3090)
[α] _D ²⁵	+141° (c 0.20, CHCl ₃)	+38° (c 0.12, CHCl ₃)
UV (MeOH) λ _{max} (log ε) nm	283 (4.04), 386 (3.52)	270 (4.04), 385 (3.52)
IR ν _{max} (NaCl) cm ⁻¹	3521, 2968, 2879, 1730, 1660, 1645, 1456	3468, 2971, 2936, 1734, 1678, 1645, 1454

Pharmaceuticals culture collection. The strain was sub-cultured and fermented using the same conditions described above.

Isolation of Active Constituents of *Bipolaris brizae* (Nisikado) Shoem

The freeze-dried fermentation broth (2 liters) of *Bipolaris brizae* (Nisikado) Shoem was extracted 3 times with CH₂Cl₂-MeOH (1:1) and evaporated to dryness under vacuum. The dry extract (14.5 g) was partitioned 3 times between hexane and 90% MeOH in H₂O (1:1). The 90% MeOH portion was adjusted to 70% with H₂O, and then partitioned 3 times with CHCl₃. The active CHCl₃ fraction (1.7 g) was subjected to reverse phase preparative HPLC (isocratic elution for 70 minutes; mobile phase: 0.1% HCOOH in acetonitrile+0.1% HCOOH in H₂O (65:35); flow rate: 10 ml/minute) to give compounds **1** (6 mg), cochlioquinone A (**2**) (1 mg) and isocochlioquinone A (**3**) (3 mg). Cochlioquinone A (**2**) and isocochlioquinone A (**3**) were identified by comparison of their spectral properties with those reported in the literature^{8~10}.

Isolation of Active Constituents of *Stachybotrys chartarum*

The freeze-dried fermentation broth (2 liters) of *Stachybotrys chartarum* was extracted 3 times with CH₂Cl₂-CH₃OH (1:1) and evaporated to dryness under vacuum. The crude extract (2 g) was partitioned 3 times between hexane and 90% MeOH in H₂O (1:1). The active hexane fraction (0.5 g) was subjected to reverse phase preparative HPLC (gradient elution; mobile phase: 0.1% HCOOH in acetonitrile+0.1% HCOOH in H₂O (50:50→100:0 over 30 minutes); flow rate: 12 ml/minute) to give (**4**) (3 mg) and *epi*-cochlioquinone A (**5**) (1 mg). *Epi*-cochlioquinone A (**5**) was identified by comparison of its spectral properties with those previously reported¹¹.

Structure Elucidation

The molecular formula (C₃₁H₄₆O₉) of compound **1** was established by analysis of the ¹³C NMR, multiplicity-edited HSQC spectrum and positive HR-ESIMS. The physico-chemical properties of **1** are given in Table 1. The IR spectrum showed characteristic absorption bands from OH

Table 2. ^1H (500 MHz) and ^{13}C (125 MHz) NMR data for 17-methoxycochlioquinone A (**1**) in CDCl_3 .

No.	^{13}C (δ , m) ^a		^1H δ , (m, J in Hz)	COSY	HMBC ^1H to ^{13}C
1	39.5	t	1.42 m 2.45 m	1b, 2, 2b 1a, 2, 2b	3, 10
2	22.5	t	1.44 m 1.64 m	1a, 1b, 2b, 3 1b, 1b, 2, 3	1, 3, 24
3	86.0	d	3.23 dd (12, 2)	2a, 2b	5, 24, 25, 26
5	84.2	d	3.15 dd (12, 2)	6a, 6b	3, 6, 10, 27
6	26.2	t	1.54 m 1.72 m	5, 6b, 7a, 7b 5, 6a, 7a, 7b	5, 10
7	38.4	t	1.90 m 2.08 dt (11.5, 3)	6a, 6b, 7b 6a, 6b, 7a	5, 6, 9, 28
8	84.7	s			
9	52.8	d	1.69 d (10)	11	5, 10, 11, 12, 28
10	37.7	s			
11	64.1	d	4.89 d (10)	9	9, 10, 12, 13, 18
12	118.4	s			
13	152.8	s			
15	183.0	s			
16	132.8	s			
17	157.3	s			
17-OMe	62.8	q	4.01 s		17
18	185.5	s			
19	33.0	d	3.38 m	20, 29	15, 16, 17, 20, 29
20	78.1	d	5.25 br d (10)	19	16, 19, 21, 22, 20- OCOMe, 30
20-OCOMe	171.5	s			
20-OCOMe	22.0	q	1.88 s		OCOMe
21	36.8	d	1.69 m	30	22, 30
22	28.1	t	1.15 m 1.27 m	22b, 23 22a, 23	21, 30
23	12.9	q	0.91 t (7.5)	22a, 22b	21, 22
24	72.8	s			
25	27.0	q	1.18 s		3, 24, 26
26	24.7	q	1.16 s		3, 24, 25
27	13.3	q	1.00 s		5, 9, 10
28	21.9	q	1.31 s		7, 8, 9
29	16.2	q	1.19 d (7)	19	16, 19, 20
30	13.1	q	0.93 d (7)	21	20, 21, 22

^a Assignments based on COSY, multiplicity-edited HSQC and HMBC NMR data.

(broad, 3521 cm^{-1}), ester (1730 cm^{-1}) and carbonyl groups (1660 and 1645 cm^{-1}). The ^{13}C NMR spectrum of **1** (Table 2) showed 31 resolved peaks, which could be classified into seven methyl, one methoxyl, five methylene, three methines, four oxygenated methines and eleven quaternary carbons. The ^1H NMR spectrum of **1** (Table 2) displayed signals for four oxymethine groups (δ_{H} 5.25, br d; 4.89, d; 3.23, dd; 3.15, dd), five methylene groups (δ_{H} 1.4~2.5, m), one methoxy group (δ_{H} 4.01, s) and eight methyl groups (δ_{H} 0.91, t; 0.93, d; 1.00, s; 1.16, s; 1.18, s; 1.19, d; 1.31, s; and 1.88, s). These data were similar to that of cochlioquinone A (**2**)^{8~10} except for the presence of a methoxy group and the absence of aromatic methine H-17 (δ_{H} 6.56) present in cochlioquinone A (**2**)^{8~10}. The

placement of the methoxy at C-17 was confirmed by the observation of a 3J (OMe to C-17) correlation in HMBC NMR spectrum. This was also in accord with the observed carbon resonance of C-17 at δ_{C} 157.3 since it is α to both an oxygen and a carbonyl function. These observations supported the assignment of **1** as the 17-methoxy derivative of cochlioquinone A.

Examination of the NMR data of compound **4** (Table 3) suggested that **4** was a cochlioquinone derivative related to *epi*-cochlioquinone A (**5**). The physico-chemical properties of **4** are given in Table 1. A molecular formula of $\text{C}_{31}\text{H}_{46}\text{O}_8$ was determined for compound **4** on the basis of positive HR-ESIMS and ^{13}C NMR data, which was different to **5** by a CH_2 unit. Comparison of ^1H and ^{13}C NMR data of **4** with

Table 3. ^1H (500 MHz) and ^{13}C (125 MHz) NMR data for 11-*O*-methyl-*epi*-cochlioquinone A (**4**) in CDCl_3 .

No.	^{13}C (δ , m) ^a	^1H δ (m, <i>J</i> in Hz)	COSY	gHMBC ^1H to ^{13}C
1	37.7	t	1.35 m	1b, 2a, 2b
			2.05 m	1a, 2a, 2b
2	22.7	t	1.48 m	1a, 1b, 2b, 3
			1.62m	1a, 1b, 2a, 3
3	85.2	d	3.15 dd (12, 2)	2a, 2b
5	84.2	d	3.08 dd (12, 2)	6a, 6b
6	24.5	t	1.57 m	5, 6b, 7a
			1.80 m	5, 6a
7	37.5	t	1.72 m	6a, 7b
			2.35 m	7a
8	80.1	s		
9	53.4	d	1.60 d (9)	11
10	35.5	s		
11	68.4	d	4.08 d (9)	9
11-OMe	57.9	q	3.40 s	
12	119.0	s		
13	153.3	s		
15	182.2	s		
16	150.0	s		
17	135.0	d	6.55 s	
18	187.0	s		
19	35.5	d	3.20 m	20, 29
20	79.5	d	5.02 dd (8, 4)	19, 21
20-OCOMe	171.4	s		
20-OCOMe	21.7	q	1.93 s	
21	37.0	d	1.60 m	30
22	27.5	t	1.10 m	22b, 23
			1.27 m	22a, 23
23	12.7	q	0.89 t (7)	22a, 22b
24	72.8	s		
25	24.6	q	1.16 s	
26	27.0	q	1.18 s	
27	12.7	q	0.60 s	
28	27.3	q	1.31 s	
29	17.8	q	1.17 d (7)	19
30	14.0	q	0.91 d (7)	21

^a Assignments based on COSY, HSQCED and HMBC.

those of *epi*-cochlioquinone A (**5**)¹¹ revealed that **4** had a methyl ether group (δ_{H} 3.40, s; δ_{C} 57.9) in addition to the *epi*-cochlioquinone A (**5**) skeleton. The observation of an HMBC correlation from the methyl ether protons to δ_{C} 68.4 allowed placement of the methyl ether group at C-11 and assignment of the structure of **4** as the 11-*O*-methyl derivative of *epi*-cochlioquinone A. The relative stereochemistry of H-11 was deduced to be α from the $J_{9\alpha-11}$ value of 9 Hz, which required H-11 and H-9 α to be in a *cis*-diaxial arrangement.

Biological Activities

17-Methoxycochlioquinone A (**1**), cochlioquinone A (**2**), isocochlioquinone A (**3**), 11-*O*-methyl-*epi*-cochlioquinone

A (**4**) and *epi*-cochlioquinone A (**5**) were found to compete effectively with MIP-1 α for binding to human CCR5 with IC_{50} values of 100, 11, 50, 7 and 4 μM respectively. Comparison of the activity of **2** and **3** suggested that the quinoid skeleton increased the activity of the cochlioquinones, while substitution of a methoxy group at C-17 in **1** significantly decreased the activity compared to **2**. The IC_{50} values **4** and **5** were similar to that of cochlioquinone A (**2**) (11 μM), which has the same 12,13,15-trisubstituted quinone unit but different stereochemistry at C-8. Cochlioquinones family have been previously reported to inhibit NADH-ubiquinone reductase¹⁰ and acyl-CoA cholesterol acyltransferase¹¹.

Acknowledgments

We would like to acknowledge A/Prof. LAI from the Department of Chemistry, National University of Singapore, for obtaining IR spectra and A/Prof. LOH for the use of the polarimeter. The financial support of GlaxoSmithKline, the Economic Development Board of Singapore and Institute of Molecular and Cell Biology are gratefully acknowledged.

References

- 1) KEDZIERSKA, K.; S. M. CROWE, S. TURVILLE & A. L. CUNNINGHAM: The influence of cytokines, chemokines and their receptors on HIV-1 replication in monocytes and macrophages. *Rev. Med. Virology* 13: 39~56, 2003
- 2) LOCATI, M. & P. M. MURPHY: Chemokines and chemokine receptors: biology and clinical relevance in inflammation and AIDS. *Ann. Rev. Med.* 50: 425~440, 1999
- 3) LEHNER, T.: The role of CCR5 chemokine ligands and antibodies to CCR5 coreceptors in preventing HIV infection. *Trends in Immunology* 23: 347~351, 2002
- 4) KAZMIERSKI, W. M.; L. BOONE, W. LAWRENCE, C. WATSON & T. KENAKIN: CCR5 chemokine receptors: gatekeepers of HIV-1 infection. *Current Drug Targets: Infectious Disorders* 2: 265~278, 2002
- 5) DE CLERCQ, E.: New anti-HIV agents and targets. *Med. Res. Rev.* 22: 531~565, 2002
- 6) YOGANATHAN, K.; C. ROSSANT, S. NG, Y. HUANG, M. S. BUTLER & A. D. BUSS: 10-Methoxydihydrofusicin, fusicarin and fusicin, novel antagonists of the human CCR5 receptor from *Oidiodendron griseum*. *J. Nat. Prod.* 66: 1116~1117, 2003
- 7) CAO, S.; C. ROSSANT, S. NG, A. D. BUSS & M. S. BUTLER: Phenolic derivatives from *Wigandia urens* active against the chemokine receptor CCR5. *Phytochemistry* 64: 987~990, 2003
- 8) LIM, C.-H.; H. MIYAGAWA, M. AKAMATSU, Y. NAKAGAWA & T. UENO: Structures and biological activities of phytotoxins produced by the plant pathogenic fungus *Bipolaris cynodontis* cynA. *J. Pesticide Sci.* 23: 281~288, 1998
- 9) MIYAGAWA, H.; S. NAGAI, T. TSURUSHIMA, M. SATO, T. UENO & H. FUKAMI: Phytotoxins produced by the plant pathogenic fungus *Bipolaris bicolor* El-1. *Biosci. Biotech. Biochem.* 58: 1143~1145, 1994
- 10) LIM, C.-H.; H. UENO, H. MIYOSHI, H. MIYAGAWA, H. IWAMURA & T. UENO: Phytotoxic compounds cochlioquinones are inhibitors of mitochondrial NADH-ubiquinone reductase. *J. Pesticide Sci.* 21: 213~215, 1996
- 11) FUJIOKA, T.; K. YAO, K. HAMANO, T. HOSOYA, T. KAGASAKI, Y. FURUKAWA, H. HARUYAMA, S. SATO, T. KOGA & Y. TSUJITA: *Epi*-cochlioquinone A, a novel acyl-CoA: Cholesterol acyltransferase inhibitor produced by *Stachybotrys bisbyi*. *J. Antibiotics* 49: 409~413, 1996
- 12) COOK, N. D.: *Drug Discovery Today* 1: 287~295, 1996