VI. 浜北市根堅遺跡出土人骨の弗素含有量

田 辺 義 一

静岡県浜北市根堅の石灰岩採掘場より出土した数個の人骨破片の相対的地質年代を推測する目的で、それらの人骨に含まれている弗素の量を測定し、同遺跡出土の獣骨についてしらべた弗素含有量と比較考察した。また、さきに測定した静岡県三ケ日、愛知県牛川などの人骨、獣骨の弗素含有量（田辺, 1962）との比較考察を行なった。

分析資料及び分析結果

人骨は数次にわたって採集されたものを、その都度鈴木教授より提供していただいたもので、最初は上腕骨と鎖骨を同遺跡の獣骨片3個とともに1961年の秋に分析した。その後前頭骨、頭頂骨の破片、また別に脛骨及び人骨とみられる大腿骨が発見され、それぞれ分析に供されたが、対照とするのに適当な獣骨が得られなかつたので、このことにつき鈴木教授に依頼しておいたところ、同教授の連絡により、当時この遺跡一帯を独自に調査しておられた国立科学博物館の長谷川善和氏より根堅遺跡出土のトラとシカの骨10数個が分析用に提供された。長谷川氏によると、これらの獣骨は2つの別の層から採集されたもので、1つの層はトラとシカを含み（A地点、第I-5図）、も1つの層はシカのみでトラを含まない（D地点、第I-5図）。同氏はこの層をそれぞれ *Felis Bed*, *Homo Bed* と呼んでおられ、ともに洪積世後期に属し、前者は後者より古いものと考えておられた。そこでこれらの獣骨を本遺跡の *pilot series* とすることにし、1964年春、人骨とともにこれらの獣骨の分析を行なった。更に1965年の暮れに人の腰骨1個を分析した。これらの人骨のうち頭骨片は鈴木教授のしらべにより、すべて同一個体に属するものであることがわかつた。

分析には骨の緻密質からドリルで 50 mg 余を採取したものをを用いた。分析方法はさきに三ケ日人骨などについて行なったのと同じである。

分析結果は一括表示した（第VI-1表）。このうち獣骨A及びBとしたものは長谷川氏の提供になるもので、それぞれ同氏の *Homo Bed* 及び *Felis Bed* の獣骨にあたるものである。また獣骨Cとしたものは1961年に分析した獣骨である。また人骨はその後の地質学的検討により、脛骨（と大腿骨）は *Felis Bed* に属し、そのほかはすべて *Homo Bed* に属するものと結論されたといわれるので2つの群に分けて記載した。

考 察

(1) 浜北人骨と獣骨の比較

分析結果によると、*pilot series* とした獣骨のA（*Homo Bed*）とB（*Felis Bed*）とでは弗素の量においてかなりはつきりしたちがいがみとめられ、同じシカの骨でもBに属するものはAのものより多量の弗素を含んでいる。Aの弗素量は0.33~0.43%にわたり、Bのそ

本研究の一部は日本人類学会・日本民族学会第19回連合大会（京都、1964）で報告した。

Table VI-1. Fluorine contents of bones from Hamakita

Animal		F (%)	Loss of Ignition (%)	Man		F (%)
A Cervus	femur	0.33	12.63	A	humerus	0.64
	cannon bone 1	0.43	10.76		clavicula	0.55
	" " 2	0.35	12.28		iliac bone	0.50
	finger bone	0.33	12.31		frontal bone	0.41
	$\bar{x}$	0.36	12.00		parietal bone	0.39
	u^2	0.0023	0.7030		$\bar{x}$	0.50
B Felis	humerus	0.83	10.57	B		0.0106
	femur 1	0.72	9.48		tibia	0.77
	" 2	0.87	11.50		(femur	0.69)
	cannon bone	0.91	8.17		$\bar{x}$	0.73
	tibia	0.80	10.57		u^2	0.0032
	finger bone	0.76	10.39			
	mandible	0.90	8.11			
	dentin (C)	0.82	—			
	Cervus cannon bone	0.88	7.78			
	" finger bone	0.90	9.84			
	$\bar{x}$	0.84	9.62			
	u^2	0.0053	1.7211			
C		0.36				
		0.34				
		0.38				
	$\bar{x}$	0.36				
	u^2	0.0004				

れは0.72~0.91%で、全く重複するところがなく、それぞれの平均値0.36%と0.84%とは1%の危険率をもつて明らかな有意差がある。またそれらの灼熱減量も平均値において1%の危険率でAの方がBよりも多いと云える。従つて獣骨AがBより地質年代においてより新しいことはほぼ確実である。

獣骨CはAと殆ど同じ弗素量を示し、Aと同一地層である可能性がよい。

次に人骨の弗素量についてみてゆくと、全体的にみると獣骨AとBの間にあるようにみえる(弗素量平均0.56%, 分散 u^2 0.0204)。しかしこれを地質学的結論に従つてA群、B群に分けたものについてみると、A群は0.39%~0.64%にわたり、獣骨Aと一部重複し、B群は0.69%~0.77%で獣骨Bと一部重複する。そこで人骨A、B群及び獣骨A、B相互間の弗素量の平均値の差の有意性をしらべてみると、まず人骨AとBとでは5%の危険率で有意差がみとめられる。次に獣骨との比較では、人骨Bは獣骨Aとは1%の危険率で有意差があるが、獣骨Bとは5%の危険率においても有意差があるとはいえないという結果になる。このことから人骨Bの弗素量は獣骨Bのひろがりの中に完全には入らないが、平均的にみれば、人骨Bは獣骨Bとはゞ同一、あるいはこれに近い地質年代に属すると見做して大体矛盾はないと考えられる。一方人骨Aは獣骨Bとは1%の危険率で有意差があり、また獣骨Aとも5%の危険率で有意差がみとめられる。したがって人骨Aは獣骨A、Bの間のAにより近い位置にあ

るようである。ところでこの人骨Aとしたもののうち頭骨は比較的弗素量が少く獣骨Aと重複するので、人骨Aを弗素量にもとずいて頭骨(A-a)とその他の部分(A-b)とに一応分けてみることも考えられる。そしてそのように分けて検定してみると、それぞれのちがいは漸進的であり、人骨A-a(弗素量平均0.40%)は獣骨B、人骨Bと1%の危険率で、また人骨A-bとも5%の危険率で有意差があるが、獣骨Aとは有意差があるとはいえない。また人骨A-b(平均0.56%)は獣骨BともAとも1%の危険率で有意差があるが、人骨Bとは有意差があるとはいえないという微妙な傾向を示す。しかしこのような細分は人為的であるのでこゝではこのような見方もできるということを示すにとどめる。そこで人骨の弗素量について総括すると、

- 1) 人骨Bは獣骨Bに近く、これと同一地質年代と見做すことに矛盾はないと考えられる。
- 2) 人骨Aは一部獣骨Aと重複するが、平均値においては獣骨Aと若干ちがいがみられ、これと同一地層であると断定することはできない。しかしBとくらべれば獣骨Aにずっと近いと考えることができる。

簡単に式で示すと： 獣骨B $\geq$ 人骨B > 人骨A > 獣骨A

(2) 浜北人骨と他の遺跡の人骨との比較

次に浜北人骨の弗素量を、さきにしらべた三ケ日や牛川などで出土した骨のそれと対比してその相対的年代関係を考察する。そのために浜北、牛川、三ケ日の人骨、獣骨、及び嵩山の獣骨の各群の弗素量の平均値についてその差の有意性を検定した。第VI-1図はその結果を図示したものである。ここで浜北の人骨はA、B群に、また獣骨もA、B両群に分けてある。

	F (%)	U Man	U Animal	H Animal B	M Animal	H Man B	M Man	H Man A	H Animal A	S Animal
U Man	0.93		—	—	—	—	>>	>>	>>	>>
U Animal	0.91			—	—	—	>	>>	>>	>>
H Animal B	0.84				—	—	>>	>>	>>	>>
M Animal	0.81					—	—	>	>>	>>
H Man B	0.73						—	>	>>	>>
M Man	0.64							>	>>	>>
H Man A	0.50								>	>>
H Animal A	0.36									>
S Animal	0.23									

Fig. VI-1. Comparison of fluorine contents of groups of bones from various sites

—: not significant

>: significant at 5% level

>>: significant at 1% level

U: Ushikawa, M: Mikkabi, H: Hamakita, S: Suse

この結果によると、浜北獣骨Bは牛川の人骨及び獣骨、三ケ日獣骨と有意差があるとはいえないが、三ケ日人骨、嵩山獣骨とは有意差がある。浜北獣骨Aは他のすべてとの間に有意

差がみとめられる。一方浜北人骨Bは牛川人骨及び獣骨、三ケ日獣骨及び人骨と有意差があるとはいえないが、嵩山獣骨とは有意差がある。浜北人骨Aは他のすべてとの間に有意差がみとめられる。

これによつて弗素量の近いものをあつめて群化してみると、〔牛川人骨及び獣骨、浜北獣骨B及び人骨B、三ケ日獣骨〕がそれぞれ弗素量にいくらかずつ差異をもちながら1つの群をなし、〔浜北人骨B、三ケ日獣骨及び人骨〕が別の群をなし、〔浜北人骨A〕、〔浜北獣骨A〕及び〔嵩山獣骨〕はそれぞれ単独の群をなしているようである。上のうち三ケ日獣骨と浜北人骨Bは2つの群にまたがっているが、前者は弗素量にかなり変異があり、分散の大きいことが原因していると思われ、後者は両群の境界的位置にあるためと思われる。また最後の3群も1%の危険率を基準にすれば隣り合つた同志は比較的近い関係にある。

異つた遺跡から出土した骨の弗素量を比較する場合、その地域差を十分考慮に入れる必要があり、弗素量のちがいを直ちに年代のちがいにむすびつけることは危険であるが、ここにとりあげた4遺跡は東海地方に比較的近接して存在し、地質条件も似かよつていたので、弗素量のちがいがかなりの程度年代のちがいをあらわすものと見做すことができるであろう。それで一応上のような大別がなされることは浜北人骨の相対的年代位置を決めるのに役立つものであろう。そしてこれらのうち同じ浜北遺跡で獣骨A、Bという、地質学的にも、弗素量においても明らかなちがいのある2つの群の存することは、この2群が浜北だけでなく、その近隣を含んだ地域をしらべる場合の1基準となりうるものと考えられる。そして浜北獣骨A、B両群の間に三ケ日人骨によつて代表されるような1つの群が存在する可能性も推測され、また三ケ日の獣骨も本来は2つ位の群からなつていてのではないかということも考えさせられる。

文 献

田辺義一、1962：三ケ日遺跡出土人骨の弗素含有量。人類誌，70，41-45。

（お茶の水女子大学家政学部）

Hamakita Man and the Site of Nekata Limestone Quarry at Hamakita

I. Site of Nekata

Hisashi SUZUKI¹⁾ and Banri ENDO²⁾

As compared with the remarkable progress of the study of the pre-ceramic culture, the knowledge of Pleistocene man in Japan has been deficient. This may be explained by the fact that most of the pre-ceramic sites so far discovered are open-sites in the loam, which are not suitable for preservation of skeletal remains, and that most of skeletal remains of Pleistocene man, as were found at the sites of Ushikawa, Mikkabi and Kuzuu, are mere fragments occurring in the deposits in the limestone fissures, having been transported and laid by water. The site of Nekata, where the Hamakita man was found, is one of such fissure deposits.

The site of the Nekata limestone quarry (Maruhon Co.) at Hamakita is located 16 km. north-northeast of Hamamatsu City, Central Japan, and 22 km. away from the Pacific Coast (Fig. I-1). The site is situated on the southern border of hills sloping down to a plane which is cultivated into rice field (Figs. I-2 and I-11). The quarry is working to Palaeozoic limestone having caves and fissures which are filled with fossiliferous Pleistocene deposits. Plans of the quarry in which the site is located are shown in Figs. I-3~5 and its birdseye-view in Fig. I-12.

The present paper deals with the result of the surveys and excavations of the site performed between 1960 and 1963. In the course of five surveys carried out between 1960 and 1962, several fragments of human bone were obtained with many mammalian fossils as follows (cf. Figs. I-3~5): skull, hip bone and ulna at Locality C; humerus and clavicle at Locality D; tibia at Locality B; Lower molar at Locality D. From June to August, 1962, excavation was made in the Main Cave (Areas I, II and III in Fig. I-5) at the site, for the purpose of obtaining mammalian fossils as many as possible and clarifying the situation of the Pleistocene deposits so as to determine the geologic age of the site (Figs. I-8 and I-9). In June, 1963, excavation was made at another cave (Area IV in Fig. I-4) for the same purpose.

*¹⁾Department of Anthropology, Faculty of Science,
University of Tokyo; ²⁾Department of Anthropology,
The University Museum, University of Tokyo*

II. Skeletal Remains of Hamakita Man

Hisashi SUZUKI

Human Skeletal remains were found in the deposits of the cavelike fissure

(Main Cave in Chaps. I and II) at the site of the Nekata limestone quarry at Hamakita between 1960 and 1962. The deposits can be divided into the upper chocolate-coloured clay layer and the lower gravel-bearing brownish one. From the upper layer were found fragments of human skull vault, clavicle, humerus, ulna and ilium and a lower third molar, whereas the lower layer yielded one fragment of tibia. In colour, condition of preservation, specific gravity and fluorine content (Table II-1) the bones from the upper layer are different from that of the lower layer; the latter is considered to be of a somewhat older geologic age than the former.

The bones from the upper layer may belong to an individual about twenty years old, but no estimation can be made on the bone from the lower layer.

The fragment of skull (Fig. II-1 and Pl. II-1) is a part of calvaria belonging to that of *Homo sapiens*. It is small, the size being about the same as that of the pygmy. The bone is relatively thick (Table II-2). The form of skull seems to be brachycephalic or close to it.

The zygomatic bone (Pl. II-2, A, B), the only facial bone of the remains, is extremely small and the swelling on its anterior surface is weak. The incisura inframalaris and the fossa canina of the maxilla are less developed, but the tuberosity at the origin of the masseter on the lower surface of the body of the malar bone is well developed and comparable with that of Jomon-Age man. The lower third molar shows a reduced form.

As to the bones of the extremities, the restored lengths of the clavicle, humerus and ulna are apparently short, consequently the stature must be small and seems to belong to the category of the pygmy. On the other hand, the bones are relatively stout.

Other morphological characteristics of each bone are summarized as follows.

The clavicle (Pl. II-3, A, B) shows a vertically compressed form. Its sagittal breadth at the middle of the shaft is not much different from that at the acromial end. Its Claviculo-Humeral Index (Table II-4) falls within the range of the variation of the Mongolian races such as the Japanese.

The shape of the humerus (Pl. II-4-1, A, B, C) differs from that of the Ushikawa man and belongs to that of *Homo sapiens*. The general shape is closest to that of the Jomon-Age man, but the lower part of the shaft of Hamakita humerus is almost cylindrical (Table II-5, 6).

The ulna (Pl. II-3-2, A, C) is different from any others hitherto unearthed in Japan in showing no crista-like development along the interosseal margin and having the narrow anterior surface (Fig. II-2, Table II-7).

The human bone from the lower layer is only one fragment of the right tibia (Pl. II-4-B, A~C). The form of its cross-section belongs to the eurycnemia. The anterior crest of the bone is obtuse, and its medial surface is remarkably swollen by which the bone is distinguished from the tibia of the Jomon-Age man.

From the facts described above, it is clear that the Upper-Layer man of Hamakita belongs to a human being different from the Ushikawa man and to *Homo sapiens*; his characteristics are similar to those of the Jomon Age man on one hand, and on the other hand they resemble the European Neanderthal man in some points. Moreover, the stature of the Hamakita Man seems to have been very low as is true in other fossil man in Japan (Mikkabi, Ushikawa

Man etc.)

It is impossible to clarify the characteristics of the Lower-Layer man owing to small fragment.

From the points mentioned above it is considered that the Hamakita Man from the upper layer may have very closely resembled to the Mikkabi man and is a probable ancestor of the Jomon-Age man.

Provided that the Pleistocene man in Japan can be regarded as the ancestor of the Jomon-Age man, the remarkable increase in the stature of the latter (Table II-8) may be ascribed to the changes in some of the living conditions, as is the case with the Bushman (TOBIAS, 1962) whose stature has remarkably increased in the last few decades.

*Department of Anthropology, Faculty of
Science, University of Tokyo*

III. The Deposits at the Nekata Cave-site, Hamakita City, Shizuoka and their Geologic Age

Kiyotaka CHINZEI

The fossil human bones were found in the deposits filling up a cave opened in the Palaeozoic limestone lens at the Maruhon limestone quarry. At this site, two caves were excavated for scientific investigation. The main cave is a northnortheasterly trending narrow cave about 20 m long (areas I-III of Fig. I-4), and the other is a smaller cave (area IV of Fig. I-4) having an elliptical profile, about 2 m in longer diameter. The human bones were found in the main cave. The longitudinal vertical section of this cave is shown in Fig. III-2. The cave opens probably at its western end.

The deposits filling up the main cave are divided into the Pleistocene Gansuiji formation (HASEGAWA, 1964) and the overlying younger sediments. The stratigraphic succession is shown in Table III-1, and the cross-sections in Figs. III-3~10.

The lower part, about 1.6 m thick, of the Gansuiji formation is composed of yellowish brown clay, sandy silt and laminated medium-grained sand. The lower part grades upward into gravel-bearing brownish porous clay containing abundant mammalian bones. The brownish porous clay is interfingering with olive-coloured massive silt and limestone rubble beds. They are covered by porous, snail-bearing brown clay in the western part of the cave, and by chocolate-coloured greasy clay in the eastern part, which constitute the uppermost of the formation. A slight stratigraphic discontinuity can be expected between the main part and the uppermost part. A reddish brown soil, 50 cm thick, is found in the overlying younger sediments.

Most of the human bones in question are concluded to have occurred in the chocolate-coloured clay, the uppermost part of the Gansuiji formation, and a fragment of tibia seems to have occurred in the gravel bearing brown clay of the middle part of the formation.

The Quaternary terraces developed in the area surrounding the site are shown in Fig. III-11 and in Tab. III-2.

The main cave opens on the slope of the Mikataga-hara terrace. The cave deposits are probably the secondary deposits derived from the gravel beds of the terrace, judging from their constituent materials and their geographic position (Fig. III-12). The bottom of the cave (76 m above sea level) is nearly at the same level as the top of the gravel beds (75 m) of the Ringyo-shikenjo terrace 1.5 km east of the cave. Since any caves or fissures occurring near the main cave do not have their bottom lower than the level of the gravel beds, it is most likely that the main cave has been formed close to the groundwater table at the time of the Ringyo-shikenjo gravel deposition. Thus, deposition of the Gansuiji formation is considered nearly contemporaneous with, or later than, the Ringyo-shikenjo gravel beds. On the other hand, the reddish soil beds similar to the reddish brown soil overlying the Gansuiji formation are found on the top of the gravel beds older than the Hamakita terrace. This means that the deposition of the uppermost part of the formation, the *Homo* bed, was earlier than the Hamakita terrace.

There are two opinions on the correlation of the Quaternary terraces in the Hamamatsu area with the Kanto region which is the standard area of the Japanese Quaternary chronology. According to KOBAYASHI (1963, 65) the Hamamatsu formation is correlative with the Shimosueyoshi formation of the Kanto region, while TSUCHI (1960) referred the Mikataga-hara gravels to the Shimosueyoshi formation. It has been generally accepted that the Shimosueyoshi formation is the deposits at the time of the last high sea level in the Pleistocene, i. e., the Riss-Würm interglacial. In both opinions, the Ringyo-shikenjo terrace is younger than the Shimosueyoshi. Thus, the Gansuiji formation is concluded to be the deposits of the Würm glacial age. In the present stage of our knowledge, more detailed correlation with the European standard should not be attempted. KOBAYASHI (1963) made a tentative correlation of the Quaternary succession of the Hamamatsu area to the European standard (Tab. III-3). According to him, the Ringyo-shikenjo terrace is in the middle of the Hauptwürm.

*Geological Institute, Faculty of Science,
University of Tokyo*

IV. Horizons yielding Human Bones

Yoshikazu HASEGAWA

The human bones discovered in Gansuiji formation comprise seven pieces; skull, clavicle, humerus, ulna, ilium, lower third molar and tibia. The horizon and location of the skull and the ilium were confirmed by the present writer immediately after the excavation. As to the molar, its horizon and location have been ascertained later, on the basis of the information from the person who excavated it. These specimens are supposed to have been contained in a horizon corresponding to, or a somewhat older than, the Shell bed which rests on the Felis bed. The horizon of the clavicle, humerus and ulna has not been

confirmed yet, but the writer thinks it is the same as the above-mentioned horizon. These six bones are the ones described as *Homo* sp. A (human bone A by TANABE, and upper layer bones by SUZUKI) in the *Homo* bed.

The horizon of the last specimen, tibia, was questionable. However, it has been concluded that the specimen is older than the other bones and occurred in the *Felis* bed, judging from the fluorine content and the condition of fossilization, as well as from the circumstances at the time of the excavation as informed by the excavator. This is the specimen described as *Homo* sp. B (human bone B by TANABE, and lower layer bone by SUZUKI).

Thus, the writer thinks that the human bones excavated from the Gansuiji formation were contained in two different horizons.

*Department of Paleontology, National
Science Museum, Tokyo*

V. Vertebrate Fossils from the Gansuiji Formation

Fuyuji TAKAI¹⁾ and Yoshikazu HASEGAWA²⁾

The fossil fauna in the Gansuiji formation resembles in composition that of the Upper Kuzuuian formation and the Tadaki formation of the late Pleistocene, and seems to be contemporaneous with the latter. This fauna comprises three horizons, *Pices* bed, *Felis* & *Macaca* bed and *Homo* or *Shell* bed, in ascending order. The constituent species are shown in Tab. V-1.

¹⁾*Geological Institute, Faculty of Science, University
of Tokyo*; ²⁾*Department of Paleontology, National
Science Museum, Tokyo*

VI. Fluorine Contents of Human Bones from Pleistocene Deposits of Hamakita

Giichi TANABE

The fluorine dating method was applied to the human bones found at the Nekata site, Hamakita city, Shizuoka Prefecture, Japan. The bones, divided geologically into two groups, A and B, were compared with the pilot series consisting of two groups of animal bones A and B, excavated from different layers of the same site. From the result of analysis (Table VI-1) it was concluded that animal bone A was quite significantly newer than B, and human bone B was of almost the same geological age as animal B, and human bone A was rather close to animal A.

The fluorine content of the Hamakita bones was compared with that of the bones from neighbouring sites (TANABE, 1962), (Fig. VI-1).

Ochanomizu Women's University, Tokyo