Review Article

Cardiovascular Disease in Japanese Patients with Type 2 Diabetes Mellitus

Kazuya Fujihara, MD, PhD and Hirohito Sone, MD, PhD, FACP

Individuals with diabetes have a two- to four-fold increased risk of coronary artery disease (CAD) and higher mortality rates than those without diabetes. Because not only microvascular but also macrovascular disease in patients with diabetes are known to predispose patients to a lower quality of life as well as lead to higher mortality rates, identifying and managing risk factors of CAD is of clinical relevance in diabetes care. A number of antihyperglycemic drugs are currently approved for the treatment of hyperglycemia in patients with type 2 diabetes mellitus (T2DM), with several new drugs having been developed during the last decade. Diabetes-related complications have been substantially reduced worldwide. However, in view of the current situation in which both the prevalence of obesity and glucose abnormality have increased worldwide, including Japan, diet and exercise remain the crucial means of treatment for patients with diabetes. Furthermore, predicting the development of CAD is essential. This review summarizes data from recent studies on cardiovascular disease in patients with T2DM, focusing on clinical trials and big data, including studies involving Japanese individuals.

Keywords: diabetes, cardiovascular disease, coronary artery disease, stroke, antihyperglycemic drugs

Introduction

Coronary artery disease (CAD) is a major cause of morbidity and mortality in patients with type 2 diabetes mellitus (T2DM).¹⁾ Although comprehensive and intensive

management of multiple cardiovascular risk factors in patients with T2DM is recommended to reduce the risk of cardiovascular events, a considerable number of patients develop CAD even under intensive management.²⁾ A number of antihyperglycemic drugs are currently approved for the treatment of hyperglycemia in patients with T2DM, and diabetes-related complications have been substantially reduced worldwide.³⁾ However, in view of the current situation in which both the prevalence of obesity and glucose abnormality have increased worldwide, including Japan, diet and exercise remain the crucial means of treatment for patients with diabetes. This review summarizes data from recent epidemiological and clinical studies on cardiovascular disease (CVD) in patients with T2DM, focusing on clinical trials and big data, including studies on Japanese patients.

Characteristics of CVD in Patients with T2DM

CAD during the early stages of glucose abnormalities and in individuals with diabetes was observed to be associated with worsening of the mortality rate following acute myocardial infarction.^{2,4,5)} The incidence of CAD among people with diabetes in Japan is lower than that in the US and other western countries as well as among the general population^{6,7)} (Table 1). Hotta et al. showed that the frequency of CVD as a cause of death was 24.0% and 31.6% in men and women, respectively, between 1991 and 2000.⁸⁾ A more recent study conducted between 2001 and 2010 showed that the frequency of CVD as a cause of death decreased; the frequencies reported were 10.7% and 12.6% in men and women, respectively.⁹⁾

Patients with diabetes without a previous myocardial infarction have a risk of myocardial infarction as high as that in patients without diabetes with a previous myocardial infarction.¹⁰⁾ Thus, the presence of diabetes and underlying systemic insulin resistance and metabolic abnormalities due to diabetes may accelerate the development of CAD. Impaired fasting glucose and glucose tolerance are also associated with modest increases in the risk of CVD.^{11–16)} Booth et al. revealed that diabetes confers a risk of CAD equivalent to aging 15 years in western

Department of Internal Medicine, Niigata University Faculty of Medicine, Niigata, Niigata, Japan

Received: November 1, 2017; Accepted: November 15, 2017
Corresponding author: Hirohito Sone, MD, PhD, FACP. Department of Internal Medicine, Niigata University Faculty of Medicine, 1-754 Asahimachi, Niigata, Niigata 951-8510, Japan
Tel: +81-25-368-9026, Fax: +81-29-368-9300
E-mail: sone@med.niigata-u.ac.jp

©2018 The Editorial Committee of Annals of Vascular Diseases. This article is distributed under the terms of the Creative Commons Attribution License, which permits use, distribution, and reproduction in any medium, provided the credit of the original work, a link to the license, and indication of any change are properly given, and the original work is not used for commercial purposes. Remixed or transformed contributions must be distributed under the same license as the original.

Table 1 Summary of the association between the incidence of or death from cardiovascular disease and glucose abnormalities

Study	Country	Glucose tolerance status	Age (years), mean or range	BMI (kg/m ²), mean	% Men	No. participants total	Covariate	Endpoint	Hazard ratios
Funagata Study Tominaga et al. (1999)	Japan	IGT/DM	IGT: 63 DM: 66	ND	44	2,534	Age	CVD death	IGT, 2.22 (1.08–4.58); DM, 2.27 (1.07–4.84)
JPHC study Saito et al. (2009)	Japan	IFG/DM	IFG: 52 DM: 53	23.9	36	31,192	Age, sex, BMI, hypertension, fasting status, community, dyslipidemia, smoking, regular alcohol drinking, sports and exercise	CHD incidence	IFG, 1.61 (1.01–2.57); DM, 4.05 (2.16–7.56)
JPHC study Cui et al. (2009)	Japan	Borderline/DM	Borderline: 55 DM: 56	23.9	37	22,528	Age, BMI, SBP, fasting status, community, LDLC, HDLC, TG, smoking, regular alcohol drinking, and use antihypertensive medication	All types of stroke incidence	Borderline, 1.01 (0.76–1.34) in men and 1.26 (0.88–1.81) in women; DM, 1.64 (1.21–2.23) and 2.19 (1.53–3.12)
Hisayama Study Doi et al. (2010)	Japan	IGT/DM	58	23.0	43	2,421	Age, BMI, SBP, ECG abnormality, LDLC, HDLC, smoking, alcohol intake, and regular exercise	Ischemic stroke incidence CHD incidence	IGT, 0.91 (0.44–1.89) in men and 0.88 (0.46–1.70) in women; DM, 2.54 (1.40–4.63) and 2.02 (1.07–3.81) IGT, 1.11 (0.62–2.00) in men and 0.82 (0.31–2.15) in women; DM, 1.26 (0.67–2.35) in women and 3.46 (1.59–7.54) in men
Suita Study Kokubo et al. (2010)	Japan	IFG/DM	IFG: 58 DM: 60	22.5	47	5,321	Age, BMI, hypertension, dyslipidemia, smoking, and drinking status	All types of stroke incidence CHD incidence	IFG, 0.97 (0.64–1.46) in men and 1.36 (0.84–2.19) in women; DM, 1.78 (1.00–3.12) in women and 2.66 (1.22–5.80) in men IFG, 1.31 (0.87–1.96) in men and 1.83 (1.01–3.32) in women; DM, 1.69 (0.86–3.31) in women and 4.32 (1.81–10.3) in men
Large claim data Fujihara et al. (2017)	Japan	Prediabetes/DM	31–60	23.6	100	111,621	BMI, SBP, LDLC, HDLC, and current smoking	CAD incidence	Age 31–40 years (males); prediabetes, 2.89 (1.02–8.19); DM 17.3 (6.36–47.0) Age 41–50 years (males); prediabetes, 0.88 (0.61–1.28); DM 2.74 (1.85–4.05); Age 51–60 years (males); prediabetes 1.62 (1.16–2.32); DM, 2.47 (1.69–3.58)

BMI: body mass index; CAD: coronary artery disease; CHD: coronary heart disease; CVD: cardiovascular disease; DM: diabetes mellitus; ECG: electrocardiogram; HDLC: high-density lipoprotein cholesterol; IFG: impaired fasting glucose; IGT: impaired glucose tolerance; JPHC: Japan Public Health Center-based Prospective Study; LDLC: low-density lipoprotein cholesterol; ND: not described; SBP: systolic blood pressure; TG: triglycerides

populations.¹⁷⁾ Similarly, diabetes confers a risk of CAD equivalent to aging 20 years in Japanese men.¹²⁾ These findings suggest that the metabolic abnormalities reflecting underlying systemic insulin resistance, which include unfavorable lipid parameter levels and other metabolic abnormalities, contribute to the development of atherosclerosis.

A meta-analysis of 102 prospective studies that included approximately 700,000 patients showed that independent of other conventional risk factors, diabetes was associated with a two-fold increase in the risk of

CVD [CAD, 2.00 (95% confidence interval, 1.83–2.19); ischemic stroke, 2.27 (1.95–2.65); and hemorrhagic stroke, 1.56 (1.19–2.15)].¹⁸⁾ Fox et al. showed that a 50% reduction in the rate of CVD events was observed among adults with diabetes in the late 1900s compared with that in the middle 1900s.¹⁹⁾ However, the absolute risk of CVD remained two-fold higher among persons with diabetes than that among those without diabetes, as shown in previous western and Japanese studies.^{19–21)} In the Hisayama study, diabetes was shown to be an independent risk factor of ischemic stroke in both sexes and

Table 2 Incidence rate of cardiovascular disease per 1,000 person-years based on overall and glucose tolerance status

Study	Country	Glucose tolerance status	% Men	No. participants total	Endpoint	Sex	Categories	Rate per 1,000 person-years
General population								
Suita Study Okamura et al. (2009)	Japan	NA	46	4,694	MI incidence			0.14
JALS-ECC study Tanabe et al. (2009)	Japan	NA	40	22,430	AMI incidence			0.65
					Stroke incidence			3.29
According to glucose tolerance status								
JPHC study Saito et al. (2009)	Japan	IFG/DM	36	31,192	CHD incidence	Men	NGT	1.0 [†]
							Borderline	1.37 [†]
							DM	2.06 [†]
						Women	NGT	0.30 [†]
							Borderline	0.29 [†]
							DM	1.02 [†]
Hisayama Study Doi et al. (2010)	Japan	NGT/IGT/DM	43	2,421	Ischemic stroke incidence	Men	FPG <5.6	5.4 [†]
							5.6–6.0	4.0 [†]
							6.1–6.9	4.7 [†]
							≥7.0	11.7 [†]
						Women	FPG <5.6	3.4 [†]
							5.6–6.0	3.9 [†]
							6.1–6.9	7.1 [†]
							≥7.0	9.6 [†]
					CHD incidence	Men	FPG <5.6	7.0 [†]
							5.6–6.0	4.7 [†]
							6.1–6.9	7.3 [†]
							≥7.0	9.9 [†]
						Women	FPG <5.6	1.4 [†]
							5.6–6.0	1.8 [†]
							6.1–6.9	2.5 [†]
							≥7.0	7.0 [†]
Large claim data Fujihara et al. (2017)	Japan	Prediabetes/DM	100	123,746	CAD incidence	Men aged 31–40 years	NGT	0.06
							Prediabetes	0.23
							DM	1.91
						Men aged 41–50 years	NGT	0.52
							Prediabetes	0.62
							DM	2.52
						Men aged 51–60 years	NGT	1.12
							Prediabetes	2.24
							DM	4.14
T2DM cohort								
JDDM Yokoyama et al. (2011)	Japan	T2DM	63	2,984	CHD			4.4
					Stroke			3.1
JDCS Sone et al. (2011)	Japan	T2DM	53	1,771	MI	Men		4.95
						Women		2.62
					Stroke	Men		8.70
						Women		6.07
Western countries								
UKPDS (1998)	UK	T2DM			MI		Intensive/conventional	14.7/17.4
					Stroke incidence			5.6/5.0

[†]Age adjusted. AMI: acute myocardial infarction; CAD: coronary artery disease; CHD: coronary heart disease; DM: diabetes mellitus; FPG: fasting plasma glucose; IFG: impaired fasting glucose; IGT: impaired glucose tolerance; JDCS: Japan Diabetes Complications Study Group; JDDM: Japan Diabetes Clinical Data Management Study Group; JPHC: Japan Public Health Center-based Prospective Study; MI: myocardial infarction; NA: not applicable; NGT: normal glucose tolerance; T2DM: type 2 diabetes mellitus; UKPDS: UK Prospective Diabetes Study

of coronary heart disease (CHD) in women.²²⁾ Also, there has been no clear change in the incidence of acute myocardial infarction.²³⁾ Although the overall risk was lower

in Japanese individuals compared with that in the western population (Table 2), individuals with diabetes had an approximately two- to four-fold increased risk of CVD

compared with those without diabetes in both Japan and western countries.

In the US, between 1990 and 2010, the rates of the three complications of diabetes surveyed were substantially decreased: acute myocardial infarction, stroke, and amputation decreased by 67.8% (59.3%–76.2%), 52.7% (40.9%–64.4%), and 51.4% (34.5%–68.2%).³⁾ Performance of health systems and production of a number of new drugs for the treatment of diabetes in the past decade might have contributed to the reduction in the rate of these complications. However, considering the pandemic of diabetes and obesity worldwide,¹⁾ diet and exercise remain the most crucial treatment for patients with diabetes. In fact, among adults with T2DM diagnosed for <10 years, a lifestyle intervention compared with standard care resulted in a change in glycemic control that did not reach the criterion for equivalence, but was in a direction consistent with benefit.²⁴⁾

Prevalence and Risk Factors of CVD in Patients with T2DM

In the Japan Diabetes Complications Study (JDCS), the crude incidence per 1,000 patient-years of myocardial infarction in patients with diabetes was 3.84 (4.95 in men and 2.62 in women),²⁵⁾ which was higher than that in the general population (0.65–1.42 per 1,000 patient-years)^{26,27)} (Table 2). However, the prevalence was lower compared with that in the western diabetic and general populations.^{28,29)} Silent infarctions were more common in individuals with diabetes compared with those without.³⁰⁾ Moreover, traditional and emerging cardiac risk factors were not associated with abnormal stress tests.³¹⁾ Therefore, detecting groups at a high risk of developing CAD among patients with T2DM is difficult. In western cohorts with T2DM, elevated concentrations of low-density lipoprotein cholesterol (LDLC), decreased concentrations of high-density lipoprotein cholesterol (HDL), hyperglycemia, hypertension, and smoking were observed to be risk factors of CAD.^{32,33)} In JDCS, male sex, elevated concentrations of LDLC, and hypertriglyceridemia are risk factors of CAD.²⁵⁾ Moreover, hypertriglyceridemia is a useful predictor of CAD in women with diabetes.³⁴⁾ No difference was observed in the prevalence of stroke between patients with T2DM in Japan and western countries (Table 2). The crude incidence per 1,000 patient-years of all strokes was 7.45 (8.70 in men and 6.07 in women); among these, the incidence of brain infarction was 6.29 (7.44 in men and 5.03 in women).²⁵⁾ Similarly, in the UK Prospective Diabetes Study (UKPDS), the incidence per 1,000 patient-years of all strokes was 5.0 and 5.6 in men and women, respectively.³⁵⁾ High blood pressure (BP) was shown to be the most important risk factor for stroke in patients with

T2DM in JDCS as well in the general population.²⁵⁾

CAD among working age people is relevant and urgent not only because of its association with premature death but also because of its enormous socioeconomic impact. A largescale claims database in Japan showed that compared with men with normal glucose tolerance, diabetes was associated with a 17.3-fold, 2.74-fold, and 2.47-fold risk in men aged 31–40 years, 41–50 years, and 41–50 years, respectively¹²⁾ (Table 1). Being a prediabetic confers an approximately three-fold higher risk of CAD in men aged 31–40 years. Impact of diabetes on CAD was markedly higher in men aged 31–40 years compared with that in men aged 41–60 years, and diabetes conferred an approximately 20-fold higher risk of CAD independently of other conventional risk factors in men aged 31–40 years. That risk was equal to those aged 51–60 years with normal glucose tolerance, similar to findings by Booth et al. that diabetes confers a risk of CAD equivalent to aging 15 years in western populations.¹⁷⁾

Relationship between Glucose Abnormality, Metabolic Phenotype, and CAD

Metabolically healthy and unhealthy obese individuals differ in terms of the cardiovascular risk.³⁶⁾ The impact of obesity and metabolically unhealthy phenotype on the development of CAD based on the glucose tolerance status showed that nonobese persons in a metabolically unhealthy state had an approximately two- to three-fold increased risk of CAD compared with those not in a metabolically unhealthy state across all glucose tolerance categories.³⁷⁾ In contrast, the presence of obesity alone showed no increase in the risk of CAD in all glucose tolerance categories. Compared with nonobese individuals with normal glucose tolerance not in a metabolically unhealthy state, the risk of CAD in obese prediabetic individuals in a metabolically unhealthy state was four-fold, which was equal to that in nonobese individuals with diabetes not in a metabolically unhealthy state.³⁷⁾

Diet, Physical Activity, and CVD in Patients with T2DM

In JDCS, increased dietary fiber, particularly soluble fiber, and intake of vegetables and fruits was associated with a lower incidence of stroke but not cerebrovascular disease in patients with T2DM.³⁸⁾ After adjustment for confounders, the hazard ratio for CVD in patients in the fourth quartile of sodium intake (mean, 15.0 g/day) after adjustment for confounders, the hazard ratio for CVD in patients in the fourth quartile of sodium intake (mean 15.0 g/day) compared with the first quartile (mean 6.4 g/day) was 2.17 (1.21–3.90).³⁹⁾ These results demonstrated that dietary

fiber intake and sodium restriction were important aspects of diet therapy in Japanese patients with T2DM.

Results of the JDCS showed that leisure-time physical activity of 15.4 metabolic equivalents (Mets) or more was associated with a significantly lower risk of stroke partly through ameliorating combinations of cardiovascular risk factors.⁴⁰⁾ The amount of daily exercise in the top tertile was >2.2 Mets·h, which was equivalent to walking >30 min per day at a speed of 5.6 km/h. However, the amount of average daily exercise for this group was 5.3 Mets·h, which was reached in about 70 min of walking with quick steps. These findings were consistent with the results of a meta-analysis,⁴¹⁾ suggesting that starting exercise contributes to the prevention of macrovascular disease in patients with T2DM without an exercise habit.

Effects of Diabetes Treatment on CVD

Intensive glycemic control and CVD in patients with T2DM

The effect of intensive glycemic control on cardiovascular outcomes in patients with T2DM is controversial. In the Kumamoto study, 110 Japanese patients with T2DM were randomly assigned to multiple insulin injection therapy groups and administered three or more daily insulin injections or assigned to conventional insulin injection therapy and followed up for 6 years.⁴²⁾ Intensive glycemic control delayed the onset and progression of the early stages of diabetic microvascular complications.⁴²⁾ The Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial investigated 4,733 patients with T2DM who were randomized to two different blood glucose targets (<6.0% and 7.0%–7.9%) and followed up for 3.4 years. This trial showed no significant differences between the groups in the primary endpoints nonfatal myocardial infarction, nonfatal stroke, and cardiovascular mortality.⁴³⁾ The reduction in cardiovascular morbidity in the intensified therapy group did not reach statistical significance. Moreover, this study was prematurely interrupted because of increased mortality in the intensive treatment group; although the intensified therapy relatively rapidly improved blood glucose control within 1 year (median HbA1c, 8.1% to 6.4%), weight gain of >10 kg was observed in 27% of the participants, and of these, 16.2% of patients experienced severe hypoglycemia. Hypoglycemia stimulates the release of catecholamines, which have profound effects on the myocardium and blood vessels.⁴⁴⁾ Catecholamines increase myocardial contractility, myocardial workload, and cardiac output. The lack of a significant reduction was observed for CVD events with intensive glycemic control in the Action in Diabetes and Vascular Disease: Preterax and Diamicron Modified Release Controlled Evaluation (ADVANCE) and Veterans Affairs Dia-

betes Trial (VADT).^{45,46)} In a subanalysis of ACCORD, the risk of death with the intensive strategy linearly increased from approximately 6% to 9% HbA1c and seemed to be higher with the intensive than that with the standard strategy only when the mean HbA1c was >7%.⁴⁷⁾ In a subanalysis of ADVANCE, severe hypoglycemia was associated with a significant increase in the adjusted risks of death from any cause [hazard ratio, 3.27 (2.39–4.65)] and cardiovascular events [3.79 (2.36–6.08)].⁴⁸⁾ These findings suggest the necessity of various types of care depending on patients' characteristics such as age, duration of diabetes, glycemic control, and comorbidity based on a patient-centered approach.

In a meta-analysis of five randomized controlled trials [UKPDS, PROspective pioglitAzone Clinical Trial In macroVascular Events (PROactive), ACCORD, ADVANCE, and VADT] that included 33,040 patients, patients in the intensive glycemic control group had a significant reduction in nonfatal myocardial infarction and CHD events by 17% and 15%, respectively.⁴⁹⁾ However, another meta-analysis of 14 clinical trials, including 28,614 patients, showed that intensive glycemic control did not reduce all-cause and cardiovascular mortality in patients with T2DM.⁵⁰⁾ Thus, there is no certain conclusion on the effect of strict blood glucose control on macrovascular disease. Future studies are needed to assess these issues, particularly focusing on older age, weight gain, influence of hypoglycemia, and rapid improvement of glycemic control.

Metabolic memory and the legacy effect

Although long observation periods are needed to reveal the advantage of glycemic control, its benefits are sustained for a relatively long period. Intensive diabetes therapy during the Diabetes Control and Complications Trial (DCCT) (6.5 years) had long-term beneficial effects on the incidence of CVD and macrovascular complications and was associated with a decrease in the glomerular filtration rate in type 1 diabetes that persisted for up to 30 years.^{51–53)} Similarly, prolonged benefits of well-controlled blood glucose were observed in UKPDS in patients with T2DM.⁵⁴⁾ These results illustrated that intensive glucose control should be initiated in the early phase of diabetes. The ACCORD Follow-on Study (ACCORDION) showed that intensive glycemic control had no beneficial effects on the incidence of death and nonfatal cardiovascular events. The increased risk of cardiovascular death reported during ACCORD remained significant during the long-term follow-up, although the magnitude of the risk was less.⁵⁵⁾ No prolonged benefit of well-controlled blood glucose was observed in VADT,⁵⁶⁾ whereas prolonged intensive treatment reduced the incidence of CVD in the ADVANCE posttrial observational study (ADVANCE-ON).⁵⁷⁾ These

findings should be cautiously interpreted because the participants in ACCORD, ADVANCE, and VADT were at a high risk of developing CVD compared with those in DCCT and UKPDS.

Effect of Various Hypoglycemic Agents on CVD

Metformin

The UKPDS trial investigated 3,277 patients with newly diagnosed T2DM who were randomized to four different treatments [conventional treatment, insulin, sulfonylurea, or metformin (>120% ideal bodyweight)] and followed up for 5 years. Metformin significantly reduced the risk of myocardial infarction and death from any cause that persisted over time compared with the control group.⁵⁴⁾ Based on this result, metformin was recommended as a first-line treatment option for T2DM in the consensus statement from the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD).⁵⁸⁾ However, these guidelines should be interpreted cautiously because impaired insulin secretion had a greater impact compared with insulin resistance on the background of T2DM in a Japanese population.^{59,60)}

Alpha-glucosidase inhibitor

Treatment with the alpha-glucosidase inhibitor acarbose significantly reduced the progression of carotid intima-media thickness by 46%⁶¹⁾ and CVD by 49%.⁶²⁾ Meta-analysis of Risk Improvement under Acarbose (MeRIA7), which evaluated the association between the control of postprandial hyperglycemia and the occurrence of cardiovascular events, showed a significant 64% decrease in the relative risk of myocardial infarction and a 35% decrease in the relative risk of any cardiovascular event.⁶³⁾ Interestingly, the HbA1c difference between the treatment and placebo groups was only 0.57%, suggesting that clinicians need to pay close attention to the control of postprandial hyperglycemia.

Thiazolidinediones

The PROactive study assessed pioglitazone for the secondary prevention of macrovascular events in patients with T2DM. Pioglitazone showed a nonsignificant 10% decrease in the relative risk of the primary composite endpoint (all-cause mortality; nonfatal myocardial infarction, including silent myocardial infarction; stroke; acute coronary syndrome; endovascular or surgical intervention in the coronary or leg arteries; and amputation above the ankle) and a significant 16% decrease in the relative risk of main secondary endpoints (death, myocardial infarction, and stroke) after a mean of 2.9 years.⁶⁴⁾

Thiazolidinediones were associated with increased body

weight, heart failure, fracture, and malignancy.^{65,66)} Moreover, the relationship between rosiglitazone and CVD has gained interest worldwide. The US Food and Drug Administration stated that pioglitazone may pose an increased risk of bladder cancer^{65,67)}; hence, it is contraindicated for patients at risk of or with bladder cancer. Similar statements were made by the European Medicines Agency.⁶⁸⁾ However, no association between the cumulative use of pioglitazone or rosiglitazone and the incidence of bladder cancer was observed in a large pooled multipopulation analysis.⁶⁹⁾ Further studies are needed before drawing conclusions on this issue.

Glinides

The Nateglinide and Valsartan in Impaired Glucose Tolerance Outcomes Research trial investigated 9,306 persons with impaired glucose tolerance randomized to two different therapies (nateglinide or placebo) and followed up for 5.0 years. This trial showed no significant differences between the groups in the primary endpoints incidence of diabetes and cardiovascular outcomes (cardiovascular death, nonfatal myocardial infarction, nonfatal stroke, or hospitalization for heart failure).⁷⁰⁾ However, the nateglinide group had an overall higher mean body weight throughout the study compared with the placebo group (mean difference, 0.35 kg). Thus, these findings suggest that the examination of treatment effects considering weight change in this type of study is necessary.

Dipeptidyl peptidase-4 inhibitors

Dipeptidyl peptidase-4 inhibitors (DPP-4Is) are key drugs for Asians, who are more likely to have impaired insulin secretion compared with Caucasians. The DPP-4I prescription rate has markedly increased in Japan in recent years.^{71–74)} A recent meta-analysis indicated that DPP-4Is exhibited better glucose-lowering efficacy in studies consisting of $\geq 50\%$ Asians compared with those consisting of $< 50\%$ Asians.⁷⁵⁾ Sitagliptin attenuated the progression of carotid intima-media thickness in insulin-treated Japanese patients with T2DM.⁷⁶⁾ In a number of randomized controlled trials that evaluated the effectiveness of DPP-4Is, no evidence showed that DPP-4Is prevent CVD.^{77–79)} The short observation period could have affected these results; hence, further studies are needed to conclude the effectiveness of DPP-4Is for CVD.

Sodium-glucose co-transporter 2 inhibitors

In the Empagliflozin Cardiovascular Outcome Event Trial in Type 2 Diabetes Mellitus Patients (EMPA-REG OUTCOME), empagliflozin added to the standard of care reduced the risk of three-point major adverse cardiovascular events (MACE) in patients with T2DM and established CVD as follows: composite of cardiovascular

death, nonfatal myocardial infarction, or nonfatal stroke by 14%, cardiovascular death by 38%, hospitalization for heart failure by 35%, and all-cause mortality by 32%.⁸⁰⁾ The same results were observed in Asians.⁸¹⁾ However, these results should be interpreted cautiously because this study investigated empagliflozin for secondary prevention in those who were at a high risk of developing CAD. Indeed, the mean age, body mass index (BMI), and duration of diabetes were 63 years, $\geq 30 \text{ kg/m}^2$, and > 10 years, respectively. Moreover, metabolic changes in addition to improvements in glycemic control by empagliflozin, such as weight loss, reduced BP, increased HDLC, and decreased uric acid, may affect CVD reduction. The Canagliflozin cardioVascular Assessment Study (CANVAS) showed that canagliflozin reduced cardiovascular events by 14% and decreased the rate of renal decline by 40%, but it also doubled the risk of lower-limb amputation.⁸²⁾

Glucagon-like peptide-1 receptor agonists

In the Liraglutide Effect and Action in Diabetes: Evaluation of Cardiovascular Outcome Results (LEADER) trial, the rate of the first four-point MACE (composite of death from cardiovascular causes, nonfatal myocardial infarction, or nonfatal stroke) among patients with T2DM was lower with liraglutide than that with placebo.⁸³⁾ Similar results were observed in the Semaglutide in Subjects with Type 2 Diabetes (SUSTAIN-6) study.⁸⁴⁾ In contrast, no superior effect on CVD was observed in the Evaluation of Lixisenatide in Patients with T2DM and Acute Coronary Syndrome (ELIXA) study and Exenatide Study of Cardiovascular Event Lowering (EXSCEL).^{85,86)} The difference in the reduction of all-cause mortality was dependent on differences in the baseline cardiovascular risk among the sample population.

Severe Hypoglycemia, Comorbidities, and CVD

Prevention of severe hypoglycemia is an important issue in the treatment of diabetes. Severe hypoglycemia causes an approximately two-fold increase in the risk of CVD.⁸⁷⁾ In this analysis, comorbid severe illness alone may not explain the association between hypoglycemia and CVD.⁸⁷⁾ Thus, clinicians should avoid the risk of hypoglycemia in the treatment of diabetes. Future studies need to clarify the relationship between each class of drug use and CVD and mortality.

Weight Gain

Weight gain is a serious issue in the treatment of diabetes. Despite improved glycemic control, the weight gain in ACCORD and VADT was 3.5 kg and 8.2 kg, respectively,

compared with baseline BMI (28.0 kg/m^2 in ACCORD and 31.3 kg/m^2 in VADT).^{43,46)} Basically, the goal of diabetes treatment is the reduction of obesity through diet and exercise. These findings emphasize the necessity of achieving glycemic control while avoiding weight gain.

Intensive BP Control and CVD in Patients with T2DM

High BP has long been recognized as a major risk factor for CVD in patients with and without diabetes.^{32,88)} Recent results of the Framingham study showed that patients with diabetes with hypertension had an increased risk of CVD death compared with those with diabetes without hypertension.⁸⁹⁾ In UKPDS, no legacy effect was observed with regard to intensive treatment of hypertension.⁹⁰⁾ Thus, ensuring long-term achievement of BP control may be essential for the reduction of CVD in patients with T2DM.

Antihypertensive therapy has been shown to be an effective method for reducing the macrovascular complications in individuals with T2DM. However, the optimal target BP levels are still under debate. The ACCORD trial investigated 4,733 patients with T2DM who were randomized to two different systolic BP targets (120 mmHg and 140 mmHg) and followed up for 4.7 years. This trial showed no significant differences between the groups in the primary endpoints nonfatal myocardial infarction, nonfatal stroke, and cardiovascular mortality.⁹¹⁾ Results of a meta-analysis suggested that impaired fasting glucose, impaired glucose tolerance, or a systolic BP treatment goal of 130 to 135 mmHg was acceptable in patients with T2DM.⁹²⁾ In addition, with more aggressive goals ($< 130 \text{ mmHg}$), target organ heterogeneity was observed in that the risk of stroke continued to decrease, but benefits with respect to the risk of other macrovascular events were not observed and the risk of serious adverse events increased. A recent meta-analysis of 49 clinical trials also showed that antihypertensive treatment reduced the risk of mortality and cardiovascular morbidity in patients with diabetes with a baseline systolic BP of $\geq 140 \text{ mmHg}$; for those a baseline systolic BP of $< \text{systolic blood pressure (SBP)} 140 \text{ mmHg}$, further treatment was associated with a rather increased risk of cardiovascular death.⁹³⁾

A meta-analysis of 40 studies showed that BP lowering was associated with improved mortality and other clinical outcomes, with lower relative risks observed among those with a baseline systolic BP of $\geq 140 \text{ mmHg}$.⁹⁴⁾ Angiotensin-converting enzyme inhibitors also reduced all-cause mortality, cardiovascular mortality, and major cardiovascular events in patients with diabetes, whereas angiotensin-receptor blockers had no benefits on these outcomes in patients with diabetes.⁹⁵⁾ Nevertheless, al-

though achieving the target level of BP first is essential, and continuing the discussion of mortality and protection from organ damage among each class of drugs, including the drug effect, is necessary.

Lipid Control and CVD in Patients with T2DM

A number of studies showed that statin use reduces cardiovascular events in patients with T2DM. The Collaborative Atorvastatin Diabetes Study (CARDS) randomized 2,838 patients with T2DM aged 40–75 years with at least one other vascular risk factor to atorvastatin 10 mg or placebo groups. This study found a 40% reduction in the LDLC level, a 37% risk reduction in time to the first cardiovascular event, a 31% reduction in the risk of coronary revascularization, and a 48% reduction in the risk of stroke over 3.9 years with atorvastatin.⁹⁶⁾

Management of Elevated Cholesterol in the Primary Prevention Group of Adult Japanese (MEGA) randomized 7,832 patients with hypercholesterolemia without a history of CHD or stroke to diet alone or diet plus low-dose pravastatin (10–20 mg/day) groups. Diet plus pravastatin reduced the mean LDLC by 18%, and this reduction was associated with a 33% lower incidence of CHD events compared with that in the diet group during the 5.3 years of follow-up.⁹⁷⁾ In a subanalysis, these effects were confirmed in patients with diabetes and/or impaired fasting glucose.⁹⁸⁾

An LDLC reduction of 2.0–3.0 mmol/L (77–116 mg/dL) may be able to suppress vascular events by approximately 40%–50%. A meta-analysis investigating almost 170,000 people showed a significant reduction of 22% per 1.0 mmol/L (38.7 mg/dL) in the risk of LDLC.⁹⁹⁾ A meta-analysis of 14 prospective studies that included 18,868 patients showed that independent of other conventional risk factors, statin therapy was associated with a 13% reduction in the risk of mortality.¹⁰⁰⁾ However, these findings are about the absolute value of LDLC and do not mention the quality of LDLC. In patients with T2DM, hypertriglyceridemia and low HDLC are frequently observed rather than high total cholesterol and LDLC due to a decrease in insulin action. Indeed, hypertriglyceridemia is comparable to LDLC as a risk factor for CVD in East Asians, including Japanese.^{25,101)} Moreover, not LDLC but remnant, small dense LDL and glycated LDL are potent risk factors as atherogenic factors. In addition, the relationship between LDLC and other lipid variables, such as HDLC, is important. Thus, although we mainly focused on LDLC, clinicians must also pay attention to the control of the entire lipid profile.

With regard to the effects of long-term fenofibrate therapy on cardiovascular events in 9,795 people with

T2DM (the FIELD study), although fenofibrate did not significantly reduce the primary outcome consisting of incident CHD, cardiovascular mortality, or total mortality in patients with T2DM treated for 5 years, it reduced CHD.¹⁰²⁾ In a subanalysis of the benefit from long-term fenofibrate in patients with T2DM and various eGFR categories in comparison with placebo, risks of side effects of fenofibrate were not increased in any eGFR category.¹⁰³⁾ In the ACCORD Lipid study, a benefit was shown in a prespecified subanalysis of participants with both higher triglycerides (≥ 204 mg/dL) and low HDLC (≤ 34 mg/dL) at baseline.¹⁰⁴⁾ Although statins are recommended as first-line therapy for CVD prevention, these findings suggested the effectiveness of fibrate treatment, particularly in East Asians because hypertriglyceridemia is a risk factor of CVD in this population.^{25,105)}

Aspirin for Primary Prevention of CVD in Patients with T2DM

The Japanese Primary Prevention of Atherosclerosis with Aspirin for Diabetes (JPAD) study examined the efficacy of low-dose aspirin for the primary prevention of cardiovascular events in a randomized 163-center, open-label trial conducted in 2,539 Japanese patients with T2DM but without a history of CVD. The patients were assigned to either aspirin (81–100 mg daily) or no aspirin groups and were followed up for a mean of 4.4 years. This trial revealed no significant differences between the groups in the primary endpoints fatal or nonfatal ischemic heart disease, fatal or nonfatal stroke, and peripheral arterial disease [hazard ratio, 0.80 (0.58–1.10)].¹⁰⁶⁾ In subgroup analyses, however, among participants aged >65 years, the incidence of the primary endpoint was lower with aspirin [hazard ratio, 0.68 (0.46–0.99)]. Based on these studies, a position statement of the ADA, American Heart Association, and American College of Cardiology Foundation noted that aspirin should not be recommended for CVD prevention in adults with diabetes at a low risk of CVD (men <50 years and women <60 years without major additional CVD risk factors; 10-year CVD risk of $<5\%$) because the potential adverse effects of bleeding offset the potential benefits.^{107,108)}

In the Principal Results of the Diabetic Atherosclerosis Prevention by Cilostazol (DAPC) Study, the investigators compared the efficacy of cilostazol (100–200 mg/day) and aspirin (81–100 mg/day) on carotid atherosclerosis in a randomized, open, blinded endpoint study conducted in four East Asian countries on participants with T2DM. The patients were assigned to either the phosphodiesterase inhibitor cilostazol or no aspirin group and were followed for 2.0 years. Compared with aspirin, cilostazol potentially inhibited the progression of carotid intima-media thickness.¹⁰⁹⁾

Effect of Control of Multiple Risk Factors for CAD in Patients with T2DM

The Steno-2 study investigated 160 patients with T2DM randomized to two different therapies (intensive therapy and usual treatment) and followed up for approximately 8 years. Intensive multifactorial treatment (including glycemic control) reduced CVD by 53%.¹¹⁰⁾ Moreover, intensive intervention showed beneficial effects with respect to vascular complications over a period of time.¹¹¹⁾ The Japan Diabetes Optimal Integrated Treatment Study for 3 Major Risk Factors of Cardiovascular Diseases (J-DOIT3) trial investigated 2,540 patients with T2DM aged 45 to 69 years who were randomized to two different therapies (intensive therapy and usual treatment) and followed up for approximately 8.5 years.¹¹²⁾ Although the reduction in the primary outcome was nonsignificant, adjustment for variables, such as smoking, showed that a significantly reduced risk of these outcomes was found among the intensive therapy group.¹¹²⁾

Prediction of CVD in Japanese Patients with T2DM

Epidemiology shows the mean or trends of a specific group. By integrating the risk factor for each complication of CVD, future complications in each patient with T2DM can be possibly predicted. The risk engine (Japan Diabetes Complications Study/the Japanese Elderly Diabetes Intervention Trial risk engine) made from data of the JDCS and The Japanese Elderly Intervention Trial (J-EDIT) accurately predicted macro- and microvascular complications and would provide helpful information in risk classification in Japanese.¹¹³⁾

Future Perspectives

Although a number of clinical studies of diabetes and CVD are being conducted worldwide, clinical studies involve a large cost, enormous effort, and cooperation of participants. To resolve such issues, using new and alternative research sources, such as big data, is essential. Big data has the following characteristics that can complement the abovementioned issues: use of available existing datasets, low cost, available data accumulated over a long period, and expectation of an increasing number of events with time. In the future, comprehensive big medical databases covering lifestyle habits and genetic information will make it possible to achieve more precise treatment of patients with T2DM.

Conclusion

Considering that East Asian patients with T2DM, including Japanese, have different features compared with western patients, applying the same guidelines for the treatment of these two groups is difficult. Clinical studies demonstrated that intensive glucose control should be initiated in the early phase of diabetes for macrovascular disease accompanied by lipid and BP control. In the future, in addition to standardized therapies, comprehensive medical big databases covering lifestyle habits and genetic information will make it possible to achieve precision in the treatment of patients with T2DM.

Acknowledgments

This work is supported by JSPS. No potential conflicts of interest relevant to this article were reported. The sponsor had no role in the design and conduct of the study. The authors also thank Mami Haga and Natsuko Tada, Niigata University Faculty of Medicine for excellent secretarial assistance.

Disclosure Statement

K. Fujihara has no conflicts of interest. H. Sone has received donations for research from Kyowa Hakko Kirin, MSD, Boehringer Ingelheim, Sumitomo Dainippon, Ono, Teijin Pharma, Kowa Souyaku, Takeda, Taishotoyama, and Daiichi Sankyo.

Author Contributions

Review conception: KF, HS

Writing: KF

Critical review and revision: all authors

Final approval of the article: all authors

Accountability for all aspects of the work: all authors

References

- 1) IDF Diabetes Atlas 2015 (International Diabetes Federation) (<http://www.diabetesatlas.org/>). 2015.
- 2) National Diabetes Fact Sheet, National Center for Chronic Disease Prevention and Health Promotion, US (http://www.cdc.gov/diabetes/pubs/pdf/ndfs_2011.pdf).
- 3) Gregg EW, Li Y, Wang J, et al. Changes in diabetes-related complications in the United States, 1990–2010. *N Engl J Med* 2014; 370: 1514–23.
- 4) Kannel WB, McGee DL. Diabetes and cardiovascular disease. The Framingham study. *JAMA* 1979; 241: 2035–8.
- 5) Stamler J, Vaccaro O, Neaton JD, et al. Diabetes, other risk factors, and 12-yr cardiovascular mortality for men screened in the Multiple Risk Factor Intervention Trial.

- Diabetes Care 1993; 16: 434-44.
- 6) Ueshima H, Sekikawa A, Miura K, et al. Cardiovascular disease and risk factors in Asia: a selected review. *Circulation* 2008; 118: 2702-9.
 - 7) Yokoyama H, Matsushima M, Kawai K, et al. Low incidence of cardiovascular events in Japanese patients with Type 2 diabetes in primary care settings: a prospective cohort study (JDDM 20). *Diabet Med* 2011; 28: 1221-8.
 - 8) Hotta N, Nakamura J, Iwamoto Y, et al. Causes of death in Japanese diabetics: a questionnaire survey of 18,385 diabetics over a 10-year period. *J Diabetes Investig* 2010; 1: 66-76.
 - 9) Nakamura J, Kamiya H, Haneda M, et al. Causes of death in Japanese patients with diabetes based on the results of a survey of 45,708 cases during 2001–2010—report from the Committee on the Cause of Death in Diabetes Mellitus—. *J Japan Diab Soc* 2016; 59: 667-84.
 - 10) Haffner SM, Lehto S, Ronnema T, et al. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N Engl J Med* 1998; 339: 229-34.
 - 11) Tominaga M, Eguchi H, Manaka H, et al. Impaired glucose tolerance is a risk factor for cardiovascular disease, but not impaired fasting glucose. The Funagata Diabetes Study. *Diabetes Care* 1999; 22: 920-4.
 - 12) Fujihara K, Igarashi R, Yamamoto M, et al. Impact of glucose tolerance status on the development of coronary artery disease among working-age men. *Diabetes Metab* 2017; 43: 261-4.
 - 13) Huang Y, Cai X, Mai W, et al. Association between pre-diabetes and risk of cardiovascular disease and all cause mortality: systematic review and meta-analysis. *BMJ* 2016; 355: i5953.
 - 14) Ford ES, Zhao G, Li C. Pre-diabetes and the risk for cardiovascular disease: a systematic review of the evidence. *J Am Coll Cardiol* 2010; 55: 1310-7.
 - 15) Barr EL, Zimmet PZ, Welborn TA, et al. Risk of cardiovascular and all-cause mortality in individuals with diabetes mellitus, impaired fasting glucose, and impaired glucose tolerance: the Australian Diabetes, Obesity, and Lifestyle Study (AusDiab). *Circulation* 2007; 116: 151-7.
 - 16) Henry P, Thomas F, Benetos A, et al. Impaired fasting glucose, blood pressure and cardiovascular disease mortality. *Hypertension* 2002; 40: 458-63.
 - 17) Booth GL, Kapral MK, Fung K, et al. Relation between age and cardiovascular disease in men and women with diabetes compared with non-diabetic people: a population-based retrospective cohort study. *Lancet* 2006; 368: 29-36.
 - 18) Sarwar N, Gao P, Seshasai SR, et al. Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: a collaborative meta-analysis of 102 prospective studies. *Lancet* 2010; 375: 2215-22.
 - 19) Fox CS, Coady S, Sorlie PD, et al. Trends in cardiovascular complications of diabetes. *JAMA* 2004; 292: 2495-9.
 - 20) Kannel WB, McGee DL. Diabetes and glucose tolerance as risk factors for cardiovascular disease: the Framingham study. *Diabetes Care* 1979; 2: 120-6.
 - 21) Fujishima M, Kiyohara Y, Kato I, et al. Diabetes and cardiovascular disease in a prospective population survey in Japan: the Hisayama Study. *Diabetes* 1996; 45 Suppl 3: S14-6.
 - 22) Doi Y, Ninomiya T, Hata J, et al. Impact of glucose tolerance status on development of ischemic stroke and coronary heart disease in a general Japanese population: the Hisayama study. *Stroke* 2010; 41: 203-9.
 - 23) Hata J, Ninomiya T, Hirakawa Y, et al. Secular trends in cardiovascular disease and its risk factors in Japanese: half-century data from the Hisayama Study (1961–2009). *Circulation* 2013; 128: 1198-205.
 - 24) Johansen MY, MacDonald CS, Hansen KB, et al. Effect of an intensive lifestyle intervention on glycemic control in patients with type 2 diabetes: a randomized clinical trial. *JAMA* 2017; 318: 637-46.
 - 25) Sone H, Tanaka S, Tanaka S, et al. Serum level of triglycerides is a potent risk factor comparable to LDL cholesterol for coronary heart disease in Japanese patients with type 2 diabetes: subanalysis of the Japan Diabetes Complications Study (JDCS). *J Clin Endocrinol Metab* 2011; 96: 3448-56.
 - 26) Tanabe N, Iso H, Okada K, et al. Serum total and non-high-density lipoprotein cholesterol and the risk prediction of cardiovascular events—the JALS-ECC. *Circ J* 2010; 74: 1346-56.
 - 27) Okamura T, Kokubo Y, Watanabe M, et al. Low-density lipoprotein cholesterol and non-high-density lipoprotein cholesterol and the incidence of cardiovascular disease in an urban Japanese cohort study: the Suita study. *Atherosclerosis* 2009; 203: 587-92.
 - 28) Hadamitzky M, Achenbach S, Al-Mallah M, et al. Optimized prognostic score for coronary computed tomographic angiography: results from the CONFIRM registry (COronary CT Angiography EvaluationN For Clinical Outcomes: an InteRnational Multicenter Registry). *J Am Coll Cardiol* 2013; 62: 468-76.
 - 29) Rana JS, Dunning A, Achenbach S, et al. Differences in prevalence, extent, severity, and prognosis of coronary artery disease among patients with and without diabetes undergoing coronary computed tomography angiography: results from 10,110 individuals from the CONFIRM (COronary CT Angiography EvaluationN For Clinical Outcomes): an InteRnational Multicenter Registry. *Diabetes Care* 2012; 35: 1787-94.
 - 30) Janand-Delenne B, Savin B, Habib G, et al. Silent myocardial ischemia in patients with diabetes: who to screen. *Diabetes Care* 1999; 22: 1396-400.
 - 31) Wackers FJ, Young LH, Inzucchi SE, et al. Detection of silent myocardial ischemia in asymptomatic diabetic subjects: the DIAD study. *Diabetes Care* 2004; 27: 1954-61.
 - 32) Turner RC, Millns H, Neil HA, et al. Risk factors for coronary artery disease in noninsulin dependent diabetes mellitus: United Kingdom prospective diabetes study (UKPDS: 23). *BMJ* 1998; 316: 823-8.
 - 33) Kramer CK, Rodrigues TC, Canani LH, et al. Diabetic retinopathy predicts all-cause mortality and cardiovascular events in both type 1 and 2 diabetes: meta-analysis of observational studies. *Diabetes Care* 2011; 34: 1238-44.
 - 34) Sone H, Tanaka S, Tanaka S, et al. Comparison of various lipid variables as predictors of coronary heart disease in Japanese men and women with type 2 diabetes: subanalysis of the Japan Diabetes Complications Study. *Diabetes Care* 2012; 35: 1150-7.

- 35) UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998; **352**: 837-53.
- 36) Stefan N, Haring HU, Hu FB, et al. Metabolically healthy obesity: epidemiology, mechanisms, and clinical implications. *Lancet Diabetes Endocrinol* 2013; **1**: 152-62.
- 37) Fujihara K, Matsubayashi Y, Yamamoto M, et al. Impact of body mass index and metabolic phenotypes on coronary artery disease according to glucose tolerance status. *Diabetes Metab* 2017; **43**: 543-6.
- 38) Tanaka S, Yoshimura Y, Kamada C, et al. Intakes of dietary fiber, vegetables, and fruits and incidence of cardiovascular disease in Japanese patients with type 2 diabetes. *Diabetes Care* 2013; **36**: 3916-22.
- 39) Horikawa C, Yoshimura Y, Kamada C, et al. Dietary sodium intake and incidence of diabetes complications in Japanese patients with type 2 diabetes: analysis of the Japan Diabetes Complications Study (JDCS). *J Clin Endocrinol Metab* 2014; **99**: 3635-43.
- 40) Sone H, Tanaka S, Tanaka S, et al. Leisure-time physical activity is a significant predictor of stroke and total mortality in Japanese patients with type 2 diabetes: analysis from the Japan Diabetes Complications Study (JDCS). *Diabetologia* 2013; **56**: 1021-30.
- 41) Kodama S, Tanaka S, Heianza Y, et al. Association between physical activity and risk of all-cause mortality and cardiovascular disease in patients with diabetes: a meta-analysis. *Diabetes Care* 2013; **36**: 471-9.
- 42) Shichiri M, Kishikawa H, Ohkubo Y, et al. Long-term results of the Kumamoto Study on optimal diabetes control in type 2 diabetic patients. *Diabetes Care* 2000; **23 Suppl 2**: B21-9.
- 43) Gerstein HC, Miller ME, Byington RP, et al. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008; **358**: 2545-59.
- 44) Desouza CV, Bolli GB, Fonseca V. Hypoglycemia, diabetes, and cardiovascular events. *Diabetes Care* 2010; **33**: 1389-94.
- 45) Patel A, MacMahon S, Chalmers J, et al. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2008; **358**: 2560-72.
- 46) Duckworth W, Abairra C, Moritz T, et al. Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med* 2009; **360**: 129-39.
- 47) Riddle MC, Ambrosius WT, Brillon DJ, et al. Epidemiologic relationships between A1C and all-cause mortality during a median 3.4-year follow-up of glycemic treatment in the ACCORD trial. *Diabetes Care* 2010; **33**: 983-90.
- 48) Zoungas S, Patel A, Chalmers J, et al. Severe hypoglycemia and risks of vascular events and death. *N Engl J Med* 2010; **363**: 1410-8.
- 49) Ray KK, Seshasai SR, Wijesuriya S, et al. Effect of intensive control of glucose on cardiovascular outcomes and death in patients with diabetes mellitus: a meta-analysis of randomised controlled trials. *Lancet* 2009; **373**: 1765-72.
- 50) Hemmingsen B, Lund SS, Glud C, et al. Intensive glycaemic control for patients with type 2 diabetes: systematic review with meta-analysis and trial sequential analysis of randomised clinical trials. *BMJ* 2011; **343**: d6898.
- 51) Nathan DM, Cleary PA, Backlund JY, et al. Intensive diabetes treatment and cardiovascular disease in patients with type 1 diabetes. *N Engl J Med* 2005; **353**: 2643-53.
- 52) de Boer IH, Sun W, Cleary PA, et al. Intensive diabetes therapy and glomerular filtration rate in type 1 diabetes. *N Engl J Med* 2011; **365**: 2366-76.
- 53) Diabetes Control and Complications Trial (DCCT)/Epidemiology of Diabetes Interventions and Complications (EDIC) Study Research Group. Intensive diabetes treatment and cardiovascular outcomes in type 1 diabetes: the DCCT/EDIC study 30-year follow-up. *Diabetes Care* 2016; **39**: 686-93.
- 54) Holman RR, Paul SK, Bethel MA, et al. 10-year follow-up of intensive glucose control in type 2 diabetes. *N Engl J Med* 2008; **359**: 1577-89.
- 55) ACCORD Study Group. Nine-year effects of 3.7 years of intensive glycemic control on cardiovascular outcomes. *Diabetes Care* 2016; **39**: 701-8.
- 56) Hayward RA, Reaven PD, Wiitala WL, et al. Follow-up of glycemic control and cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2015; **372**: 2197-206.
- 57) Zoungas S, Chalmers J, Neal B, et al. Follow-up of blood-pressure lowering and glucose control in type 2 diabetes. *N Engl J Med* 2014; **371**: 1392-406.
- 58) American Diabetes Association. Standards of medical care in diabetes—2017. *Diabetes Care* 2017; **40**: S1-135.
- 59) Fukushima M, Suzuki H, Seino Y. Insulin secretion capacity in the development from normal glucose tolerance to type 2 diabetes. *Diabetes Res Clin Pract* 2004; **66 Suppl 1**: S37-43.
- 60) Morimoto A, Tatsumi Y, Deura K, et al. Impact of impaired insulin secretion and insulin resistance on the incidence of type 2 diabetes mellitus in a Japanese population: the Saku study. *Diabetologia* 2013; **56**: 1671-9.
- 61) Hanefeld M, Chiasson JL, Koehler C, et al. Acarbose slows progression of intima-media thickness of the carotid arteries in subjects with impaired glucose tolerance. *Stroke* 2004; **35**: 1073-8.
- 62) Chiasson JL, Josse RG, Gomis R, et al. Acarbose treatment and the risk of cardiovascular disease and hypertension in patients with impaired glucose tolerance: the STOP-NIDDM trial. *JAMA* 2003; **290**: 486-94.
- 63) Hanefeld M, Gagatay M, Petrowitsch T, et al. Acarbose reduces the risk for myocardial infarction in type 2 diabetic patients: meta-analysis of seven long-term studies. *Eur Heart J* 2004; **25**: 10-6.
- 64) Dormandy JA, Charbonnel B, Eckland DJ, et al. Secondary prevention of macrovascular events in patients with type 2 diabetes in the PROactive Study (PROspective pioglitazone Clinical Trial In macroVascular Events): a randomised controlled trial. *Lancet* 2005; **366**: 1279-89.
- 65) Piccinni C, Motola D, Marchesini G, et al. Assessing the association of pioglitazone use and bladder cancer through drug adverse event reporting. *Diabetes Care* 2011; **34**: 1369-71.
- 66) Yki-Järvinen H. Thiazolidinediones. *N Engl J Med* 2004; **351**: 1106-18.
- 67) Tseng CH. Pioglitazone and bladder cancer: a population-based study of Taiwanese. *Diabetes Care* 2012; **35**: 278-80.
- 68) European Medicines Agency clarifies opinion on pio-

- glitazone and the risk of bladder cancer (http://www.ema.europa.eu/docs/en_GB/document_library/Press_release/2011/10/WC500116936.pdf). 2011.
- 69) Levin D, Bell S, Sund R, et al. Pioglitazone and bladder cancer risk: a multipopulation pooled, cumulative exposure analysis. *Diabetologia* 2015; **58**: 493-504.
 - 70) Holman RR, Haffner SM, McMurray JJ, et al. Effect of nateglinide on the incidence of diabetes and cardiovascular events. *N Engl J Med* 2010; **362**: 1463-76.
 - 71) Oishi M, Yamazaki K, Okuguchi F, et al. Changes in oral antidiabetic prescriptions and improved glycemic control during the years 2002–2011 in Japan (JDDM32). *J Diabetes Investig* 2014; **5**: 581-7.
 - 72) Kohro T, Yamazaki T, Sato H, et al. Trends in antidiabetic prescription patterns in Japan from 2005 to 2011. *Int Heart J* 2013; **54**: 93-7.
 - 73) Fujihara K, Hanyu O, Heianza Y, et al. Comparison of clinical characteristics in patients with type 2 diabetes among whom different antihyperglycemic agents were prescribed as monotherapy or combination therapy by diabetes specialists. *J Diabetes Investig* 2016; **7**: 260-9.
 - 74) Fujihara K, Igarashi R, Matsunaga S, et al. Comparison of baseline characteristics and clinical course in Japanese patients with type 2 diabetes among whom different types of oral hypoglycemic agents were chosen by diabetes specialists as initial monotherapy (JDDM 42). *Medicine (Baltimore)* 2017; **96**: e6122.
 - 75) Kim YG, Hahn S, Oh TJ, et al. Differences in the glucose-lowering efficacy of dipeptidyl peptidase-4 inhibitors between Asians and non-Asians: a systematic review and meta-analysis. *Diabetologia* 2013; **56**: 696-708.
 - 76) Mita T, Katakami N, Shiraiwa T, et al. Sitagliptin attenuates the progression of carotid intima-media thickening in insulin-treated patients with type 2 diabetes: the Sitagliptin Preventive Study of Intima-Media Thickness Evaluation (SPIKE): a randomized controlled trial. *Diabetes Care* 2016; **39**: 455-64.
 - 77) White WB, Cannon CP, Heller SR, et al. Alogliptin after acute coronary syndrome in patients with type 2 diabetes. *N Engl J Med* 2013; **369**: 1327-35.
 - 78) Scirica BM, Bhatt DL, Braunwald E, et al. Saxagliptin and cardiovascular outcomes in patients with type 2 diabetes mellitus. *N Engl J Med* 2013; **369**: 1317-26.
 - 79) Green JB, Bethel MA, Armstrong PW, et al. Effect of sitagliptin on cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2015; **373**: 232-42.
 - 80) Zinman B, Wanner C, Lachin JM, et al. Empagliflozin, cardiovascular outcomes, and mortality in type 2 diabetes. *N Engl J Med* 2015; **373**: 2117-28.
 - 81) Kaku K, Lee J, Mattheus M, et al. Empagliflozin and cardiovascular outcomes in Asian patients with type 2 diabetes and established cardiovascular disease—results from EMPA-REG OUTCOME®. *Circ J* 2017; **81**: 227-34.
 - 82) Neal B, Perkovic V, Mahaffey KW, et al. Canagliflozin and cardiovascular and renal events in type 2 diabetes. *N Engl J Med* 2017; **377**: 644-57.
 - 83) Marso SP, Daniels GH, Brown-Frandsen K, et al. Liraglutide and cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2016; **375**: 311-22.
 - 84) Marso SP, Bain SC, Consoli A, et al. Semaglutide and cardiovascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2016; **375**: 1834-44.
 - 85) Pfeffer MA, Claggett B, Diaz R, et al. Lixisenatide in patients with type 2 diabetes and acute coronary syndrome. *N Engl J Med* 2015; **373**: 2247-57.
 - 86) Holman RR, Bethel MA, Mentz RJ, et al. Effects of once-weekly exenatide on cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2017; **377**: 1228-39.
 - 87) Goto A, Arah OA, Goto M, et al. Severe hypoglycaemia and cardiovascular disease: systematic review and meta-analysis with bias analysis. *BMJ* 2013; **347**: f4533.
 - 88) Ogden LG, He J, Lydick E, et al. Long-term absolute benefit of lowering blood pressure in hypertensive patients according to the JNC VI risk stratification. *Hypertension* 2000; **35**: 539-43.
 - 89) Chen G, McAlister FA, Walker RL, et al. Cardiovascular outcomes in framingham participants with diabetes: the importance of blood pressure. *Hypertension* 2011; **57**: 891-7.
 - 90) Holman RR, Paul SK, Bethel MA, et al. Long-term follow-up after tight control of blood pressure in type 2 diabetes. *N Engl J Med* 2008; **359**: 1565-76.
 - 91) Cushman WC, Evans GW, Byington RP, et al. Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med* 2010; **362**: 1575-85.
 - 92) Bangalore S, Kumar S, Lobach I, et al. Blood pressure targets in subjects with type 2 diabetes mellitus/impaired fasting glucose: observations from traditional and bayesian random-effects meta-analyses of randomized trials. *Circulation* 2011; **123**: 2799-810.
 - 93) Brunström M, Carlberg B. Effect of antihypertensive treatment at different blood pressure levels in patients with diabetes mellitus: systematic review and meta-analyses. *BMJ* 2016; **352**: i717.
 - 94) Emdin CA, Rahimi K, Neal B, et al. Blood pressure lowering in type 2 diabetes: a systematic review and meta-analysis. *JAMA* 2015; **313**: 603-15.
 - 95) Cheng J, Zhang W, Zhang X, et al. Effect of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers on all-cause mortality, cardiovascular deaths, and cardiovascular events in patients with diabetes mellitus: a meta-analysis. *JAMA Intern Med* 2014; **174**: 773-85.
 - 96) Colhoun HM, Betteridge DJ, Durrington PN, et al. Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): multicentre randomised placebo-controlled trial. *Lancet* 2004; **364**: 685-96.
 - 97) Nakamura H, Arakawa K, Itakura H, et al. Primary prevention of cardiovascular disease with pravastatin in Japan (MEGA Study): a prospective randomised controlled trial. *Lancet* 2006; **368**: 1155-63.
 - 98) Tajima N, Kurata H, Nakaya N, et al. Pravastatin reduces the risk for cardiovascular disease in Japanese hypercholesterolemic patients with impaired fasting glucose or diabetes: diabetes subanalysis of the Management of Elevated Cholesterol in the Primary Prevention Group of Adult Japanese (MEGA) Study. *Atherosclerosis* 2008; **199**: 455-62.
 - 99) Baigent C, Blackwell L, Emberson J, et al. Efficacy and safety of more intensive lowering of LDL cholesterol: a

- meta-analysis of data from 170,000 participants in 26 randomised trials. *Lancet* 2010; **376**: 1670-81.
- 100) Kearney PM, Blackwell L, Collins R, et al. Efficacy of cholesterol-lowering therapy in 18,686 people with diabetes in 14 randomised trials of statins: a meta-analysis. *Lancet* 2008; **371**: 117-25.
 - 101) Chan WB, Tong PC, Chow CC, et al. Triglyceride predicts cardiovascular mortality and its relationship with glycaemia and obesity in Chinese type 2 diabetic patients. *Diabetes Metab Res Rev* 2005; **21**: 183-8.
 - 102) Keech A, Simes RJ, Barter P, et al. Effects of long-term fenofibrate therapy on cardiovascular events in 9,795 people with type 2 diabetes mellitus (the FIELD study): randomised controlled trial. *Lancet* 2005; **366**: 1849-61.
 - 103) Ting RD, Keech AC, Drury PL, et al. Benefits and safety of long-term fenofibrate therapy in people with type 2 diabetes and renal impairment: the FIELD Study. *Diabetes Care* 2012; **35**: 218-25.
 - 104) Ginsberg HN, Elam MB, Lovato LC, et al. Effects of combination lipid therapy in type 2 diabetes mellitus. *N Engl J Med* 2010; **362**: 1563-74.
 - 105) Sone H, Tanaka S, Ishibashi S, et al. The new worldwide definition of metabolic syndrome is not a better diagnostic predictor of cardiovascular disease in Japanese diabetic patients than the existing definitions: additional analysis from the Japan Diabetes Complications Study. *Diabetes Care* 2006; **29**: 145-7.
 - 106) Ogawa H, Nakayama M, Morimoto T, et al. Low-dose aspirin for primary prevention of atherosclerotic events in patients with type 2 diabetes: a randomized controlled trial. *JAMA* 2008; **300**: 2134-41.
 - 107) Pignone M, Alberts MJ, Colwell JA, et al. Aspirin for primary prevention of cardiovascular events in people with diabetes: a position statement of the American Diabetes Association, a scientific statement of the American Heart Association, and an expert consensus document of the American College of Cardiology Foundation. *Diabetes Care* 2010; **33**: 1395-402.
 - 108) Pignone M, Alberts MJ, Colwell JA, et al. Aspirin for primary prevention of cardiovascular events in people with diabetes: a position statement of the American Diabetes Association, a scientific statement of the American Heart Association, and an expert consensus document of the American College of Cardiology Foundation. *Circulation* 2010; **121**: 2694-701.
 - 109) Katakami N, Kim YS, Kawamori R, et al. The phosphodiesterase inhibitor cilostazol induces regression of carotid atherosclerosis in subjects with type 2 diabetes mellitus: principal results of the Diabetic Atherosclerosis Prevention by Cilostazol (DAPC) study: a randomized trial. *Circulation* 2010; **121**: 2584-91.
 - 110) Gæde P, Vedel P, Larsen N, et al. Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. *N Engl J Med* 2003; **348**: 383-93.
 - 111) Gæde P, Lund-Andersen H, Parving HH, et al. Effect of a multifactorial intervention on mortality in type 2 diabetes. *N Engl J Med* 2008; **358**: 580-91.
 - 112) Ueki K, Sasako T, Kato M, et al. Effect of an intensified multifactorial intervention on cardiovascular outcomes and mortality in type 2 diabetes (J-DOIT3): an open-label, randomised controlled trial. *Lancet Diabetes Endocrinol* 2017; Oct 24. DOI: 10.1016/S2213-8587(17)30327-3 [Epub ahead of print]
 - 113) Tanaka S, Tanaka S, Iimuro S, et al. Predicting macro- and microvascular complications in type 2 diabetes: the Japan Diabetes Complications Study/the Japanese Elderly Diabetes Intervention Trial risk engine. *Diabetes Care* 2013; **36**: 1193-9.