How to Do It

Tailor-Made Tapering Grafts for Large-Neck Aorta

Takuro Shirasu, MD, PhD,^{1,2} Masaru Kimura, MD, PhD,¹ Takanori Kaneko, MD,¹ Takatoshi Furuya, MD, PhD,¹ Kaito Fukuda, MD,¹ Motoki Nagai, MD,¹ and Yukihiro Nomura, MD¹

Patients having a large aortic neck poses a challenge in abdominal aortic aneurysm surgery both in endovascular and open aneurysm repair, sometimes necessitating paravisceral or thoracoabdominal aneurysm repair which carries considerable perioperative risk. Here, we describe techniques of using a tailor-made tapering graft in open surgery that can be adjusted for large neck morphology. This technique helps avoid discrepancies between the proximal aorta and graft, and postoperative acute kidney injury by clamping at lower levels. The conscientious use of this technique in selected patients realizes satisfactory outcomes both in the short term and midterm in the demanding anatomy of large aortic necks.

Keywords: abdominal aortic aneurysm, large neck, hostile neck, open aneurysm repair, aortomegaly

Introduction

A large aortic neck is one of the hindrances to endovascular aneurysm repair; if it is operated on endovascularly, the risk of type Ia endoleak and subsequent death due to rupture could carry over even five years later in various major devices.^{1–3} The aneurysms sometimes constrain paravisceral aneurysm repair, which is more invasive with

substantial morbidity and mortality both in endovascular and open repair.⁴ Open repair for a large-neck abdominal aortic aneurysm (AAA) is more durable but not straightforward due to discrepancies in sizes between the proximal aorta and general prosthetic graft. We use tailor-made tapering grafts to address the predicaments when dealing with large proximal aortas.

Case Report

An 82-year-old male patient was diagnosed with pararenal AAA of 56 mm in a maximal diameter, that had gradually grown in the 10-year follow-up. The aortic diameters at celiac, superior mesenteric/right renal, and left renal artery levels were 33 mm, 32 mm, and 43 mm, respectively. He chose to undergo open aneurysm repair with informed consent. His aorta was large (aortomegaly); therefore, we planned to clamp it just below the superior mesenteric/right renal artery level (32 mm) and reconstruct the left renal artery with bypass grafting. Through a midline incision, the left renal vein was divided, and the bilateral renal arteries and proximal aorta were looped for clamping (Fig. 1A). After confirming the diameter at the anastomo-

Fig. 1 Preparation of the proximal neck. (A) The left renal vein was dissected. (B) Both renal arteries and the inter-renal aorta were looped after the division of the left renal vein. The level and size of the proximal aorta were carefully inspected before anastomosis.

¹Department of Surgery, Asahi General Hospital, Asahi, Chiba, Japan

²Department of Surgery, University of Virginia, Charlottesville, VA, USA

Received: October 6, 2021; Accepted: November 29, 2021
Corresponding author: Takuro Shirasu, MD, PhD. Department of Surgery, University of Virginia, 409 Lane Rd MR4, Charlottesville, VA, USA 22903
Tel: +1-434-924-5772, Fax: +1-434-924-1218
E-mail: shirasu-tky@umin.ac.jp

©2022 The Editorial Committee of Annals of Vascular Diseases. This article is distributed under the terms of the Creative Commons Attribution License, which permits use, distribution, and reproduction in any medium, provided the credit of the original work, a link to the license, and indication of any change are properly given, and the original work is not used for commercial purposes. Remixed or transformed contributions must be distributed under the same license as the original.

Fig. 2 Schematic of a tailor-made tapering graft. (A) Transection of the trunk of the bifurcated graft. (B) An isosceles triangle is cut from the detached body of the graft. An acute rather than obtuse isosceles triangle is preferred. (C) The longitudinal cut of the forefront of the trunk of the bifurcated graft—note that the length of the incision is the same as the equal sides of the isosceles triangle. (D) The isosceles triangle was sutured to the trunk with a 3-0 monofilament polypropylene suture.

Fig. 3 Postoperative imaging of a tailor-made tapering graft. Contrast-enhanced computed tomography 1 year after surgery. The tailor-made tapering graft was created from Hemashield Gold 18mm x 9mm. An arrowhead shows the proximal anastomosis. Note that there is no graft-related complication or thrombus, and both graft limbs were anastomosed to the normal-sized iliac arteries.

sis, a tapering graft was tailor-made on the back table. Using a transected portion of Hemashield Gold (Maquet, Rastatt, Germany), 18mm x 9mm (Fig. 2A), we created an acute isosceles triangle (Fig. 2B). The forefront of the trunk of the bifurcated graft was cut longitudinally of the same length as the equal side of the isosceles triangle (Fig. 2C). Each vertex of the triangle together with the part on the trunk of the main graft was held with curved Kelly forceps. Two separate running sutures were performed with 3-0 Surgipro™ (Covidien, Dublin, Ireland) for each edge, beginning from the acute vertex (Fig. 2D). After clamping, the left kidney was irrigated with cold Ringer's solutions. Reconstruction was performed using the tailor-made tapering bifurcated graft and left renal artery bypass. The proximal anastomosis was reinforced by an expanded polytetrafluoroethylene felt (Fig. 1B). The operation duration was 496 min, and the estimated blood loss was 1725 mL. His postoperative course was uneventful, with the highest serum creatinine levels at 1.40 mg/dL on Day 1. He was discharged on Day 6 and returned to his normal activities. Contrast-enhanced computed tomography taken 1 year after surgery indicated no problems at the site of anastomosis or the tapering graft (Fig. 3). At 3 years, he neither had renal impairment (serum creatinine level 0.79 mg/dL) nor anastomotic degeneration. He gave written informed consent for case presentation.

Discussion

A large aortic neck is not rare in patients with AAA. In real-world registries, proximal aortic diameters >25 mm and >30 mm were observed in 37.6% and 7.7% of patients/the population, respectively, both of which lead to higher risks of delayed type Ia endoleak after conventional endovascular aneurysm repair.^{1,2)} Considering the natural enlargement of ectatic aortas (26–29 mm) together with the factors of radial force and oversizing of stent grafts, this observation is legitimate, especially in the long term.^{5,6)} Therefore, open repair is a reasonable option for good-risk patients having AAA combined with a large neck.

The tailor-made tapering graft presented in the current report can be applied to any diameter at the anastomosis. Tips should include an adequate length of the trunk because an acute rather than obtuse isosceles triangle makes for a natural taper (Figs. 2B and 2D). We have employed this method for five other patients with large proximal aortas (aortic diameters at anastomosis ranging from 30 to 35 mm). We preferred using expanded polytetrafluoroethylene felts at the anastomoses. Compared to other pararenal open repairs with conventional grafts (n=22), tailor-made tapering grafts treated significantly larger aortas (3.2 ± 0.27 cm vs. 2.4 ± 0.47 cm, $p=.0007$)

and yet contributed to withholding renal ischemia (0 min vs. 42 ± 27 min, $p = .002$) and showing better trends in operation time (257 ± 33 min vs. 307 ± 69 min, $p = .13$), estimated blood loss (515 ± 173 mL vs. 806 ± 393 mL, $p = .12$), and postoperative acute kidney injury (0% vs. 36%, $p = .28$). During a mean nine-year follow-up, no graft-related problems were observed. One patient with an anastomotic dilatation (42 mm) had been observed non-operatively without any event. Our results were further supported by others' observations wherein reconstruction at the level just below the renal arteries performed better in the short term without increasing anastomotic degeneration at 3 years.⁷⁾

As described in the literature, there are two distinct types of arterial enlargements without reaching aneurysms: general arteriomegaly and local arteriomegaly (or aortomegaly).⁸⁾ Commercially available larger-size grafts (e.g., $24\text{ mm} \times 12\text{ mm}$ or $22\text{ mm} \times 11\text{ mm}$) may be used for the former cases. However, there are a few techniques available in the case of a large-neck aorta with a huge discrepancy in size from the distal arteries—composite grafts made of different sizes of prosthetic grafts or beveled grafts. Although we applied the inclusion technique for proximal anastomosis, any size discrepancy should be avoided as much as possible because it can lead to either leakage or wrinkles, which could potentially cause luminal narrowing of the renal arteries. Tailor-made tapering grafts can increase the graft diameter by up to 50% in ex-vivo experiments (Fig. 4). We also conform to grafts with a curved edge, which is estimated to increase the perimeter by up to 25%. However, widely used beveled grafts (e.g., 30 degrees) will increase the perimeter by up to 15%. Supposing Dacron grafts expand by 25% when implanted; 18 mm grafts could accommodate up to 34 mm, 28 mm, and 26 mm aortas in tapering, curved, and beveled grafts, respectively. Considering the contemporary concern regarding adequate experience and performance level among young vascular surgeons,^{9–11)} open surgical

techniques ensuring safe and effective outcomes are highly desirable.

Conclusion

The advantages of our tapering grafts lie in their flexibility to handle the widest ranges of aortic diameters and natural configurations without crushing the orifices of the renal arteries. Anastomosis at aneurysms should be avoided. In selected cases with a large-neck aorta such as aortomegaly, tailor-made tapering grafts contribute to decent outcomes both in the short term and midterm.

Disclosure Statement (COI)

The authors declare that there is no conflict of interest.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Ethical Approval

The ethics committee of Asahi General Hospital approved this study (registration number 2016071908).

Informed Consent

Written informed consent was obtained from all individual participants included in the study.

Author Contributions

Study conception: TS, TF

Data collection: all authors

Analysis: TS, MK, TK, KF

Investigation: all authors

Writing: TS

Funding acquisition: none

Critical review and revision: all authors

Final approval of the article: all authors

Accountability for all aspects of the work: all authors

	Tapering graft	Curved graft	Beveled graft
Graft perimeter	+50%	+25%	+15%
After implantation	~34mm	~28mm	~26mm
Configuration	natural	natural	unnatural

Fig. 4 Characteristics of different types of grafts used in large-neck aortas. The size after implantation takes +25% growth in size into consideration. The tapering graft handles the largest diameter range with natural configuration.

References

- 1) Oliveira NFG, Gonçalves FB, Ultee K, et al. Patients with large neck diameter have a higher risk of type IA endoleaks and aneurysm rupture after standard endovascular aneurysm repair. *J Vasc Surg* 2019; **69**: 783-91. doi: 10.1016/j.jvs.2018.07.021
- 2) Howard DPJ, Marron CD, Sideso E, et al. Editor's choice: influence of proximal aortic neck diameter on durability of aneurysm sealing and overall survival in patients undergoing

Shirasu T, et al.

- endovascular aneurysm repair. Real world data from the Gore Global Registry for Endovascular Aortic Treatment (GREAT). *Eur J Vasc Endovasc Surg* 2018; **56**: 189-99. doi: 10.1016/j.ejvs.2018.03.027
- 3) van den Ham LH, Holden A, Savlovskis J, et al. Editor's choice-occurrence and classification of proximal Type I endoleaks after endovascular aneurysm sealing using the Nellix™ device. *Eur J Vasc Endovasc Surg* 2017; **54**: 729-36. doi: 10.1016/j.ejvs.2017.09.016
 - 4) Doonan RJ, Girsowicz E, Dubois L, et al. A systematic review and meta-analysis of endovascular juxtarenal aortic aneurysm repair demonstrates lower perioperative mortality compared with open repair. *J Vasc Surg* 2019; **70**: 2054-64. e3. doi: 10.1016/j.jvs.2019.04.464
 - 5) Devaraj S, Dodds SR. Ultrasound surveillance of ectatic abdominal aortas. *Ann R Coll Surg Engl* 2008; **90**: 477-82. doi: 10.1308/003588408X301064
 - 6) Kret MR, Tran K, Lee JT. Change in aortic neck diameter after endovascular aortic aneurysm repair. *Ann Vasc Surg* 2017; **43**: 115-20. doi: 10.1016/j.avsg.2016.11.013
 - 7) Wang LJ, Tsougranis GH, Tanious A, et al. The removal of all proximal aneurysmal aortic tissue does not affect anastomotic degeneration after open juxtarenal aortic aneurysm repair. *J Vasc Surg* 2020; **71**: 390-9. doi: 10.1016/j.jvs.2019.02.072
 - 8) Callum KG, Lea Thomas M, Browse NL. A definition of arteriomegaly and the size of arteries supplying the lower limbs. *Br J Surg* 1983; **70**: 524-9. doi: 10.1002/bjs.1800700907
 - 9) Shirasu T, Furuya T, Nagai M, et al. Learning curve analysis to determine operative requirements for young vascular surgeons learning open abdominal aortic aneurysm repair. *Circ J* 2019; **83**: 1868-75. doi: 10.1253/circj.CJ-19-0386
 - 10) Varkevisser RRB, O'Donnell TFX, Swerdlow NJ, et al. Fenestrated endovascular aneurysm repair is associated with lower perioperative morbidity and mortality compared with open repair for complex abdominal aortic aneurysms. *J Vasc Surg* 2019; **69**: 1670-8. doi: 10.1016/j.jvs.2018.08.192
 - 11) Deery SE, O'Donnell TFX, Bodewes TCF, et al. Early reintervention after open and endovascular abdominal aortic aneurysm repair is associated with high mortality. *J Vasc Surg* 2018; **67**: 433-40.e1. doi: 10.1016/j.jvs.2017.06.104