S2-1

A novel pest control strategy: Targeting insect chitin degradation during moulting

Qing Yang, Lei Chen, Mingbo Qu

School of Bioscience and Biotechnology, Dalian University of Technology, Dalian, P. R. China

Insects possess a greater number of chitinases than any other organisms. Here is the first report of unliganded and oligosaccharide-complexed crystal structures of the insect chitinase OfChtI from *Ostrinia furnacalis*. OfChtI is essential to insect moulting. A structural comparison with other chitinases revealed that OfChtI contains a long substrate-binding cleft similar to the bacterial chitinase SmChiB from *Serratia marcescens*. However, unlike the exoacting SmChiB, which has a blocked and tunnel-like cleft, OfChtI possesses an open and groove-like cleft. The crystal structures reveal the unique structural characteristics of an insect chitinase. Since chitinase inhibitors have potential applications as pesticides, fungicides, and antiasthmatics. Here fully deacetylated chitooligosaccharides were determined to be inhibitors by thermodynamics and crystallography. Crystal structures reveal a competitive inhibition mode, first exploring fully deacetylated chitooligosaccharides acting as chitinase inhibitors and perhaps providing a structural basis for developing eco-friendly inhibitors against chitinases.

S2-2

Development, Ultrastructure and Morphology of Cuticle of a Beetle

Mi Young Noh and Yasuyuki Arakane

Department of Applied Biology, Chonnam National University, Gwangju 500-757, Korea

Insect cuticles, a biomaterial primarily formed from structural proteins and the polysaccharide chitin. Despite this limited compositional palette, cuticle has remarkably diverse properties, ranging from soft and flexible to hard and rigid. Chitin fibers are embedded in a matrix of protein molecules, which are cross-linked by catechol derivatives, forming a dual-network composite material. In a process called sclerotization (hardening and coloration), cuticular proteins (CPs) are cross-linked by quinones derived from the oxidation of catechols, and by formation of bonds between tyrosine side chains, resulting in hardening of the exoskeleton. Some CPs contain a conserved amino acid sequence known as the R&R motif, which has function as a chitin-binding region, coordinating interactions between the chitin fibers and protein network. The factors that lead to synthesis and assembly of cuticle with differing mechanical properties are not well understood.

In this study, we investigated development and differentiation of rigid cuticle using the red flour beetle, *Tribolium castaneum* adult, as a model insect. *T. castaneum* as a beetle is superior model for studying rigid cuticle formation because they have a highly modified (sclerotized and pigment) forewing (elytron), which can be separated from other tissues easily and cleanly. We analyzed ultrastructure of elytral cuticle during development by transmission electron microscopy (TEM). In 3 d-old pupae, pupal cuticle separated from the epidermal cells (apolysis), and the outermost envelop of adult cuticle was being formed. Protein-rich epicuticle and procuticle composed numbers of chitinous horizontal laminae and vertical oriented canals (pore canals) were formed at 4 and 5 d-old pupal stages (before molting to adult). Furthermore, additional thick horizontal chitinous laminae and brick-like layers were formed by 10 days after adult eclosion.

We further analyzed by immunogold-labeling TEM rigorous protein localization of several CPs that are highly abundant into the rigid cuticle of a beetle. We also discuss functional importance of chitin metabolism related genes such as chitinases and chitin deacetylases in insect molting, ultrastructure and organization of chitinous insect cuticle. These results will contribute understanding cuticle formation and differentiation in insect during post-embryonic development.

This work was supported by NRF (NRF-2012R1A2A1A01006467).