(J. H. Müller:—Journ. of Bact. VII. 300–325, 1922. Journ. of Biolog. chem. LVI. No 1. 1923.) The sulphur compound, isolated by the author from yeast, has entirely the same properties with it, except a little difference in the rotatory power. Although the constitution and the distribution of this compound are now in the course of investigation, it is clear that it must have been produced by autolysis of of yeast protein.

The author desires to express his sincere thanks to prof. Dr. U. Suzuki, for kind advice and giving the opportunity of publishing this paper.

Sinomenine and Dehydrosinomenine.

Part IV.

By KAKUJI GOTO.

In the last report (this Journal, this vol., p. 50), it could not be decided whether the amine, obtained from sinomenine by the fusion with kali was methylvinylamine or methylethylamine. For, the melting point of the gold salt coincided with that of the methylethylaminechloroaurate, but the chloroplatinate melted seventeen degrees higher than that of the methylethylamine, while the analysis of the both salts could scarcely show any difference in these two amines.

The decision could be given, if sinomenineiodomethylate was fused with kali. For, in this case, the amine to be liberated must naturally be tertiary and the ethanolamine, if it was produced at all, could not loose water intermolecularly.

In this supposition sinomenineiodomethylate was boiled with 66 per cent caustic kali in the same way as with sinomenine, and the sinomenol was obtained in 40 per cent yield of the theoretical. But the amine, isolated in this case was undoubtedly dimethylethylamine (m. p. of the chloroaurate 222° with decomposition, $\text{Au}=47.87$ per cent; calc. as $\text{C}_3\text{H}_{11}\text{NAuCl}_4\text{H}$, $\text{Au}=47.81$ percent and m.p. of the chloroplatinate 239° with decomposition, $\text{Pt}=35.38$ per cent; calc. as $(\text{C}_3\text{H}_{11}\text{N})_2\text{PtCl}_6\text{H}_2$ $\text{Pt}=35.22$ per cent.). From this fact, it may be inferred that the amine obtained in the kali fusion of sinominine must have been methylethylamine and not methylvinylamine. The fact that the m. p. of its chloroplatinate seventeen degrees higher than that given in the literature was perhaps one of examples often met with auri-

and platinichlorides of the lower fatty amines

The decomposition of sinomenine in the fusion with kali can, therefore, be expressed in the following scheme.

If so, sinomenol must be dioxydimethoxyphenanthrene ($C_{16}H_{14}O_4$) itself, and not its dihydride, as was supposed in the previous report. The comparison of the analytical data with the theoretical shows also clearly this fact. The molecular formulae of sinomenol and its derivatives, given in the same report, must, accordingly, be corrected as follows.

		Found		Calculated		Calculated on the basis sinomenol $C_{16}H_{14}O_4$	
		C	H	C	H	C	H
Sinomenol	$C_{16}H_{14}O_4$	70.71	5.83	71.08	5.22	70.58	5.89
Diacetyl-sinomenol.	$C_{20}H_{18}O_6$	1.67.45	5.54	67.79	5.12	67.41	5.62
		2.67.88	6.24				
		3.67.32	6.00				
Dibenzoyl-sinomenol.	$C_{30}H_{22}O_6$	1.75.59	5.39	75.28	4.62	75.00	5.00
		2.75.39	5.15				
Dibenzoyl-sinomenol- chinone.	$C_{30}H_{20}O_8$	69.42	4.07	70.86	3.96	—	—
Dimethyl-sinomenol,	$C_{18}H_{18}O_4$	A. 72.59	6.51	72.45	6.08	72.00	6.71
		B. 72.01	6.39				

Sinomenine was methylated by diazomethane in status nascendi. Methylsinomenine, thus obtained, melts at 169° and contains three methoxyl groups (26.51 per cent; calc. as three methoxyl groups in $C_{20}H_{25}NO_4$, 27.11 per cent). Its hydrochloride melts at 265° and is very sparingly soluble in water. Methylsinomenine gives neither diazoreaction nor ferric chloride and potassium ferriocyanide reactions. It gives a semicarbazone (m. p. $149-151^\circ$, N=14.54 per cent; calc. as $C_{21}H_{26}O_5N_5$, N=13.69 per cent.) so that the existence of the carbonyl and hydroxyl groups independently each other, could be shown on sinemenine itself.

Department of Chemotherapy, Kitasato Institute.

(Received, May 21st, 1925)