<i>d-l</i> -α amino- <i>n</i> -butyric acid	act. amylalc.	81.5	32.5	79	0.6(13.2 L)	—
<i>d-l</i> - <i>n</i> -valine	<i>n</i> -butylalc.	70.5	57	70	3.5(30)	49
<i>d-l</i> , valine	amylalc. B. P.(123~30°)	65	40~43	59	30.36(30)	40*
<i>d-l</i> , leucine	isoamylalc.	63	57.5	—	2.0(23)	38
asparagine	—	—	—	—	4.0(30)	84.5
glutaminic acid	amylalc.	—	—	—	24(30)	88
control	trace of amyl alc.				3.0(20)	54*

* together with added yeast.

On the Glycerin Fraction in saké.

By Masakazu YAMADA

(Received October 29, 1934)

(1) The glycerin fraction of saké (10 L) and moto-mash (50 L) were isolated by ordinary alcohol-ether extraction method from the extracts and 0.384% in saké and 0.754% in moto-mash were found. The fraction consists of 2·3-butylenglycol (B. P. 177~185°) and glycerin (B. P. 156~169°/7 mm). The ratio of two compounds are 1:20 in saké and 1:10 in moto mash.

(2) Diphenyl-carbamate of 2·3-butylenglycol prepared melts at 201° and contains nitrogen 8.61% (8.54% calculated for $C_{18}H_{20}N_2O_4$).

(3) Specific rotatory power of butylenglycol:

$$[\alpha]_D^{20} = +9.70$$

(4) Both constituents are necessary for the refined sweet taste of saké.

On the Variation of Sugar-content in saké.

By Masakazu YAMADA

Katsuichirō TAKAGISHI

(Governmental Institute of Brewing, Takinogawa, Tokyo.)

(Received October 29, 1934)

The sweet taste of saké is mainly due to sugar in it, while we found that sugar content was not constant but increased day by day with average speed of 0.0284% a day until pasteurization after the stage of compression of 'moromi.'*

* Moromi=saké lees.