Studies on Phenolic Lactones

Part V. General Route to Substituted β -Benzyl- γ -butyrolactones and an Alternate Synthesis of ($\pm$) Hibalactone

By Kyôhei YAMASHITA and Masanao MATSUI

Faculty of Agriculture, University of Tokyo

Received June 13, 1960

Substituted β -benzyl- γ -butyrolactones were synthesized. Hydrolysis and decarboxylation of ethyl α -allyl- α -benzyl malonate gave α -allyl- α -benzyl acetic acid. Esterification of the acid and subsequent reduction with lithium aluminum hydride afforded β -benzyl- β -allyl ethanol. Oxidation of the double bonds with potassium permanganate followed by periodate treatment, silver oxide oxidation and lactonization yielded β -benzyl butyrolactones. Thus β -benzyl-, β -piperonyl-, β -3,4,5-trimethoxybenzyl- butyrolactone were prepared.

Condensation of piperonal and β -piperonyl- γ -butyrolactone with sodium ethylate afforded ($\pm$) hibalactone.

The authors were interested in preparing substituted β -benzyl- γ -butyrolactones as intermediates for further synthetic work. It is a well known fact that the condensation of substituted malonic esters with ethylene oxide or

substituted β -benzyl butyrolactones.

The authors attempted several other routes for the synthesis of β -benzyl- γ -butyrolactones. For the first time the authors attempted the lactonization of γ -bromo butyronitriles as follows:

lactonization of γ -haloesters afford α -benzyl- γ -butyrolactones. For the synthesis of substituted β -benzyl- γ -butyrolactones, however, only a few reports have appeared in the literature. Reduction of β -substituted succinic acid β -mono thioalkyl ester with Raney nickel catalyst¹⁾ or hydroxymethylation of β -benzoyl propionic acid followed by catalytic hydrogenation²⁾ afford

But in the course of this route the reaction of one mole of sodium cyanide with one mole of dibromide afforded only a small amount of bromo-nitrile and the product was mainly dinitrile and recovered dibromide. Accordingly the yield of β -benzyl butyrolactone was very poor. So the authors devised an another scheme as follows:

1) S. Okumura, M. Masumura, T. Horie and S. Kori, *J. Soc. Org. Synth. Chem. Japan*, **17**, 461 (1951).

2) J. Rothe and H. Zimmer, *J. Org. Chem.*, **24**, 586 (1959).

Hydrolysis and decarboxylation of ethyl α -allyl- α -benzyl malonate gave α -allyl- α -benzyl

acetic acid. Esterification of the acid and subsequent reduction with lithium aluminum hydride afforded β -benzyl- β -allyl ethanol. Oxidation of the double bonds with potassium permanganate followed by periodate treatment, silver oxide oxidation and lactonization yielded β -benzyl butyrolactones. Of course, the ozonization of β -allyl- β -benzyl-ethanol was expected to yield the same results.

For the synthesis of some of the substituted β -benzyl- γ -butyrolactone reduction of β -benzoyl- γ -butyrolactones seemed to be superior method, however, β -piperonyl butyrolactone was more easily obtained by our method than that from β -piperonyl butyrolactone.

The authors attempted the condensation of piperonal with β -piperonyl- γ -butyrolactone. Although the condensation of piperonal with β -piperonyl-butylolactone by the use of potassium amide in liquid ammonia afforded hydroxylactone, the use of sodium ethylate or sodium hydride in benzene gave racemic hiballactone.

EXPERIMENTAL

β -Benzyl- γ -butyrolactone (General procedure)

a) Ethyl α -allyl- α -benzyl acetate

Ethyl allyl malonate (50 g, 0.25 mole) was alkylated with 35 g (0.28 mole) of benzyl chloride in the presence of 5.8 g (0.25 mole) of sodium in absolute ethanol. After refluxing for 2 hr. 30 g (0.54 mole) of potassium hydroxide in 100 ml of water was added and refluxed for an hour to saponify the esters. After the distillation of alcohol neutral substances were removed by ether extraction. An aqueous solution was acidified with hydrochloric acid and extracted with ether. After the evaporation of the ether the residual mass was heated to 170~180°C to decarboxylation. The crude acid was esterified by refluxing with ethanol and sulfuric acid for 2 hr. The neutral fraction was separated and distilled in vacuo. Ethyl α -allyl- α -benzyl acetate was distilled at 120~125°C, yielded 26.2 g. *Anal.* Found: C, 77.32; H, 8.06. Calcd. for $C_{14}H_{18}O_2$: C, 77.1; H, 8.2%.

b) β -Allyl- β -benzyl ethanol

Ethyl α -allyl- α -benzyl acetate, 26.2 g (0.12 mole), was reduced with 2.66 g (0.07 mole) of lithium aluminum

FIG. 1. Infrared Absorption Spectra of β -Benzyl- γ -butyrolactones.

- (I) β -Benzyl- γ -butyrolactone (film)
- (II) β -Piperonyl- γ -butyrolactone (film)
- (III) β -3,4,5-Trimethoxybenzyl- γ -butyrolactone (Nujol mull)

hydride. β -Allyl- β -benzyl ethanol was distilled at 120~121°C/6 mm, yielded 19.3 g. *Anal.* Found: C, 82.20; H, 9.06. Calcd. for $C_{12}H_{16}O$: C, 81.8; H, 9.1%.

c) β -Benzyl- γ -butyrolactone

A solution of 17.4 g (0.11 mole) of potassium permanganate in 350 ml of water was added to a cold solution of 19.3 g (0.11 mole) of β -allyl- β -benzyl ethanol in 100 ml of acetone at 0~5°C. After stirring for 2 hr. below 10°C and then 1 hour at 60°C, the reaction mixture was filtered to remove manganese dioxide. After the evaporation of acetone from the filtrate, 200 g of potassium carbonate was added. Crude triol was separated as a oily layer, weighted 16.9 g.

Crude 2-benzyl pentanetriol (1,4,5) was dissolved in 50 ml of water and was added to a solution of 18.3 g (0.08 mole) of periodic acid in 100 ml of water with stirring under cooling.

After stirring for an hour, the separated hydroxy-aldehyde was extracted with ethyl acetate. Evaporation of the solvent from the extract left 13.8 of crude β -benzyl- γ -hydroxybutyraldehyde. The crude aldehyde in 50 ml of warm ethanol was added to a suspension of silver oxide freshly prepared from 26.4 g of silver nitrate in 100 ml of 5% sodium hydroxide solution and warmed for 30 minutes at 80°C with vigorous stirring. The hot mixture was filtered and ethanol was distilled off. After the removal of neutral substances with chloroform extraction the aqueous solution was acidified with hydrochloric acid and warmed on water bath for 30 minutes. The reaction mixture was extracted with chloroform, and the extract was washed with sodium bicarbonate solution and water. After removal of solvent, the residual mass was distilled in vacuo. β -Benzyl- γ -butyrolactone was distilled at 165~166°C/6 mm, weighed 6.4 g. *Anal.* Found: C, 74.07; H, 6.93; Calcd. for $C_{11}H_{12}O_2$: C, 75.0; H, 6.8%. Infra-red spectrum showed a characteristic lactone band at 1740 cm^{-1} .

β -Piperonyl- γ -butyrolactone

Methyl α -allyl- α -piperonyl acetate was prepared from 50 g of allyl malonate and 48 g of piperonyl chloride in the similar manner as described in general procedure. It was distilled at 188°C/21 mm, yielded 32 g. *Anal.* Found: C, 67.81; H, 7.18. Calcd. for $C_{14}H_{16}O_4$: C, 67.8; H, 6.4%.

β -Allyl- β -piperonyl ethanol, b.p. 190°C/14 mm, was obtained by lithium aluminum hydride reduction of the

ester in 95% yield. *Anal.* Found: C, 70.75; H, 7.84. Calcd. for $C_{13}H_{16}O_3$: C, 70.9; H, 7.3%.

β -Piperonyl- γ -butyrolactone was obtained by the general procedure, b.p. 174~175°C/0.4 mm, weighed 7.4 g. *Anal.* Found: C, 64.91, H, 5.27. Calcd. for $C_{12}H_{12}O_4$: C, 65.5; H, 5.4%.

β -3,4,5-Trimethoxybenzyl- γ -butyrolactone

Methyl α -allyl- α -(3,4,5-trimethoxybenzyl)-acetate was prepared by the general procedure, b.p. 180~183°C/6 mm.

β -Allyl- β -(3,4,5-trimethoxybenzyl)-ethanol, b.p. 188~189°C/6 mm, was prepared by the reduction of the ester in 92% yield. *Anal.* Found: C, 67.48; H, 9.06. Calcd. for $C_{15}H_{22}O_4$: C, 67.7; H, 8.3%.

β -(3,4,5-Trimethoxybenzyl)- γ -butyrolactone was obtained by general procedure, b.p. 168~170°C/0.1 mm, m.p. 81~83°C (from methanol). *Anal.* Found: C, 63.04; H, 7.54. Calcd. $C_{14}H_{18}O_5$: C, 63.1; H, 6.8%.

($\pm$) Hibalactone

To a suspension of sodium ethylate prepared from 0.54 g of sodium in dry benzene, a solution of 4.0 g of β -piperonyl- γ -butyrolactone and 3.0 g of piperonal in benzene was added with external cooling. After stirring for 5 hr. at 0~5°C the reaction mixture was allowed to stand over night. After refluxing for 4 hr. 50 ml of water was added and refluxed for an hour. After cooling the organic layer was discarded and the aqueous solution was acidified with hydrochloric acid and warmed on water bath for 30 minutes. The mixture was extracted with ethyl acetate and the extract was washed with dilute sodium bicarbonate solution and with water, and the ethyl acetate distilled off. The residue was recrystallized from methanol. The colorless plate, m.p. 153~155°C, was obtained, weighed 1.0 g. Upon admixture with authentic ($\pm$) hibalactone the m.p. did not depressed. When sodium hydride was used instead of sodium ethylate, the same results were obtained.

Acknowledgements The authors are grateful to the members of the Analytical Laboratory of the Department of Agricultural Chemistry, University of Tokyo, for microanalytical data. They are also indebted to Mr. K. Aizawa for the measurement of infra-red spectrum.