Studies on *Streptomyces*.

Part I. *Streptomyces Pulveraceus* nov. sp., Producing New Antibiotics Zygomycin A and B

By MOTOO SHIBATA, Eiji HIGASHIDE, Toshihiko KANZAKI,
Hiroichi YAMAMOTO and Kōiti NAKAZAWA

Research Laboratories, Takeda Pharmaceutical Industries, Ltd., Osaka

Received September 15, 1960

Investigation was made on the mycological properties of a strain No. 45449 isolated from a sample of soil collected in Fukuchiyama. Since the antibiotics produced by the strain resembled hydroxymycin and paromomycin, the strain was compared with the hydroxymycin- and paromomycin-producing strains, *S. paucisporogenes* and *S. rimosus* forma *paromomycinus*, and as a result the strain No. 45449 was found to be different from the latter two strains. Among known strains, *S. flavogriseus* resembles the present strain, but they are different morphologically and in the kind of the antibiotics they produce. Thus, as the strain No. 45449 was found to be a new strain, it was named *S. pulveraceus* nov. sp. The antibiotics produced by the present strain are physiologically basic substances active against Gram-positive and negative bacteria and acid fast bacteria, and they are considered to belong to the neomycin-kanamycin group.

INTRODUCTION

Since the discovery of streptothricin and streptomycin, discovery of a new water-soluble and basic antibiotic having almost the same antibacterial activity as streptomycin but no delayed toxicity as streptothricin has long been desired. Although many basic antibiotics including neomycin were discovered meanwhile, they were not used practically because of their strong toxicity, but Umezawa et al.¹⁾ discovered at last in 1957 a new antibiotic, kanamycin, which satisfies the above conditions.

The authors have thereafter isolated a strain No. 45449 from a sample of soil collected in Fukuchiyama and found that it produces an antibiotic, zygomycin A²⁾, belonging to the

so-called neomycin-kanamycin group and a new basic antibiotic, zygomycin B²⁾, which is less toxic than streptomycin and belongs to the streptomycin group. While the present study was carrying out, two antibiotics, hydroxymycin³⁾ and paromomycin⁴⁾, which also seem to belong to the neomycin-kanamycin group and extremely resemble zygomycin A, were reported by other researchers. In the present paper there are reported the mycological properties and antibiotic activity of the strain No. 45449, and results of comparison of the strain with related strains, *S. paucisporogenes* and *S. rimosus* forma *paromomycinus*, which produce hydroxymycin and paromomycin respectively.

1) H. Umezawa et al., *J. Antibiotics Ser. A.*, **10**, 181 (1957).

2) H. Hitomi, S. Horii, T. Yamaguchi and A. Miyake, *ibid.*, under contribution.

3) G. Hagemann, G. Nominé and L. Pénasse, *Ann. Pharm. Franc.*, **16**, 585 (1958).

4) Parke, Davis & Company, Japanese patent 1958-6649 (1958. 8. 15).

EXPERIMENTAL

I. Mycological Properties

1) Morphological characteristics

The aerial mycelium generally develops well, and it is powdery and its color is drab-gray or light grayish olive. The sporophore forms spirals and the spore is spherical~ellipsoidal, $0.6\sim 0.8\ \mu \times 0.8\sim 1.6\ \mu$. The

spiral, its adhering state and the spore are shown in Fig. 1.

2) Cultural characteristics

The cultural characteristics of the original strain are shown in Table I. In order to select a strain with high potency, a mutant was produced by irradiation of the original strain with ultraviolet rays (referred to as yellow mutant). Differing from the original

TABLE I. CULTURAL CHARACTERISTICS OF *Streptomyces pulveraceus* nov. sp. (No. 45449)

Czapek's agar	Growth (G): Colorless, later faint brown. Aerial mycelium (A): Powdery, Light Drab (Rdg.* XLVI, 17''-b)~Light Grayish Olive (Rdg. XLVI, 21''-b). Soluble pigment (S): None.
Glucose Czapek's agar	Same as on Czapek's agar.
Glycerin Czapek's agar	Same as on Czapek's agar.
Glucose asparagine agar	(G): Orange (Rdg. III, 15)~Xanthin Orange (Rdg. III, 13-i). (A): Powdery, Light Grayish Olive. (S): None, after 3~4 weeks, becoming Ochraceous-Buff (Rdg. XV, 15'-b).
Nutrient agar (G):	Colorless, folded. (A): None. (S): None.
Glucose nutrient agar	(G): Colorless, wrinkled. (A): None or scant, White to Smoke Gray (Rdg. XLVI, 21''-d). (S): Brown.
Glycerin nutrient agar	(G): Wrinkled, Buckthorn Brown (Rdg. XV, 17'-i). (A): None or scant, white. (S): Brown.
Glucose broth	(G): Surface growth. (A): Smoke Gray. (S): Brown.
Starch agar	(G): Colorless~Yellow Ocher (Rdg. XV, 17'), penetrating deep into the medium. (A): Scant, Light Grayish Olive. (S): None.
Egg	(G): Hair Brown (Rdg. XLVI, 17''-i). (A): None.
Yeast extract agar	(G): Wrinkled, Chamois (Rdg. XXX, 19'-b)~Buckthorn Brown. (A): Scant, Light Grayish Olive. (S): Deep brown.
Potato plug	(G): Wrinkled, Pale Orange-Yellow (Rdg. III, 17-f), later Zinc Orange (Rdg. XV, 13'-o). (A): Powdery, Grayish Olive (Rdg. XLVI, 21''-b), Plug is blackend.
Carrot plug	(G): Colorless~faint brown. (A): Powdery, Light Grayish Olive, Plug is slightly brownish.
Milk	(G): Colorless, surface growth. (S): Brown. Peptonization without coagulation.
Nutrient gelatin	(G): Orange. (S): None. Liquefaction.
Nitrate reduction	Reduction, strong.
Löffler's serum	(G): Colorless. (A): None. (S): None.
Cellulose	No growth.
Calcium malate agar	(G): Yellow, penetrating into the medium. (A): None or scant, Smoke Gray. (S): None.
Tyrosinate agar	(G): Colorless~faint brown. (A): Light Grayish Olive. (S): None.
Starch hydrolysis	Hydrolysis, enzymatic zone/growth zone=65~68 mm/25~26 mm.

* Rdg: Ridgway, Color Standards and Color Nomenclature.

× 900

× 900

× 2600

× 5000

FIG. 1.

TABLE II. CULTURAL CHARACTERISTICS OF YELLOW-MUTANT

Czapek's agar	Growth (G): Colorless to Light Ochraceous-Salmon (Rdg. XV, 13'-d). Aerial mycelium (A): Light Buff (Rdg. XV, 17'-f) Soluble pigment (S): None.
Glucose Czapek's agar	Same as on Czapek's agar.
Glycerin Czapek's agar	(G): Colorless to Light Ochraceous-Salmon. (A): Powdery, Buff-Yellow (Rdg. IV, 19-d). (S): None.
Glucose asparagine agar	(G): Ochraceous-Orange (Rdg. XV, 15'). (A): Pale Yellow-Orange (Rdg. III, 15-f), powdery. (S): None later becoming faint brown.
Glucose broth	(G): Colorless sediment. (A): None. (S): None.
Egg	(G): Colorless, wrinkled. (A): None. (S): None.
Potato plug	(G): Orange (Rdg. III, 15). (A): White, Light Buff. Plug is blackend.
Carrot plug	(G): Orange (A): White, Light Buff. Plug is faint brown.
Milk	(G): Colorless ring. (S): Yellowish brown. Peptonization without coagulation.

TABLE III. UTILIZATION OF CARBON SOURCES

	<i>S. pulve-</i> <i>raceus</i>	Yellow- mutant	<i>S. rimosus</i> forma <i>par-</i> <i>omomycinus</i>	<i>S. rimosus</i> NRRL-2234
Arabinose	±	—	— ~ ±	≡
Rhamnose	≡	≡	—	—
Fructose	+	+	≡	≡
D-Galactose	≡	≡	≡	≡
Saccharose	±	±	—	—
D-Maltose	≡	≡	≡	≡
Lactose	≡	+ ~ ≡	+ ~ ≡	≡
Raffinose	≡	+ ~ ≡	± ~ +	± ~ +
Inuline	±	±	—	—
D-Mannitol	±	±	≡	≡
Sorbitol	+	±	≡	≡
Dulcitol	±	±	—	—
Inositol	±	±	≡	≡
Salicin	≡	≡	—	—
Na-acetate	+	+		
Na-citrate	≡	+		
Na-succinate	≡	+		
Erythritol	±			
D-Mannose	+		≡	≡
D-Xylose	≡	≡	— ~ ±	— ~ ±
Esculin	≡		—	— ~ ±
Dextran	≡			
Control	±	±		

Positive utilization: ≡ heavy growth
 ≡ good growth
 + fair growth
 No utilization: ± faint growth
 — no growth

strain, the aerial mycelium of the mutant is yellow, but other properties are almost the same as those of the original strain.

3) Utilizability of carbon source

Carbon source utilizability of the strain No. 45449 and yellow mutant was investigated by the method of Pridham et al.⁵⁾ In this case, the carbon source utilization of the paromomycin producing strains, *S. rimosus* forma *paromomycinus* and *S. rimosus*, was also referred to.

At present, the report of the properties of the hydroxymycin producing strain *S. paucisporogenes* and the paromomycin producing strain *S. rimosus* forma *paromomycinus* is very brief, and therefore, it is impossible to compare the strains with the strain No. 45449 in detail. But *S. paucisporogenes* forms no spiral, produces cracks, and does not reduce nitrate, whereas the strain No. 45449 forms spirals, produces no crack, and strongly reduces nitrate, and thus they are different morphologically as well as biochemically. On the other hand, *S. rimosus* forma *paromomycinus* forms white aerial mycelia on all the media described, produces cracks on its growth, and does not peptonize milk, while the strain No. 45449 forms drab-gray~grayish olive aerial mycelia, produces no crack on its growth, and well peptonizes milk, and in addition the

5) T.G. Pridham and D. Gottlieb, *J. Bact.*, **56**, 107 (1948).

two strains are different in carbon source utilizability as shown in Table III. *S. rimosus* NRRL-2234, which was reported to resemble *S. rimosus* forma *paromomycinus* in various properties such as morphological and cultural characteristics and the utilizability of carbon and nitrogen sources, except the utilization of arabinose, was really cultivated and compared with the strain No. 45449, finding that they are different. In addition to the above-mentioned differences, both *S. paucisporogenes* and *S. rimosus* forma *paromomycinus* produce no antibiotic belonging to the streptomycin group such as zygomycin B, and therefore, they are different from the strain No. 45449 in the kind of the antibiotics they produce.

Among known strains, *S. flavogriseus* appears to resemble the strain No. 45449, but they are different morphologically because it is described that the aerial mycelium of the former is long and straight hyphae producing a few curling tips, while the latter clearly forms spirals. Further, in view of the fact that the strain No. 45449 produces the two new antibiotics (zygomycin A and zygomycin B), it was judged to be a new strain and therefore, named *S. pulveraceus* nov. sp.

TABLE IV. ANTIBACTERIAL SPECTRUM OF *S. pulveraceus* nov. sp.

Test organism	Inhibition zone (mm)			
	Bouillon	agar	Glycerin bouillon	agar
<i>E. coli</i>	23	24	14	17
<i>P. vulgaris</i>	20	20	15	19
<i>B. subtilis</i>	29	30	22	23
<i>Staphy. aureus</i>	18	19	15	16
<i>B. cereus</i>	20	20	15	19
<i>Ps. aeruginosa</i>	0	0	—	—
<i>S. marcescens</i>	21	20	—	—
<i>Mycobact. 607</i>	—	—	20	22
<i>Mycobact. avium</i>	—	—	17	16
" (Streptomycin-resistant)	—	—	20	19

TABLE V. DIFFERENCE OF ANTIBACTERIAL POWER BY pH CHANGES

Test organism	Inhibition zone (mm)			
	pH 6.0-plate		pH 8.0-plate	
<i>B. Subtilis</i>	21	22	25	25
<i>Mycobact. avium</i> (S. M.-resistant)	30	29	37	35

II. Antibiotic Activity

The antibacterial spectrum of the strain No. 45449 was investigated by the cross-streak method using bouillon agar and glycerin bouillon agar. On the other hand, the strain was grown on agar-discs and the discs were put on plates having pH of 6.0 and 8.0 and containing *B. subtilis* and *Mycobact. avium* (streptomycin resistant), respectively, and incubated at 37°C for twenty hours and forty-four hours to investigate the change of the antibiotic activity (inhibition diameter) by the variation of pH. The results are shown in Table IV and V. According to the results the strain No. 45449 widely inhibits the growth of Gram-positive and negative bacteria and acid-fast bacteria, showing the so-called wide spectrum. And since the strain is more active in basic state, it seems to produce physiologically basic substances.

Known physiologically basic antibiotics which are active against Gram-positive and negative bacteria and acid-fast bacteria are the streptomycin, streptothricin, and neomycin groups. As the strain No. 45449 inhibits the growth of streptomycin-resistant *Mycobact. avium*, the antibiotics produced by the strain are not considered to belong to the streptomycin group, and since the activity of the strain against *B. cereus* is stronger than streptothricin group producing strains, the antibiotics produced by the strain more resemble neomycin than streptothricin. From the two features mentioned above, the antibiotics produced by the strain No. 45449 seem to belong to the neomycin group.

SUMMARY

Investigation was made on the mycological properties of a strain No. 45449 which was isolated from a sample of soil collected in Fukuchiyama. As the antibiotics produced by the strain resembled hydroxymycin and paromomycin, the strain was compared with the hydroxymycin and paromomycin-producing strains, i.e. *S. paucisporogenes* and *S. rimosus* forma *paromomycinus*, and as a result the strain No. 45449 was found to be a new species and therefore, named *S. pulveraceus* nov. sp. From the antibiotic properties of the strain, the antibiotics produced by the strain were assumed to belong to the neomycin-kanamycin group.

Acknowledgement The authors are grateful to Dr. Kikiti Sato and Dr. Satoru Kuwada for their advice and encouragement. Thanks are also due to Miss Tomiko Fudanoki for her cooperation.

[Agr. Biol. Chem., Vol. 25, No. 3, p. 176~180, 1961]

Studies on *Streptomyces*.

Part II. Difference between Antibiotic No. 45449-A (Zygomycin A) and other Antibiotics Belonging to the Neomycin Group.

By Motoo SHIBATA, Eiji HIGASHIDE, Toshihiko KANZAKI,
Hiroichi YAMAMOTO and Kōiti NAKAZAWA

Research Laboratories, Takeda Pharmaceutical Industries, Ltd., Osaka

Received September 14, 1960

Judging from its antibiotic spectrum and chemical properties, antibiotic No. 45449-A was found to be an antibiotic belonging to the neomycin group.

This antibiotic was compared with other antibiotics belonging to the neomycin group by the following three methods.

First, comparison of the antibiotic spectrum of No. 45449-A with those of neomycin, kanamycin hydroxymycin and paromomycin indicated that the former is different from the latter. Secondly, test for cross resistance using *St. pulveraceus* and *St. fradiae* as test microorganism showed that No. 45445-A is distinguishable from neomycin and kanamycin. Thirdly, investigation of diffusion assay ratios by the method of Fischer et al. also presumed No. 45449-A to be a new antibiotic differing from neomycin and kanamycin.

INTRODUCTION

As reported in the previous paper¹⁾, the antibiotic No. 45449-A (referred to as 45449 hereinafter) produced by *Streptomyces pulveraceus* nov. sp.¹⁾ is a basic substance and has antibiotic activity against Gram positive and negative bacteria, acid fast bacteria, and protozoa as seen from its antibiotic spectrum. The antibiotic shows cross resistance with neomycin but not with penicillin, streptomycin,

tetracycline, and chloromycetin²⁾. From this property and from the fact that the antibiotic is optically active (dextrorotatory) and is extracted from the culture broth in the same manner as streptomycin and neomycin³⁾, it was found to belong to the neomycin group.

Up to now the identity of an antibiotic with similar known antibiotics has been discussed from their antibiotic spectrum, sum-

1) M. Shibata, E. Higashide, T. Kanzaki, H. Yamamoto and K. Nakazawa, This Journal, **25**, 171 (1961).

2) T. Araki, Y. Oka and K. Tsuchiya, Annual Reports of the Takeda Research Laboratory, **19**, 131 (1960).

3) H. Hitomi, S. Horii, T. Yamaguchi and A. Miyake, *J. Antibiotics* in press.