L-Sorbose Oxidase from *Trametes sanguinea*

Sir:

In the previous papers, we reported on the two kinds of new enzymes concerning the formation and reduction of 5-keto-D-fructose (5-keto-L-sorbose); i.e. D-fructose dehydrogenase¹⁾ and 5-ketofructose reductase,²⁾ both of which were found in acetic acid bacteria.

While studying on the sugar metabolism of Basidiomycetes, we have found that L-sorbose was actively oxidized by the cell-free extracts of *Trametes sanguinea*, which, however, could not assimilate the ketose for growth.

The present communication concerns with the purification and some properties of the aerodehydrogenase obtained from the fungus which catalyzes the following reaction:

Trametes sanguinea IFO 4923 was reciprocally shaken at 27°C for 48 hours in the Sakaguchi flasks containing 100 ml of medium composed of koji juice and 0.3 per cent corn steep liquor. At the end of the cultivation small pellets were collected, washed, twice with cold saline, and suspended in M/10 phosphate buffer, pH 6.0, for the sonication for 15 minutes. The sonicated cells were centrifuged for 20 minutes at 3,500×g. The supernatant fluid was then ultracentrifuged for one hour at 105,000×g and to the resulting supernatant ammonium sulfate was added. The precipitate formed with 37~57 per cent saturation was collected and the procedure was repeated once again. This (ammonium sulfate precipitated) fraction was dissolved in M/10 phosphate buffer, pH 6.0, and the enzyme solution was dialyzed overnight against M/100 of the same buffer. The activity of the

enzyme preparation obtained with DEAE-cellulose column chromatography was about 30-fold of that of the crude extracts.

This preparation showed an intensive activity toward L-sorbose and the oxidation product was identified as 5-keto-D-fructose. In contrast, D-fructose was completely inactive as substrate. This enzyme specificity is noticeable, considering from the fact that D-fructose dehydrogenase obtained from acetic acid bacteria forms 5-keto-D-fructose from D-fructose but not from L-sorbose.¹⁾ Beside L-sorbose, D-glucose, D-galactose, D-xylose and maltose were oxidized by this partially purified enzyme preparation. (Table I) However, it is obscure whether or not this preparation is a mixture of several enzymes, since sugar-oxidizing enzymes of separate entity, to date, are not definitely clarified as yet.

The optimal pH was 6.0 and the range was rather broad. Michaelis constant for L-sorbose was 2.2×10^{-2} M/l at pH 6.0 and 30°C

TABLE I. SUBSTRATE SPECIFICITY

Substrates	Activity %
L-Sorbose	100
D-Fructose	0
D-Glucose	155
D-Mannose	0
D-Galactose	71
D-Xylose	42
D-Arabinose	0
L-Arabinose	0
Sorbitol	0
Mannitol	0
Erythritol	0
Sodium D-gluconate	0
Glucurono-γ-lactone	1
Sucrose	0
Maltose	15

Each reaction mixture (3.0 ml) contained 100 μmoles of substrate, 250 μmoles of phosphate buffer (pH 6.0), 0.2 ml of 1.0×10^{-8} M/l of 2,6-dichlorophenol indophenol, and 0.5 ml of the enzyme solution and half-decolorization time was measured.

1) Y. Yamada, K. Aida and T. Uemura, This Journal, **30**, 95 (1966); Y. Yamada, K. Aida and T. Uemura, Symposia on Enzyme Chemistry, Tokushima, 17th, 113 (1965).

2) K. Aida and Y. Yamada, This Journal, **28**, 74 (1964); K. Aida, Y. Yamada, and T. Uemura, Symposia on Enzyme Chemistry, Tokyo, 16th, 327 (1964).

in air. When catalase was added to the reaction mixture, the oxygen consumption was one half of that without catalase, in which the red-violet color was observed at the detection with KI-starch reaction, showing the presence of hydrogen peroxide. Also the enzyme could reduce 2,6-dichlorophenol indophenol, in the presence of which (6.7×10^{-4} M/l) Michaelis constant of the enzyme showed a rather higher value, 1.0×10^{-1} M/l under the same conditions.

The enzyme was not inhibited by 10^{-4} M/l *p*-chloromercuribenzoate, but silver and mercuric ions (10^{-4} M/l, respectively) abolished the enzyme activity. Inhibition by atabrine was not observed.

Recently, some of sugar oxidases such as hexose oxidase,³⁾ galactose oxidase^{4,5)} and xylose oxidase⁶⁾ have been precisely researched, but there is no description on the enzyme which oxidizes L-sorbose to 5-keto-D-fructose. Keilin and Hartree⁷⁾ reported that L-sorbose was not oxidized by their glucose oxidase (notatin) from *Penicillium*. Both lactose dehydrogenase from *Pseudomonas*⁸⁾ and glucose

dehydrogenase from *Aspergillus oryzae*^{9,10)} were inactive toward L-sorbose.

As for the chemical structure of dicarbonylhexose formed from D-fructose by acetic acid bacteria, two different ones, i.e. 6-ald-D-fructose¹¹⁾ and 5-keto-D-fructose¹²⁾ were independently proposed. Our present report which showed that dicarbonylhexose was formed only from L-sorbose but not from D-fructose may be another proof of the validity that it is 5-keto-D-fructose as confirmed by some authors.^{13,14)}

Yuzo YAMADA
Kazuaki IIZUKA
Kô AIDA
Teijiro UEMURA

The Institute of Applied Microbiology
The University of Tokyo, Tokyo

Received December 4, 1965

3) R. C. Bean, G. G. Porter and B. M. Steinberg, *J. Biol. Chem.*, **236**, 1235 (1961).

4) J. A. D. Cooper, W. Smith, M. Bacila and H. Medina, *J. Biol. Chem.*, **234**, 445 (1959).

5) G. Avigad, D. Amaral, C. Asensio and B. L. Horecker, *ibid.*, **237**, 2736 (1962).

6) H. Lyr, *Enzymologia*, **24**, 69 (1962).

7) D. Keilin and E. F. Hartree, *Biochem. J.*, **42**, 221 (1948).

8) Y. Nishizuka and O. Hayaishi, *J. Biol. Chem.*, **237**, 2721 (1962).

9) S. Boku and R. Sato, *Symposia on Enzyme Chemistry*, Osaka, 15th, 180 (1963); Personal Communication.

10) D. Ogura and M. Nagahisa, *Botanical Magazine, Tokyo*, **51**, 599 (1937).

11) R. Weidenhagen and G. Bernsee, *Chem. Ber.*, **93**, 2924 (1960).

12) O. Terada, S. Suzuki and S. Kinoshita, *This Journal*, **25**, 871 (1961); O. Terada, K. Tomizawa, and S. Kinoshita, *J. Agr. Chem. Soc. Japan*, **35**, 127 (1961); O. Terada, K. Tomizawa, S. Suzuki and S. Kinoshita, *ibid.*, 131; O. Terada, S. Suzuki and S. Kinoshita, *ibid.*, 178.

13) J. G. Carr, R. A. Coggins and G. C. Whiting, *Chemistry and Industry*, p. 1279 (1963).

14) G. Avigad and S. England, *J. Biol. Chem.*, **240**, 2290 (1965); S. England and G. Avigad, *ibid.*, 2297; S. England, G. and L. Proskey, *ibid.*, 2302.