Biochemical Studies on Ricin

Part IV. Amino Acid Analysis, Carboxyl and Amino Terminal Amino Acids of Ricin D

By Masatsune ISHIGURO, Gunki FUNATSU and Masaru FUNATSU

*Laboratory of Biochemistry, Faculty of Agriculture,
Kyushu University, Fukuoka, Japan*

Received October 7, 1970

Amino acid analysis of ricin D was carried out with an automatic amino acid analyzer. The following composition was determined: Asp₅₉Thr₃₅Ser₃₇Glu₄₆Pro₂₆Gly₃₆Ala₃₇Val₂₉Ile₃₅Leu₄₂Tyr₂₀Phe₁₈Lys₉His₆Arg₃₂Met₆1/2Cys₁₂Try₈ indicating that ricin D contains approximately 493 amino acid residues. The molecular weight of this protein was calculated to be about 55,000 agreeing well with the value obtained earlier by Archibald's method.

The C-terminal amino acids determined by carboxypeptidase A and by the hydrazinolysis technique were found to be serine (0.90 mole/mole) and phenylalanine (0.58 mole/mole).

The N-terminal amino acids measured by DNP-method of Sanger proved to be alanine (0.90 mole/mole) and isoleucine (0.52 mole/mole).

In an earlier paper,¹⁾ it was shown that ricin, a toxic protein of the castor bean, could be prepared in highly purified state by chromatographies on DEAE- and CM-cellulose. Moreover, after crystallization ricin appeared monodisperse on ultracentrifugation and electrophoresis. The crystalline ricin, designated as ricin D, was also found to be homogeneous on various biological assays exhibiting only toxicity and lacking hemagglutination, proteolytic activity and non-specific protein coagulation. The major physicochemical properties of this protein were reported subsequently.²⁾

The present paper deals with an amino acid composition, carboxyl and amino terminal amino acid determination of ricin D.

MATERIALS AND METHODS

Crystalline ricin D. Ricin D was prepared according to the method described previously.¹⁾ Twice re-

crystallized ricin D was used throughout this study except for the determination of N-terminal amino acid for which ricin D was prepared by the new method of Ishiguro, Funatsu and Funatsu³⁾ using gel filtration on Sephadex G-75.

Hydrazine was distilled according to the method of Kusama⁴⁾ just before use. Three times recrystallized and DFP-treated carboxypeptidase A purchased from Worthington Biochemical Corporation (Freehold, N. J., U.S.A.) was used after reprecipitation as described by Vallee *et al.*⁵⁾

Amino acid analysis. Aliquots of 1.459 mg of lyophilized ricin D were hydrolyzed with redistilled hydrochloric acid in evacuated, sealed tubes at 108°C for 23, 48, and 72 hr, respectively. The hydrolyzates were dried *in vacuo* to remove hydrochloric acid. The residues were dissolved in citrate buffer, pH 2.2, and analyzed with a Beckman automatic amino acid analyzer using columns of 150- and 15-cm lengths. Free cysteine was estimated amperometrically⁶⁾ and

3) M. Ishiguro, G. Funatsu and M. Funatsu, The preceding paper.

4) K. Kusama, *J. Biochem.*, **44**, 375 (1957).

5) B. J. Vallee *et al.*, *J. Biol. Chem.*, **235**, 64 (1960).

6) R. E. Benesch, H. H. Lardy and R. J. Benesch, *ibid.*, **216**, 663 (1955).

1) M. Ishiguro, T. Takahashi, G. Funatsu, K. Hayashi and M. Funatsu, *J. Biochem.*, **55**, 587 (1964).

2) M. Ishiguro, T. Takahashi, K. Hayashi and M. Funatsu, *ibid.*, **56**, 325 (1964).

spectrophotometrically.⁷⁾ Tryptophan was determined by the method of Goodwin and Morton.⁸⁾ Cystine and methionine were determined as cysteic acid and methionine sulfone after performic acid oxidation of ricin for 2.5 hr at $-7\sim-10^{\circ}\text{C}$ followed by acid hydrolysis.⁹⁾

Determination of C-terminal amino acid of ricin D.

A) *Hydrazinolysis.* Ricin D (3.0 mg) dried *in vacuo* over P_2O_5 was heated with 0.5 ml of anhydrous hydrazine in an evacuated, sealed tube at 108°C for 10 hr in presence and in absence of hydrazine sulfate. The hydrazinolysate was dried in an evacuated desiccator over conc. sulfuric acid for 24 hr. The residue was dissolved in 1.0 ml of citrate buffer, pH 2.2, without removal of the hydrazides with an aldehyde. After adjusting the pH to 2.2 with conc. hydrochloric acid the solution was analyzed with an amino acid analyzer.

B) *Digestion of ricin D with carboxypeptidase A.* About $0.5\ \mu\text{M}$ of ricin D was dissolved in 5.0 ml of 0.1 M sodium phosphate buffer, pH 8.5, and dialyzed against the distilled water and above mentioned buffer for 24 hr at 4°C to remove traces of amino acids and peptides. After removal of the insoluble material by centrifugation, the clear supernatant solution was divided equally into five test tubes. To each test tube, enzyme solution was added to give a weight ratio of 1/50 of enzyme to substrate. The reaction was carried out at 27 and 37°C for various periods of time. The solution of each test tube was applied onto the column of the automatic amino acid analyzer after solution was adjusted to pH 2.2.

Determination of N-terminal amino acid of ricin D. Determination of amino terminal amino acid of ricin D was performed according to the method of Sanger.¹⁰⁾ About 100 mg of ricin D was dinitrophenylated with 1-fluoro-2,4-dinitrobenzene (FDNB) in a 66% ethanol solution and in presence of 10% sodium bicarbonate for 2.5 hr at room temperature in the dark. After the excess FDNB had been removed with ether from the alkaline solution and then after adjustment of

pH to 5, the insoluble DNP-ricin was suspended in water and dialyzed against water in the dark overnight. Aliquots of the suspension of DNP-ricin were mixed with an equal volume of conc. hydrochloric acid, and placed in test tubes that were subsequently evacuated, sealed and heated at 107°C for 5, 8, 12, 24, 48, and 72 hr, respectively. After removal of hydrochloric acid, the liberated DNP-amino acids were then extracted with ether, dried *in vacuo*, dissolved in acetone, and after removing DNP-OH by sublimation with Mills' apparatus, applied to paper for two-dimensional chromatography. For the first chromatography *n*-butanol-0.1% ammonia¹¹⁾ and for the second one 1.5 M sodium phosphate buffer, pH 6.0¹²⁾ were utilized. The water-soluble DNP-amino acids were concentrated and desalted on a talcum column and identified by one-dimensional paper chromatography employing *tert*-amylalcohol-potassium biphthalate buffer, pH 6.0.¹³⁾ The amount of DNP-amino acids were calculated from optical density at $360\text{ m}\mu$ using extinction coefficients of 17.1, 17.7 and 17.0 for DNP-Ala, -Ile and -peptide, respectively. No correction was made for the loss of DNP-amino acid during acid hydrolysis and paper chromatographies. The DNP-amino acids detected on the paper were also identified subsequently as their free amino acids. After elution from the paper chromatogram DNP-amino acids were hydrolyzed with conc. ammonia¹⁴⁾ for 6 hr (for DNP-Ala) and 16 hr (for DNP-Ile) at 107°C in sealed tubes. The resulting free amino acids were determined with the automatic amino acid analyzer.

For the investigation of the DNP-compound 3 as seen in Fig. 1-a, representing presumably a DNP-peptide, about 60 mg of DNP-ricin D was hydrolyzed at 108°C for 16 hr with 5.7 N hydrochloric acid. The DNP-peptide was then separated from DNP-Ala and DNP-Ile by paper chromatography using the above mentioned solvent systems and hydrolyzed completely for 48 hr at 108°C with 5.7 N hydrochloric acid. The ether-soluble DNP-amino acids were subjected to the same two-dimensional paper chromatography. Free amino acids in the water phase were analyzed with the automatic amino acid analyzer.

7) P. D. Boyer, *J. Am. Chem. Soc.*, **76**, 4331 (1946).

8) T. W. Goodwin and R. A. Morton, *Biochem. J.*, **40**, 628 (1946).

9) C. H. W. Hirs, *J. Biol. Chem.*, **219**, 611 (1956).

10) F. Sanger, *Biochem. J.*, **45**, 563 (1949).

11) G. Braunitzer, *Chem. Ber.*, **88**, 2025 (1955).

12) A. L. Levy, *Nature*, **174**, 126 (1954).

13) S. Blackburn and A. G. Lowther, *Biochem. J.*, **48**, 126 (1959).

14) A. G. Lowther, *Nature*, **167**, 767 (1951).

RESULTS AND DISCUSSION

Amino acid analyses were performed by hydrolyzing 1.459 mg of ricin D for 23, 48 and 72 hr, respectively. The absolute concentrations of amino acids in ricin are deduced either by extrapolation to zero time hydrolysis or by selection of maximal recovery. The contents of cystine and methionine were determined after performic acid oxidation. Assuming the recovery of cysteic acid is 90 per cent on the oxidation according to Schram *et al.*,¹⁵⁾ the value of cysteic acid was corrected, whereas the observed recovery agreed well with the value obtained by the analysis of unoxidized ricin D. Free sulfhydryl group in ricin was repeatedly determined yielding the value of less than 0.3 residues per mole of ricin D both in the aqueous and 8M urea solutions. It was, therefore, concluded that all of the half cystine determined exist as disulfide and not sulfhydryl form in ricin D. Tryptophan was determined spectrophotometrically. It is interesting to note that 48 hr was required for complete release of phenylalanine although it has been known that the longer hydrolysis time is required for valine and isoleucine. This was explained by the presence of Ile-Phe- sequence in N-terminus of ricin D which could not completely hydrolyzed by 24 hr hydrolysis as described later.

The amino acid composition of ricin D as well as that of the ricin purified by CM-cellulose and Sephadex G-75 are shown in Table I. After calculation of minimal molecular weight, the nearest integral numbers for amino acid residues were obtained assuming a molecular weight of 60,000 for the protein. The total number of amino acid was given as 493. This figure is not definite, because of a low recovery of amino acid of 91.86 per cent due to a high content of carbohydrate in this protein. Of 493 residues, 105 are acidic and 47 basic amino acids. Since the isoelectric point of ricin D was found to

TABLE I. AMINO ACID COMPOSITIONS OF RICINS

Amino acid residues	Ricin ^{a)}	Ricin ^{b)} purified by CM-cellulose and Sephadex G-75
Aspartic acid	59	59.00
Threonine	35	34.43
Serine	37	32.90
Glutamic acid	46	45.59
Proline	26	25.91
Glycine	36	35.80
Alanine	37	36.78
Valine	29	27.41
Isoleucine	35	30.57
Leucine	42	42.09
Tyrosine	20	20.71
Phenylalanine	18	16.06
Lysine	9	9.05
Histidine	6	5.64
Arginine	32	32.53
Methionine ^{c)}	6	5.15
Cystine, 2 ^{c)}	12	10.58
Tryptophan ^{d)}	8	8.60
Amide ammonia	64	
Total	493	

^{a)} These values are obtained after appropriate corrections (see the text).

^{b)} The values are obtained after 24 hr hydrolysis and uncorrected one.

^{c)} Determined after oxidation.

^{d)} Determined spectrophotometrically.

be at pH 5.9, a relatively large number of amino acids would seem to be present as amide.

The amino acid composition of ricin preparation prepared by the new method³⁾ was also included in Table I (the last column). It was, thus, demonstrated that this preparation is identical with ricin D in amino acid composition, and this ricin preparation was used for the determination of amino terminal amino acid.

C-Terminal amino acids of ricin D.

A) *Hydrazinolysis.* Analysis of hydrazinolysate of ricin D is shown in Table II. Almost equimolar amounts of serine and

15) E. Schram *et al.*, *Biochem. J.*, **57**, 33 (1954).

TABLE II. C-TERMINAL AMINO ACID DETERMINATION

A. Hydrazinolysis

0.053 μ M of ricin D was hydrazinolized with 0.5 ml of hydrazine at 108°C for 10 hr in evacuated, sealed tube.

Amino acids	Without hydrazine sulfate	With hydrazine sulfate
Serine	0.35 mole/mole	0.33 mole/mole
Phenylalanine	0.42 mole/mole	0.38 mole/mole

B. Digestion with carboxypeptidase A

Ricin D (0.1 μ M) was digested with 1/50 of carboxypeptidase A (by weight) at pH 8.5 at different temperature and for different period of time.

Amino acids	Incubation time and temperature		
	1 hr, 27°C	10 hr, 27°C	1 hr, 37°C
Serine	0	0.90 mole/mole	1.52 mole/mole
Phenylalanine	0	trace	0.58 mole/mole
Glycine	0	0.58 mole/mole	0.76 mole/mole

phenylalanine were obtained from ricin D by hydrazinolysis both in the presence and in the absence of hydrazine sulfate, respectively.

B) Digestion of ricin D with carboxypeptidase

A. Analysis of free amino acids liberated by digestion of ricin D with carboxypeptidase A is also summarized in Table II. It was found by the treatment of ricin D with this enzyme that serine was released quantitatively at 27°C for 1 hr, and that the amount of this amino acid increased to 1.52 mole per mole of ricin D at 37°C for 1 hr. The release of phenylalanine, on the other hand, was relatively slow even at 37°C for one hour. This latter result does not necessarily imply that phenylalanine be not C-terminal amino acid of ricin D since the release of C-terminal amino acid from a protein by the action of carboxypeptidase A is also dependent on the penultimate amino acid residue of a protein in question.

From these two analyses, it was concluded that C-terminal amino acids of ricin D are serine and phenylalanine.*

N-Terminal amino acids. A typical paper chromatogram of ether-soluble DNP-amino acids from a 12 hr hydrolysate of ricin D is presented in Fig. 1-a. Three main spots (compound 1, 2 and 3) and two minor spots were observed. Of the three main compounds, compound 3 was assumed to be DNP-peptide since it disappeared after the prolonged hydrolysis as shown in Fig. 1-b. Compounds 1 and 2 were found to be DNP-alanine and DNP-leucine (or -isoleucine), respectively. This finding was confirmed by heating compound 1 for 6 hr and compound 2 for 16 hr

FIG. 1-a. A Schematic Representation of Paper Chromatogram of 12 hr Hydrolysate of DNP-Ricin D.

Ether-soluble DNP-amino acids were chromatographed on Toyo Roshi No. 51 developed first with *n*-butanol-0.1% ammonia (1:1) and then with 1.5 M phosphate buffer, pH 6.0. Three major DNP-compounds were found: compound-1, -2 and -3; in addition, three minor spots were observed.

* The qualitative analyses of C-terminal amino acids of ricin D has been recently carried out by tritium-labelling technique. The result that ^3H -phenylalanine was the only amino acid detected, thus, confirms the conclusion reached above considering the fact that this method at present is limited in that C-terminal serine is not identified.

FIG. 1-b. Liberation of DNP-Alanine, -Isoleucine and Unknown Compound.

●—● DNP-alanine; ●...● DNP-isoleucine;
●—● the unknown (DNP-peptide).

at 108°C in conc. ammonia and analyzing free amino acids liberated with the amino acid analyzer. Amino acids were observed at the positions of alanine for compound 1 and of isoleucine for compound 2, respectively.

As shown in Fig. 1-b, the yield of DNP-alanine, after only 5 hr of hydrolysis, was as high as 0.90 mole/mole, whereas DNP-isoleucine was liberated in rather small amount even after 48 hr of hydrolysis. This delay may be due to the formation of a compound intermediately, possibly DNP-peptide, which is represented by compound 3 as seen in Fig. 1-a. To obtain the latter compound, about 1 μ M of DNP-protein was hydrolyzed for 16 hr at 108°C. After DNP-peptide had been separated from DNP-alanine and -isoleucine by paper chromatography, DNP-peptide was further hydrolyzed. Ether-soluble DNP-amino acid was identified by two-dimensional paper chromatography to be DNP-leucine or -isoleucine. The regeneration of this DNP-amino acid into free amino acid followed by analysis

by the amino acid analyzer demonstrated this DNP-amino acid to be DNP-isoleucine. Analysis of the water phase by the same technique revealed the free amino acid of this peptide to be phenylalanine only. The latter two findings established that compound 3 indeed represents a DNP-peptide consisting of DNP-isoleucylphenylalanine. In addition, it was also shown that in the water phase of the original hydrolyzate of DNP-ricin D was free of DNP-arginine but contained the expected amount of ϵ -DNP-lysine.

In order to elucidate the reason why N-terminal alanine was easily split from DNP-ricin at the initial stage of acid hydrolysis, an attempt was made to isolate N-terminal peptide. Since DNP-ricin was insoluble in aqueous solution and not susceptible to proteolytic enzymes, it was treated with a 1:1 mixture of acetic acid and formic acid and subsequently dialyzed against water and then digested with Nagarse and Pronase at pH 8. The digest was acidified and extracted with ethylacetate. DNP-peptide extracted from the ethylacetate phase with 1% sodium bicarbonate solution was purified on paper. A DNP-peptide having $R_f=0.1$ consisted of DNP-alanine and aspartic acid. Therefore, alanyl-peptide of N-terminus of ricin D was assumed to be Al₂-Asp (or -Asn). The facts that this DNP-peptide is extractable in alkaline solution and that DNP-alanine is liberated almost completely by acid hydrolysis for 5 hr support the terminal sequence of NH₂-Ala-Asp-.

Acknowledgement. We wish to thank Dr. Karl Schmid, Boston University School of Medicine, for his help during the preparation of this manuscript.