Syntheses of Pentadienoic Acids Structurally Related to Absciscic Acid

Takayuki ORITANI, Takashi MATSUNAGA and Kyohei YAMASHITA

*Department of Agricultural Chemistry, Faculty of Agriculture,
Tohoku University, Sendai*

Received July 19, 1972

Instead of the 1'-hydroxy-4'-keto-2',6',6'-trimethyl-2'-cyclohexen-1'-yl group of absciscic acid, 3-methylpentadienoic acids with cyclohexenyl, methylcyclohexenyl, furyl, phenyl, hydroxyphenyl and hydroxy-methylphenyl groups at their 5-position were synthesized from simple alicyclic and aromatic aldehydes. Growth inhibitory activities of these compounds were examined.

In previous papers^{1,2)} the authors reported that 2-*cis*,4-*trans*-ionylideneacetic acids (I and II) have growth inhibitory activities similar to absciscic acid (III), and that the 2-*cis*,4-*trans*-3-methylpentadienoic acid system of the side-chain of I, II and III is essential for the appearance of their activities. Here, for the elucidation of the effect of the trimethylcyclohexene ring and the methyl group of I on the physiological activity, 3-methylpentadienoic acids substituted with simple alicyclic and aromatic rings at their 5-position were synthesized as follows.

The 2', 6', 6'-tridesmethyl acid (XII) of I (R=H) was synthesized by two methods (A and B) as shown in Scheme 1. Dehydration of the cyanohydrin of cyclohexanone (IV) with phosphorus oxychloride in pyridine gave 1-cyclohexenecarbonitrile (V).³⁾ Lithium aluminum hydride reduction of methyl 1-cyclohexenecarboxylate, obtained by alkaline hydrolysis and methylation of V, afforded 1-cyclohexenemethanol which was oxidized with active manganese dioxide to give 1-cyclohexenecarboxaldehyde (VI).⁴⁾ Aldol condensation of VI with acetone in the presence of sodium hydroxide gave the methyl ketone (VII'). Because VII' showed two peaks of UV absorption at λ_{max} 224.5 and 281 m μ in ethanol, the methyl ketone was considered

to be a mixture of positional isomers of the double bond as represented by the structure VII'. The reaction of VII' with carbethoxymethylenetriphenylphosphorane at 170°C afforded ethyl esters (VIII') consisting of a stereoisomeric mixture of 2-*cis* and *trans* forms (method A).

The reaction of the Grignard reagent of cyclohexyl bromide (IX) with ethyl orthoformate, followed by treatment with dilute sulfuric acid, afforded cyclohexanecarboxaldehyde (X).⁴⁾ Aldol condensation of X with acetone in the presence of sodium *tert*-butylate gave the methyl ketone (XI). XI was brominated with N-bromosuccinimide and then dehydrobrominated by heating with quinoline to give the dienone (VII).⁴⁾ The reaction of VII with carbethoxymethylenetriphenylphosphorane, followed by silica-gel column chromatography, gave the ethyl ester (VIII) of the 2-*cis* form. Alkaline hydrolysis of VIII afforded 2-*cis*,4-*trans*-5-(1'-cyclohexene-1'-yl)-3-methylpentadienoic acid (XII), mp 168~170°C (method B). In addition, the didesmethyl esters (XVIII and XXIII) of I (R=H) were synthesized by methods similar to those used for the above syntheses of XII. 6-Methyl-1-cyclohexenecarbonitrile (XIV),⁵⁾ prepared from 2-methylcyclohexanone (XIII), was hydrolyzed with potassium hydroxide and then methylated with diazomethane to yield methyl 6-methyl-1-cyclohexenecarboxylate, which was converted

Studies on Absciscic Acid (Part VII), see Reference 2).

SCHEME 1.

to the aldehyde (XV) by lithium aluminum hydride reduction and subsequent active manganese dioxide oxidation. Aldol condensation of XV with acetone afforded the methyl ketone (XVI).⁴⁾ The reaction of XVI with carbethoxymethylenetriphenylphosphorane gave the ethyl ester (XVII) of the 2-*cis* form, which was hydrolyzed with sodium hydroxide to give 2-*cis*,4-*trans*-(6'-methyl-1'-cyclohexen-1'-yl)-3-methylpentadienoic acid (XVIII), mp 141~142°C. Similarly, aldol condensation of 2-methylcyclohexanecarboxaldehyde (XX),⁶⁾ prepared from 2-methylcyclohexyl bromide (XIX), gave the methyl ketone (XXI). XXI was brominated with N-bromo-succinimide and then dehydrobrominated with

quinoline to give the dienone (XXII). As XXII showed olefinic methyl protons at 1.72 δ (singlet) and methyl protons at 1.03 δ (doublet) in the NMR spectrum, the obtained ketone is assumed to be a mixture of positional isomers of the double bond as represented by the structure XXII. The reaction of XXII with carbethoxymethylenetriphenylphosphorane, followed by separation with silica-gel column chromatography, gave the ethyl ester of the 2-*cis* form (XXIII).

In nature, some aromatic carotenoids are known: for example, 3-hydroxy-isoreniratenene (XXIV).⁷⁾ In order to estimate the physiological activities of the C-15 products probably produced by the cleavage of these and aromati-

zation of I, II and III, 3-methylpentadienoic acids with aromatic rings were synthesized as shown in Schemes 2 and 3. Reformatsky reaction of benzalacetone (XXV) with ethyl bromoacetate, followed by dehydration using a catalyst of *p*-toluenesulfonic acid, afforded the ethyl ester (XXVI) consisting of a stereoisomeric mixture of 2-*cis* and *trans* forms. XXVI was hydrolyzed with sodium hydroxide and then chromatographed on silica-gel. Elution with benzene afforded 2-*cis*,4-*trans*-5-phenyl-3-methylpentadienoic acid (XXVII),⁸⁾ mp 154~155°C. Aldol condensation of *p*- and *o*-hydroxy-benzaldehydes with acetone gave the methyl ketones (XXVIII) mp 108°C and (XXXII), mp 137°C, which were acetylated with acetic anhydride-pyridine to afford the acetates (XXIX), mp 76°C, and (XXXIII) as an oil. The same Wittig reaction of XXIX and XXXIII gave ethyl 2-*cis* and *trans*,4-*trans*-5-(*p*- and *o*-acetoxy-phenyl)-3-methylpentadienoates (XXX and XXXIV). Alkaline hydrolysis of XXX and XXXIV gave the pentadienoic acids (XXXI), mp 191~192°C

(2-*cis* form) and (XXXV), mp 184~185°C (2-*cis* and *trans* stereoisomeric mixture). Reimer-Tiemann reaction⁹⁾ of *m*-cresol with chloroform gave the hydroxy-methylbenzaldehydes (XXXVII, XXXVIII, XXXIX), which were converted to the methyl ketones (XL, XLI and XLII) by aldol condensation with acetone. These ketones were acetylated with acetic anhydride-pyridine and reacted with carbethoxymethylenetriphenylphosphorane to give ethyl 2-*cis* and *trans*,4-*trans*-5-(acetoxy-methylphenyl)-3-methylpentadienoates (XLIV, XLVII and L), which were hydrolyzed with sodium hydroxide to yield pentadienoic acids (XLV, XLVIII and LI) consisting of stereoisomeric mixtures of 2-*cis* and *trans* forms. Further, the same Wittig reaction of furfuralacetone (LII) afforded the ethyl ester (LIII). Alkaline hydrolysis of LIII gave 5-(2'-furyl)-3-methylpentadienoic acid (LIV),⁸⁾ mp 126~127°C, which consisted of 58% 2-*cis* form and 42% 2-*trans* form.

The physiological activities of the above compounds were tested on rice seedlings with

SCHEME 2.

SCHEME 3.

the results shown in Table I. The desmethyl acids (XII and XVIII) showed weak inhibition, but the desmethyl esters (VIII, XVII and XXIII) did not. The aromatic esters and acids (XXVI, XXVII, XXX, XXXIV and LIV) showed weak activities, but the phenolic acids (XXXI and XXXV) and the esters (XLIV and LIII) did not. The acid (XXVII) showed growth promoting action at a concentration of 10^{-4} mole/liter (Table II). The aromatic acetoxy-esters (XLVII and L) did not inhibit elongation strongly, but twisted stalks and leaves at random (depriving the plant of its normal tropism) at 10^{-3} – 10^{-4} mole/liter. The phenolic acids (XLV, XLVIII and LI) showed weak promoting action on stalks at 10^{-3} mole/liter. The authors think that the above data show the necessity for the methyl group of the rings of I and III for the appearance of marked physiological activity, and that some fission

TABLE I. GROWTH INHIBITORY ACTIVITIES OF PENTADIENOIC ACIDS ON RICE SEEDLINGS

Concn. (mole/liter) compd. No.	2×10^{-3}	10^{-3}	10^{-4}	10^{-5}
XII	—	1.1 ^{a)}	2.2	2.2
XVIII	— ^{b)}	1.2	1.6	2.2
XXVII	—	1.9	2.6	2.3
XXVI	—	1.5	2.3	2.2
XXX	—	0.5	2.3	2.3
XXIV	—	2.2	2.3	2.3
XLV	2.3	2.5	2.2	2.2
XLVIII	2.5	2.6	2.1	2.1
LI	2.4	2.5	2.2	2.1
LIV	—	1.5	2.1	2.2
I(R=H) ¹⁾	—	—	1.3	1.6
III (ABA) ¹⁾	—	—	—	—
Control	2.2			

^{a)} Mean length (cm) of the second leaf sheath of Honen-Wase (an ordinary variety of *Oryzae sativa*) after growing for 7 days at 28°C.

^{b)} Complete inhibition.

TABLE II. PHYSIOLOGICAL ACTIVITIES OF 2-*cis*,4-*trans*-5-PHENYL-3-METHYLPENTADIENOIC ACID (XXVII) ON RICE SEEDLINGS

The length (cm) of different organs of the seedlings were measured after growing for 7 days at 28°C.

Length (cm) Concn. (mole/liter)	Plant height	Second leaf sheath	Root
2×10^{-3}	0.2	0	0
10^{-3}	5.0	1.9	0.7
10^{-4}	8.2	2.6	8.1
10^{-5}	5.3	2.3	6.5
Control	5.1	2.2	5.7

products of aromatic carotenoids may have certain physiological activity.

EXPERIMENTAL

Melting points and boiling points were uncorrected. IR spectra were recorded with a Model IR-E of Japan Spectroscopic Co., Ltd. NMR spectra were recorded at 60 MHz with TMS as an internal standard (Model JMN-MH-60-II of Japan Electron Optics Laboratory Co., Ltd). Gas-liquid chromatography was carried out on a Yanagimoto Model GCG-550T.

1) 1-Cyclohexenecarboxaldehyde (VI). To a stirred solution of 63 g of cyanohydrin of IV in 200 ml of pyridine was gradually added 77 g of POCl₃ at 5~10°C. After stirring at room temperature for 2 hr and then at 50~55°C for 1 hr, the ordinary treatment of the reaction mixture gave 1-cyclohexene-carbonitrile (V),³⁾ bp 97°C/28 mmHg, $n_D^{15} = 1.4838$ (yield 98%). V was hydrolyzed with ethanolic potassium hydroxide under reflux and the acid obtained was methylated with diazomethane in ether to give methyl 1-cyclohexenecarboxylate, bp 97~100°C/23 mmHg, $n_D^{17} = 1.4725$. Reduction of the ester with excess LiAlH₄ in ether at room temperature, followed by treatment with dilute sulfuric acid, gave 1-cyclohexenemethanol, bp 110~114°C/35 mmHg, $n_D^{14} = 1.4868$. The alcohol (8 g) was shaken with 100 g of active MnO₂ in 100 ml of chloroform at room temperature for 2 days to give 6 g of 1-cyclohexenecarboxaldehyde (VI),⁴⁾ bp 100°C/50 mmHg, $n_D^{16} = 1.4917$.

2) 4-Cyclohexyl-3-buten-2-one (XI). To a stirred solution of Grignard reagent, prepared from 100 g of IX and 15.5 g of Mg in 200 ml of dry ether, was added dropwise 90 g of ethyl orthoformate. The solution was refluxed for 6 hr and evaporated on a water bath. The residual mixture was treated with cold dilute hydrochloric acid and extracted with ether. The ether extract was evaporated and the residue was steam-

distilled in the presence of dilute sulfuric acid to give cyclohexanecarboxaldehyde (X),⁴⁾ bp 63°C/24 mmHg. A mixture of 14 g of X, 50 ml of acetone and 2 g of sodium *tert*-butoxide was refluxed under nitrogen for 10 hr. The reaction mixture was poured into cold water and extracted with ether. The ether extract was washed with water and dried over anhyd. MgSO₄. Evaporation of the ether gave the methyl ketone (XI), bp 112~115°C/18 mmHg, $n_D^{27} = 1.4808$. IR $\nu_{\text{max}}^{\text{film}}$ cm⁻¹: 1670, 1610, 1355, 1250, 1175, 970. Gas chromatographic analysis using a 1.5 m $\times$ 3 mm column packed with 30% silicone SE30 on celite 545, column temp: 152°C isothermal, He flow rate: 20 ml/min (1.2 kg/cm²), retention time: 5.7 min (98% purity).

3) 4-(1'-Cyclohexen-1'-yl)-3-buten-2-one (VII)⁴⁾

Method A. To a stirred solution of 6 g of VI in 100 ml of acetone was added 40 ml of 5% aqueous sodium hydroxide at 5°C under nitrogen. After stirring for 6 hr, the reaction mixture was extracted with ether. The ether extract was washed with water and dried over anhyd. MgSO₄. Evaporation of the ether gave 5.5 g of the methyl ketone (VII'), bp 131~135°C/16 mmHg, $n_D^{14} = 1.5442$. IR $\nu_{\text{max}}^{\text{film}}$ cm⁻¹: 1670, 1620, 1594, 1357, 1250, 976. UV $\lambda_{\text{max}}^{\text{EtOH}}$ m μ : 224.5 and 281. Gas chromatographic analysis using the same column as in 2), column temp: 152°C isothermal, He flow rate: 15 ml/min (0.98 kg/cm²), retention time: 5.1 min (with an overlapping shoulder).

Method B. A mixture of 4.0 g of XI and 4.5 g of N-bromosuccinimide in 30 ml of carbontetrachloride was stirred at 50~55°C for 2 hr. After cooling, the reaction mixture was filtered and the filtrate was evaporated under reduced pressure on a water bath. The residual oil was heated with 20 ml of quinoline under occasional shaking for 1 hr. The reaction mixture was poured into cold water and extracted with ether. The ether extract was washed with dilute sulfuric acid and water and dried over anhyd. MgSO₄. Evaporation of the ether gave an oil, which was chromatographed on alumina. Elution with benzene-ether (4:1) afforded 2.3 g of VII. IR $\nu_{\text{max}}^{\text{film}}$ cm⁻¹: 1660, 1620, 1595, 1355, 1250, 1170, 975, 798. UV $\lambda_{\text{max}}^{\text{EtOH}}$ m μ : 280. Gas chromatographic analysis using the same column and column conditions as in 3), retention time: 9.1 min (95% purity).

4) Ethyl 2-*cis*,4-*trans*-5-(1'-cyclohexen-1'-yl)-3-methylpentadienoate (XII)

Method A. A mixture of 1.0 g of VII' and 2.5 g of carbethoxymethylenetriphenylphosphorane was heated at 170~180°C under nitrogen for 1 hr. The reaction mixture was treated with *n*-hexane-ether to separate triphenylphosphineoxide. The residual oil was chromatographed on alumina. Elution with benzene afforded 1.0 g of the ethyl ester (VIII') as an

oil. IR $\nu_{\max}^{\text{film}}$ cm^{-1} : 3090, 1715, 1610, 1220, 1150, 1055, 970, 860. UV $\lambda_{\max}^{\text{EtOH}}$ $m\mu$: 296.

Method B. The same Wittig reaction of VII with carbethoxymethylenetriphenylphosphorane gave the ethyl esters as a stereoisomeric mixture of 2-*cis* and *trans* forms. Gas chromatographic analysis using a 1.5 m $\times$ 3 mm column packed with 10% silicone SE30 on chromosorb-G (AW-DMCS), column temp: 224°C isothermal, He flow rate: 20 ml/min (1.48 kg/cm²), retention time: 2.5 min (67% 2-*cis*-ester (VIII)), 2.9 min (33% 2-*trans*-ester). The esters were chromatographed on silica-gel. Elution with benzene afforded 2-*cis* ester (VIII) as an oil. IR $\nu_{\max}^{\text{film}}$ cm^{-1} : 3090, 1715, 1610, 1590, 1375, 1280, 1220, 1150, 970, 848. UV $\lambda_{\max}^{\text{EtOH}}$ $m\mu$: 307.5. NMR (CCl_4) δ : 1.30 (3H, *t*, $J=7$ cps), 2.05 (3H, *s*, 2-*cis*-3-CH₃), 4.12 (2H, *q*, $J=7$ cps), 5.55 (1H, *s*), 5.87 (1H, *s*), 6.45 (1H, *d*, $J=16$ cps), 7.63 (1H, *d*, $J=16$ cps). VIII was hydrolyzed with 5% ethanolic sodium hydroxide at room temperature to give the acid (XII), mp 168~169°C (unstable) from acetonitrile.

5) 4-(6'-Methyl-1'-cyclohexen-1'-yl)-3-buten-2-one (XVI). 6-Methyl-1-cyclohexenecarbonitrile (XIV)⁵ (21 g), bp 90~92°C/18 mmHg, $n_D^{16} = 1.4806$, prepared by dehydration of the cyanohydrin of XIII with POCl₃ in pyridine, was hydrolyzed with 15% ethanolic sodium hydroxide under reflux for 5 hr. By the ordinary treatment the acid obtained was methylated with diazomethane in ether to give 12 g of methyl 6-methyl-1-cyclohexenecarboxylate, bp 94~100°C/20 mmHg, $n_D^{17} = 1.4712$. Reduction of 11.2 g of the ester with excess LiAlH₄ (3.7 g) in 150 ml of dry ether at room temperature, followed by treatment with dilute hydrochloric acid, gave 6.8 g of 6-methyl-1-cyclohexenemethanol, bp 90~100°C/20 mmHg, $n_D^{12} = 1.488$, which was shaken with active MnO₂ (80 g) for 3 days to give 3.5 g of the aldehyde (XV), bp 85~90°C/21 mmHg. To a stirred solution of 3.1 g of XV in 50 ml of acetone was added 15 ml of 2% aqueous sodium hydroxide, and the solution was stirred for 5 hr at room temperature. The reaction solution was poured into water and extracted with ether. Evaporation of the ether gave the methyl ketone (XVI) as an oil, bp 139~142°C/20 mmHg, $n_D^{13} = 1.5448$, which was chromatographed on silica-gel. Elution with benzene afforded the relatively pure ketone (XVI) (2.5 g). IR $\nu_{\max}^{\text{film}}$ cm^{-1} : 1670, 1622, 1600, 1360, 1250, 1170, 975. UV $\lambda_{\max}^{\text{EtOH}}$ $m\mu$: 281.5 ($E_{1\%}^{1\text{cm}} = 1.734$). NMR (CCl_4) δ : 1.05 (3H, *s*), 5.83 (1H, *d*, $J=16$ cps), 6.90 (1H, *d*, $J=16$ cps). Gas chromatographic analysis using the same column as in 3), column temp: 173°C isothermal, He flow rate: 19 ml/min, retention time: 6.4 min (97% purity).

6) 2-*cis*, 4-*trans*-5-(6'-Methyl-1'-cyclohexen-1'-yl)-3-methylpentadienoic acid (XVIII). A mixture of

1.0 g of XVI and carbethoxymethylenetriphenylphosphorane (3.0 g) was heated at 170~175°C for 1 hr and the mixture was treated with *n*-hexane-ether to separate triphenylphosphineoxide. The oil obtained was chromatographed on silica-gel. Elution with benzene gave 300 mg of XVII as an oil. NMR (CDCl_3) δ : 1.08 (3H, *d*), 1.33 (3H, *t*), 2.08 (3H, *s*), 4.28 (2H, *q*), 5.68 (1H, *s*), 6.00 (1H, *s*), 6.61 (1H, *d*, $J=16$ cps), 7.75 (1H, *d*, $J=16$ cps). Alkaline hydrolysis of XVII with 5% ethanolic sodium hydroxide at room temperature for 12 hr afforded the acid (XVIII), mp 141~142°C from acetonitrile. NMR (CDCl_3): 1.07 (3H, *d*), 5.70 (1H, *s*), 2.12 (3H, *s*), 6.03 (1H, *s*), 6.65 (1H, *d*, $J=16$ cps), 7.68 (1H, *d*, $J=16$ cps).

7) 4-(2'-Methyl-1'-cyclohexen-1'-yl)-3-buten-2-one (XXII). a) 4-(2'-Methylcyclohexenyl)-3-buten-2-one (XXI). To a stirred solution of Grignard reagent, prepared from 75 g of XIX and 11 g of Mg in 250 ml of dry ether, was added 80 g of ethyl orthoformate. The mixture was refluxed for 6 hr and evaporated on a water bath. The residual mixture was treated with dilute hydrochloric acid and extracted with ether. After evaporation of the ether, the residual oil was steam-distilled in the presence of dilute sulfuric acid to give 2-methylcyclohexanecarboxaldehyde (XX),⁶ bp 53~60°C/18 mmHg, $n_D^{19} = 1.4532$. A mixture of 9 g of XX and 2 g of sodium *tert*-butoxide in 50 ml of acetone was stirred at room temperature for 12 hr and then under reflux for 4 hr. The reaction mixture was poured into water and extracted with ether. The ether extract was washed with water and dried over anhyd. MgSO₄. Evaporation of the ether gave 4.5 g of XXI, bp 125~135°C/17 mmHg, $n_D^{19} = 1.4790$. IR $\nu_{\max}^{\text{film}}$ cm^{-1} : 1680, 1620, 1250, 980.

b) The methyl ketone (XXII). By the same experimental procedure as used in 3) Method B, XXI was brominated with N-bromosuccinimide in carbon tetrachloride and dehydrobrominated with quinoline to give the ketone (XXII), which was purified by silica-gel column chromatography. Elution with benzene-ether (4:1) gave the relatively pure ketone (XXII) as an oil. NMR (CCl_4) δ : 1.05 (*d*, about 20% of 6'-CHCH₃), 1.72 (*s*, about 80% of 2'-C-CH₃), 2.20 (3H, *s*), 5.90 (1H, *d*, $J=16$ cps), 6.96 (1H, *d*, $J=16$ cps). Gas chromatographic analysis using the same column as in 2), column temp: 152°C isothermal, He flow rate: 16 ml/min (1.5 kg/cm²), retention time: 13.1 min (80% purity).

8) Ethyl 2-*cis*, 4-*trans*-5-(2'-methyl-1'-cyclohexen-1'-yl)-3-methylpentadienoate (XXIII). By the same experimental procedure as in 4), the Wittig reaction of 1.0 g of XXII with carbethoxymethylenetriphenylphosphorane gave 900 mg of the crude esters, which were chromatographed on silica-gel. Elution with benzene gave the ester (XXIII) as an oil. IR $\nu_{\max}^{\text{film}}$ cm^{-1} :

3080, 1720, 1630, 1610, 1240, 1220, 1155, 1050, 970. NMR (CCl₄) δ : 1.07 (*d*, about 20% 6'-CHCH₃), 1.27 (3H, *t*), 1.73 (*s*, about 80% 2'-C-CH₃), 2.00 (3H, *s*, 2-*cis*-3-CH₃), 4.10 (2H, *q*), 5.63 (1H, *s*), 6.45 (1H, *d*), 7.68 (1H, *d*).

9) 2-*cis*, 4-*trans*-5-Phenyl-3-methylpentadienoic acid (XXVII).⁸⁾ A mixture of 50 g of XXV, 60 g of ethyl bromoacetate, 23 g of mossy zinc and 100 ml of dry benzene was refluxed for 1 hr. After cooling, the mixture was decomposed with 10% aqueous acetic acid and extracted with benzene. The benzene layer was washed successively with water and aqueous NaHCO₃ and dried over anhyd. MgSO₄. After evaporation of the solvent, the residual oil was refluxed with 1.5 g of *p*-toluenesulfonic acid in 100 ml of benzene with continuous separation of the water formed for 40 min. The reaction solution was washed with aqueous NaHCO₃ and dried over anhyd. MgSO₄. Evaporation of the solvent gave 35 g of ethyl 5-phenyl-3-methylpentadienoate (XXVI), bp 180~195°C/15 mmHg. n_D^{18} = 1.6040. IR $\nu_{\max}^{\text{film}}$ cm⁻¹: 1715, 1605, 1500, 1450, 1360, 1220, 1200, 1150, 1048, 960, 870, 840. XXVI was hydrolyzed with 10% ethanolic sodium hydroxide under reflux for 1 hr to give a stereoisomeric mixture of consisting of 2-*cis* and *trans* forms, mp 125~126°C from acetonitrile. NMR (CDCl₃) δ : 2.08 (*s*, 47% 2-*cis*-3-CH₃), 2.35 (*s*, 53% 2-*trans*-3-CH₃). The acid was chromatographed on silica-gel. Elution with benzene afforded the 2-*cis*,4-*trans*-acid (XXVII), mp 154~155°C from *n*-hexene-benzene. NMR (CDCl₃) δ : 2.18 (3H, *s*, 2-*cis*-3-CH₃), 5.78 (1H, *s*, 2-*cis*- =CH-), 6.93 (1H, *d*, *J*=16 cps, 5=CH-), 7.35 (5H, *m*), 8.30 (1H, *d*, *J*=16 cps, 4=CH-). Anal. Found: C, 76.71; H, 6.37. Calcd. for C₁₂H₁₂O₂: C, 76.57; H, 6.43%. Subsequent elution with benzene gave 2-*trans*,4-*trans*-5-phenyl-3-methylpentadienoic acid,⁸⁾ mp 160°C.

10) 4-(*p*- and *o*-Acetoxy-phenyl)-3-buten-2-ones (XXIX and XXXIII). a) *p*-Acetoxy-ketone (XXIX). To a stirred mixture of 50 g of *p*-hydroxybenzaldehyde, 300 ml of acetone and 200 ml of water was added dropwise 100 ml of 25% aqueous sodium hydroxide at 10°C, with stirring. The reaction mixture was allowed to stand overnight at room temperature. After acidification with dilute hydrochloric acid, the reaction mixture was extracted with ether. The ether extract was washed with water and dried over anhyd. MgSO₄. Evaporation of the ether gave 56 g of *p*-hydroxybenzalacetone (XXVIII), mp 108°C from benzene. Acetylation of XXVIII with acetic anhydride-pyridine at room temperature for 12 hr gave *p*-acetoxybenzalacetone (XXIX), mp 76~77°C from *n*-hexane-benzene. IR $\nu_{\max}^{\text{film}}$ cm⁻¹: 1755, 1650, 1600, 1510, 1420, 1375, 1270, 1200, 1180, 1020, 1005, 930, 890, 870.

b) *o*-Acetoxy-benzalacetone (XXXIII). By using

the same experimental procedure as in a), *o*-hydroxybenzaldehyde was reacted with acetone to yield *o*-hydroxybenzalacetone (XXXII), mp 137~138°C from ethyl acetate. XXXII was treated with acetic anhydride-pyridine to give *o*-acetoxybenzalacetone (XXXIII) as an oil. IR $\nu_{\max}^{\text{film}}$ cm⁻¹: 1770, 1670, 1610, 1490, 1360, 1250, 1200, 1175, 1090, 1005, 970, 910, 840, 800, 750.

11) Ethyl 2-*cis* and *trans*, 4-*trans*-5-(*p*- and *o*-acetoxy-phenyl)-3-methylpentadienoates (XXX and XXXIV)

a) *p*-Acetoxy-ester (XXX). A solution of 2 g of XXIX and 4 g of carbethoxymethylenetriphenylphosphorane in 50 ml of dry benzene was refluxed for 24 hr. After evaporation of the benzene, the residual oil was treated with *n*-hexane-ether to separate triphenylphosphineoxide. The product was chromatographed on alumina. Elution with benzene afforded *p*-acetoxy-ester (XXX) as an oil. IR $\nu_{\max}^{\text{film}}$ cm⁻¹: 1765, 1715, 1615, 1600, 1510, 1370, 1200, 1150, 1050, 1010, 970, 910, 830. NMR (CDCl₃) δ : 1.27 (3H, *t*), 2.03 (*s*, 2-*cis*-3-CH₃), 2.20 (3H, *s*, OCOCH₃), 2.33 (*s*, 2-*trans*-3-CH₃), 4.07 (2H, *q*), 5.60 (*s*, 66% 2-*cis*- =CH-), 5.72 (*s*, 34% 2-*trans*- =CH-). XXX was hydrolyzed with 5% ethanolic sodium hydroxide at room temperature to give the *p*-hydroxy-2-*cis*-acid (XXXI), mp 191~192°C (decomp) from acetonitrile. NMR (CD₃COCD₃) δ : 2.12 (3H, *s*), 5.65 (1H, *s*), 6.73 (2H, *d*, *J*=9 cps), 6.83 (1H, *d*, *J*=16 cps), 7.30 (2H, *d*, *J*=9 cps), 8.17 (1H, *d*, *J*=16 cps).

b) *o*-Acetoxy-ester (XXXIV). Using the same experimental condition as in a) the Wittig reaction of 2.0 g of XXXIII gave 2.5 g of the *o*-acetoxy-ester (XXXIV) as an oil. IR $\nu_{\max}^{\text{film}}$ cm⁻¹: 1765, 1710, 1610, 1490, 1360, 1200, 1175, 1155, 1090, 1045, 1010, 970, 910, 840, 810, 755. NMR (CDCl₃) δ : 1.27 (3H, *t*), 2.05 (*s*, 2-*cis*-3-CH₃), 2.28 (OCOCH₃, 2-*trans*-3-CH₃), 4.07 (2H, *q*), 5.62 (*s*, 53% 2-*cis*- =CH-), 5.75 (*s*, 47% 2-*trans*- =CH-), 6.97 (4H). Alkaline hydrolysis of XXXIV with 5% ethanolic sodium hydroxide gave the *o*-hydroxy-acid (XXXV), mp 184~185°C (decomp) from acetonitrile. NMR (CD₃COCD₃) δ : 2.18 (*s*, 73% 2-*cis*-3-CH₃), 2.45 (*s*, 27% 2-*trans*-3-CH₃), 5.77 (*s*), 5.95 (*s*), 6.93 (2H), 7.50 (1H), 8.37 (*d*, *J*=16 cps).

12) 4-(Hydroxy-methylphenyl)-3-buten-2-ones (XL, XLI and XLII). Reimer-Tiemann reaction⁹⁾ of *m*-cresol with chloroform using aqueous sodium hydroxide as a basic catalyst, followed by steam-distillation and formation of the sodium bisulfite adduct, gave 4-hydroxy-2-methylbenzaldehyde (XXXVII), mp 110°C and 2-hydroxy-6-methylbenzaldehyde (XXXVIII), mp 31~32°C and 2-hydroxy-4-methylbenzaldehyde (XXXIX), mp 59~60°C. Using the same experimental procedure as in 10), the aldehydes were condensed with acetone to give XL, mp 133~134°C from ethyl

acetate, XLI, mp 143~144°C from ethyl acetate, and XLII, mp 128~129°C from benzene, respectively. XL, *Anal.* Found: C, 74.84; H, 6.55. XLI, *Anal.* Found: C, 75.04; H, 6.69. XLII, *Anal.* Found: C, 75.06; H, 6.54. Calcd for $C_{11}H_{12}O_2$: C, 74.97; H, 6.86%.

13) *Ethyl 2-cis and trans,4-trans-5-(acetoxy-methyl-phenyl)-3-methylpentadienoates (XLIV, XLVII and L).* The hydroxy-ketones (XL, XLI and XLII) were acetylated with acetic anhydride-pyridine at room temperature for 12 hr to give the acetoxy-ketones (XLIII, XLVI and IL), respectively. XLIII, mp 62~63°C. IR $\nu_{\max}^{\text{film}}$ cm^{-1} : 1770, 1670, 1600, 1500, 1375, 1200, 1020, 905, 820. XLVI: IR $\nu_{\max}^{\text{film}}$ cm^{-1} : 1770, 1670, 1610, 1465, 1375, 1190, 1030, 990, 880, 790, 750. The acetoxy-ketones (XLIII, XLVI and IL) (each 1.0 g) were reacted with carbethoxymethylenetriphenylphosphorane (3.0 g) at 170~180°C for 1 hr. The reaction mixture was treated in each case with *n*-hexane-ether to separate triphenylphosphineoxide. The products obtained were chromatographed on alumina. Elution with benzene afforded the acetoxy-esters (XLIV, XLVII and L) as oils (1.1~1.2 g) respectively. XLIV: IR $\nu_{\max}^{\text{film}}$ cm^{-1} : 1770, 1715, 1600, 1498, 1440, 1365, 1205, 1153, 1050, 960, 900. XLVII: IR $\nu_{\max}^{\text{film}}$ cm^{-1} : 1765, 1715, 1604, 1465, 1365, 1190, 1152, 1030, 970, 870, 785. L: IR $\nu_{\max}^{\text{film}}$ cm^{-1} : 1770, 1715, 1608, 1505, 1445, 1365, 1200, 1153, 1050, 970, 810. XLIV: NMR (CCl_4) δ : 1.28 (3H, *t*), 2.10 (*s*), 2.20 (3H), 2.37 (*s*), 4.12 (2H, *q*), 5.63 (*s*, 40% 2-*cis*=CH-), 5.78 (*s*, 60% 2-*trans*=CH-), 6.83 (1H), 7.38 (2H), 8.21 (*s*, *J*=16 cps). XLVII: NMR (CCl_4) δ : 1.28 (3H, *t*), 2.10 (*s*), 2.15 (*s*), 2.23 (*s*), 2.35 (*s*), 4.12 (2H, *q*), 5.65 (*s*, 44% 2-*cis*=CH-), 5.73 (*s*, 56% 2-*trans*=CH-), 5.98 (3H), 8.12 (*d*, *J*=16 cps). L: NMR (CCl_4) δ : 1.28 (3H, *t*), 2.10 (*s*), 2.30 (*s*), 2.37 (*s*), 4.10 (2H, *q*), 5.60 (*s*, 43% 2-*cis*=CH-), 5.70 (*s*, 57% 2-*trans*=CH-), 6.70 (2H), 7.33 (1H), 8.22 (*d*, *J*=16 cps). The above acetoxy-esters were hydrolyzed with 5% ethanolic sodium hydroxide at room temperature to give the hydroxy-acids (XLV, XLVIII and LI). XLV, mp 212~215°C (from acetonitrile). NMR (CD_3COCD_3) δ : 2.07 (3H), 2.10 (*s*, 56% 2-*cis*-3- CH_3), 2.45 (*s*, 44% 2-*trans*-3- CH_3). XLVIII, mp 205~208°C from acetonitrile. LI, mp 224~227°C from acetonitrile. NMR (CD_3COCD_3) δ : 2.07 (3H), 2.10 (*s*, 77% 2-*cis*-3- CH_3), 2.38 (*s*, 23% 2-*trans*-3- CH_3). XLVIII: *Anal.* Found: C, 71.40; H, 6.60. Calcd for $C_{13}H_{14}O_3$: C, 71.54; H, 6.47%.

14) *2-cis and trans,4-trans-5-(2'-Furyl)-3-methylpentadienoic acids (LIV).*⁸⁾ The same Wittig reaction of LII with carbethoxymethylenetriphenylphosphorane at 140°C for 1 hr, followed by alumina column chromatography, gave the ethyl ester (LIII) as an oil. IR $\nu_{\max}^{\text{film}}$ cm^{-1} : 1715, 1610, 1255, 1230, 1150, 1045, 1010, 960, 925, 880, 835, 800, 735. LIII was hydrolyzed with 5% ethanolic sodium hydroxide at room temperature to give the acid (LIV) consisting of a stereoisomeric mixture of 2-*cis* and *trans* forms, mp 126~127°C from acetonitrile. NMR (CDCl_3) δ : 2.12 (*s*, 44% 2-*cis*-3- CH_3), 2.37 (*s*, 56% 2-*trans*-3- CH_3), 5.73 (*s*), 5.90 (*s*), 6.41 (2H), 6.73 (1H), 7.40 (1H), 8.13 (*d*, *J*=16 cps).

15) *Physiological activities on rice seedlings.* As described in a previous report,¹⁾ the physiological activities of the synthesized compounds were tested using rice seedlings (Honen-Wase, an ordinary variety of *Oryza sativa*). The results are shown in Tables I and II.

Acknowledgements. The authors wish to express their thanks to the members of the Analytical Laboratory of the Department of Chemistry, Tohoku University, for microanalysis and also to Miss. Y. Koseki, the Department of Agricultural Chemistry, Tohoku University, for measurement of the 60 Mc NMR spectra.

REFERENCES

- 1) T. Oritani and K. Yamashita, *Agr. Biol. Chem.*, **34**, 108 (1970).
- 2) *idem, ibid.*, **36**, 362 (1972).
- 3) E. J. Boormann and R. P. Linstead, *J. Chem. Soc.*, **1935**, 258.
- 4) I. Heibron, E. R. H. Jones, R. W. Richardson and F. Sandheimer, *ibid.*, **1949**, 737.
- 5) W. S. Papson and R. G. Shuttleworth, *ibid.*, **1940**, 636.
- 6) O. Diels and K. Alder, *Ann.*, **470**, 62 (1929).
- 7) F. Arcamone, B. Camerino, E. Cotta, G. Franseschi, A. Grein, A. Penco and C. Spala, *Experientia*, **25**, 241 (1969).
- 8) M. Matsui, K. Yamashita, M. Miyano, S. Kitamura, S. Okano, A. Kobayashi, T. Sato and R. Mikami, *The Journal of Vitaminology*, **4**, 190 (1958); G. Pattenden and B. C. L. Weedon, *J. Chem. Soc.*, **1968**, 1998.
- 9) F. Tiemann and C. Schotten, *Ber.*, **11**, 767 (1878).