Short Communication

A New Enzyme “Bilirubin Oxidase”
Produced by *Myrothecium*
verrucaria MT-1

Sawao MURAO and Noriaki TANAKA

*Department of Agricultural Chemistry,
College of Agriculture,
University of Osaka Prefecture,
Sakai, Osaka 591, Japan*

Received May 20, 1981

Bilirubin, consisted of tetrapyrrole, is the product of physiological heme degradation and the principal pigment of bile. So, if the reactive enzyme to bilirubin could be found, it may be expected to decolorize urine and feces and to be an analytical tool in the clinical fields. Recently we have succeeded in isolating a strain producing potent enzyme which is reactive to bilirubin. This strain was identified as *Myrothecium verrucaria* MT-1.

The enzyme activity was assayed as follow; 2.0 ml of bilirubin dissolved in 0.2 M Tris-HCl buffer containing 1 mM EDTA, pH 8.4 (20 µg/ml), 0.2 ml of enzyme solution were incubated at 37°C. One unit was defined as the equivalent amount of enzyme which reduced a ΔA at 440 nm of 1.05/min.

For the production of the enzyme, the strain was aerobically cultured at 25°C in a jar fermentor. A medium was potato extract containing 1% glucose (pH 5.8). After 72 hr cultivation, the culture broth was filtrated to re-

move mycelium. The filtrate was brought to 80% saturation with ammonium sulfate and resulting precipitate was dissolved in water and dialyzed against 10 mM Na₂CO₃-NaHCO₃ buffer (pH 9.2). The enzyme solution was decolorized by the treatment of charcoal powder twice [0.7% (w/v) pH 8.8 and 0.2% (w/v) pH 7.4]. This supernatant was adsorbed on a QAE-Sephadex A-50 column equilibrated with 15 mM Na₂CO₃-NaHCO₃ buffer (pH 9.3). Then the enzyme was eluted with a linear gradient of Na₂CO₃-NaHCO₃ buffer with a concentration from 0.015 M to 0.8 M. Active fractions were pooled and concentrated on a Diaflo PM-10 membrane filter and applied to a column of Sephadex G-100,

FIG. 1. Polyacrylamide Gel Disc Electrophoresis of Bilirubin Oxidase.

Electrophoresis was carried out using 7% gel of pH 9.5. The protein (about 30 µg) was stained with Coomassie Brilliant Blue.

TABLE I. PURIFICATION OF BILIRUBIN OXIDASE FROM *Myrothecium verrucaria* MT-1

Step	Total protein (A280)	Total activity (units)	Specific activity (units/A280)	Yield (%)
Culture filtrate	31500	57100	1.81	100
AmSO ₄	4670	32200	6.90	56
Charcoal treatment 1	1160	29500	25.4	52
Charcoal treatment 2	765	27600	36.1	48
QAE-Sephadex A-50	111	12600	114	22
Sephadex G-100	48	9900	206	17

equilibrated with 30 mM Na_2CO_3 - NaHCO_3 buffer (pH 9.4). The specific activity of the final preparation was 206 units/A280. The final enzyme preparation was submitted to polyacrylamide gel disc electrophoresis at pH 9.5. A single band was obtained, as shown in Fig. 1. A summary of the purification was presented in Table I.

The general property of the enzyme thus obtained was as follow: the molecular weight was estimated to be about 52,000 by gel filtration method on Sephadex G-100. The isoelectric point was determined to be pH 4.1 by means of isoelectric focusing. The enzyme had a maximum activity at pH 8.0. The highest reaction rate was observed at 40°C. In the absence of oxygen, the enzyme was inactive. In the enzyme, NAD, NADP, FAD and FMN were not detected. It turned out that the copper was contained by the atomic absorption spectroscopy. In reaction mixtures from this enzyme and bilirubin, the substance with light absorption maxima at 375 and 660~670 nm corresponding to biliverdin was contained. So, this enzyme was designated as

bilirubin oxidase. The hydrogen peroxide in reaction mixtures was measured by using peroxidase, 4-aminoantipyrine, and phenol,¹⁾ but was not detected.

The presence of the bilirubin oxidase activity was expected in the brain of guinea pig, but the pure enzyme was not obtained.²⁾ Therefore, the property of bilirubin oxidase was not clear. But we purified bilirubin oxidase produced by microorganism, and found to be homogeneous by disc-gel electrophoresis.

Bilirubin oxidase was effective to decolorize urine and feces and also expected to be an analytical tool in the clinical fields.

More detailed investigations on bilirubin oxidase will be published elsewhere.

REFERENCES

- 1) C. C. Allain, L. S. Poon, C. S. G. Chan, W. Richmond and P. C. Fu, *Clin. Chem.*, **20**, 470 (1974).
- 2) R. Brodersen and P. Bartels, *Eur. J. Biochem.*, **10**, 468 (1969).