Note

Reduction of α,β -Unsaturated Carbonyl Compounds by Some ActinomycetesAlain KERGOMARD, Michel F. RENARD
and Henri VESCHAMBRELaboratoire de Chimie Organique Biologique,
E.R.A. 392 du CNRS,
Université de Clermont II,
B.P. 45, 63170 Aubière, France

Received June 6, 1984

In previous communications, we have reported that eucaryote organisms (six fungi)¹⁾ and procaryotes (a *Clostridium* and three actinomycetes)²⁾ reduced the carbon-carbon double bond of α,β -unsaturated carbonyl compounds, in particular variously substituted cyclohexenones.

The stereochemical characteristics of this reaction have been described.³⁾

The purpose of this work is to demonstrate that this reaction is widespread among actinomycetes. We report here detailed results concerning 20 other actinomycetes.

MATERIAL AND METHODS

The preparation of 2-methyl-2-cyclohexen-1-one has already been described.^{1,3)} 2-Cyclohexen-1-one was of commercial grade. Some strains have been described in previous papers: *Streptomyces griseus* LAB 2142,⁴⁾ *S. albus* LAB 3840,⁵⁾ *S. griseorubiginosus* LAB 4915.⁶⁾ The natural rubber degrading strains have been described elsewhere: *S. phaeofaciens*, *S. griseoincarnatus*, *S. anulatus*, *S. griseolus*, *Nocardia asteroides* COB 911 and *N. asteroides* COB 906.⁷⁾

Medium 1. *S. albus* sub. sp. NRRL B1865 was grown in the following medium: glucose (10 g), yeast extract (1 g), meat extract (1 g) and Difco casaminoacids (4 g) in 1 liter of tap water. The pH was adjusted to 7.2~7.5 by 10 N NaOH.

Medium 2. *S. griseorubiginosus* LAB 4915 was grown in the following medium: sucrose (40 g), $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (0.1 g), $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ (0.1 g), NaCl (2 g), corn steep solid (15 g), CaCO_3 (2 g) and K_2HPO_4 (0.75 g) in 1 liter of tap

water. The pH was adjusted to 7.5 with 10 N NaOH.

Medium 3. *S. albus* LAB 3840 was grown in the following medium: glucose (40 g), corn steep solid (10 g), $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (0.5 g) and CaCO_3 (5 g) in 1 liter of tap water. The pH was adjusted to 7.4~7.5 with 10 N NaOH.

Medium 4. *S. cinnamomensis* ATCC 15413 was grown in the following medium: glucose (30 g), soyabean meal (5 g), CaCO_3 (1 g), KCl (0.1 g), K_2HPO_4 (0.1 g), $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ (0.6 g) and $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (0.3 g) in 1 liter of tap water. The pH was adjusted to 8 with 5 N KOH.

Medium 5. *S. antibioticus* NRRL 8167 was grown in the following medium: glucose (10 g), beet molasses (20 g), soyabean meal (5 g), CaCO_3 (2 g) and yeast extract (1 g) in 1 liter of tap water. The pH was adjusted to 7.2 by 10 N NaOH.

Medium 6. *S. aureofaciens* NRRL 8092 was grown in the following medium: dextrin (80 g), Difco casaminoacids (6 g), $(\text{NH}_4)_2\text{SO}_4$ (1 g), beet molasses (15 g), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (0.5 g) and CaCO_3 (2 g) in 1 liter of tap water. The pH was adjusted to 6.8~7 with 10 N NaOH.

Medium 7. *S. griseus* LAB 2142 was grown in the following medium: sucrose (30 g), soyabean meal (10 g), peptone (4 g), K_2HPO_4 (1 g) and CaCO_3 (5 g) in 1 liter of tap water. The pH was adjusted to 7 with 10 N NaOH.

Medium 8. *S. griseus* ATCC 10137 was grown in the following medium: glucose (20 g), soyabean meal (5 g), yeast extract (5 g), K_2HPO_4 (5 g) and NaCl (5 g) in 1 liter of tap water. The pH was adjusted to 7 with 10 N NaOH.

Medium 9. *Nocardia rubra* ATCC 15906 was grown in the following medium: glucose (10 g), Difco casaminoacids (4 g), yeast extract (1 g), meat extract (4 g) and NaCl (2.5 g) in 1 liter of tap water. The pH was adjusted to 7~7.2 with 10 N NaOH.

Experiments were carried out in 500 ml conical flasks containing 100 ml of each culture medium. After inoculation, the flasks were placed on a rotary shaker at 27°C and 200 rpm for 24 hr.

Fifty mg of substrate were then added to each flask, the agitation being continued for 48 hr more.

After filtration, the filtrate was saturated with ammonium sulfate and extracted three times with ether. After evaporation of the solvent, the residue was analyzed by GPC (stainless steel column, 6 ft $\times$ 0.125 inch, packed with 20% carbowax on chromosorb W, oven temperature 100°C, hydrogen pressure 1 Bar).

The crude products were purified by column chromatography with Merck 60 silicagel and pentane-ether (90:10) as the eluant.

Optical rotations. The optical activity was determined using a Perkin-Elmer 141 polarimeter at 25°C at the wavelength of the mercury J line ($\lambda = 578 \text{ nm}$) in a CHCl_3 solution.

RESULTS

The results obtained with 20 strains of actinomycetes

TABLE I. REDUCTION OF 2-CYCLOHEXEN-1-ONE AND 2-METHYL-2-CYCLOHEXEN-1-ONE BY VARIOUS STRAINS OF ACTINOMYCETES (YIELD IN PERCENT AFTER 48 hr OF REACTION)

Micro-organisms	Culture medium							$[\alpha]_D^{25}$	$[\alpha]_D^{25}$
<i>Streptomyces albus</i> sub. sp.									
NRRL B1476	1	0	53	47			No reaction		
<i>S. albus</i> sub. sp. NRRL B1477	1	0	56	44	27	52	−15.1°	21	+22.5°
<i>S. albus</i> sub. sp. NRRL B1478	1	0	49	51	35	34	−15.2°	31	+22.5°
<i>S. albus</i> sub. sp. NRRL B1865	1	0	45	55	62	20	−15.5°	18	+23°
<i>S. albus</i> LAB 3840 ⁵⁾	3	0	80	20	64	22	−15.7°	14	+22.8°
<i>S. antibioticus</i> NRRL 8167 ¹¹⁾	5	0	35	65	77	7		16	+22.7°
<i>S. aureofaciens</i> NRRL 8092 ¹²⁾	6	0	14	86	45	50	−16.1°	5	
<i>S. chartreusis</i> NRRL 3882 ⁸⁾	1	0	58	42	66	25	−15.9°	9	
<i>S. cinnamomensis</i> ATCC 15413 ⁹⁾	4	0	55	45	59	18	−15°	23	+22.2°
<i>S. griseorubiginosus</i> LAB 4915 ⁶⁾	2	0	31	69	64	34	−16.1°	2	
<i>S. griseus</i> LAB 2142 ⁴⁾	7	0	25	75	86	6		8	
<i>S. malachitofuscus</i> ATCC 31547 ¹⁰⁾	5	0	65	35	71	19	−15.7°	10	
<i>Nocardia asteroides</i> COB 906 ⁷⁾	9	0	9	81	20	70	−16.2°	7	
<i>N. asteroides</i> COB 911 ⁷⁾	9	5	90	5	85	8		7	
<i>N. rubra</i> ATCC 15906	9	0	100	0	62	26	−15.9°	7	
<i>Streptomyces anulatus</i> COB 2212 ⁷⁾	1	11	70	19	79	10		11	
<i>S. griseoincarnatus</i> COB 914 ⁷⁾	1	0	71	29	77	5		18	+22.6°
<i>S. griseolus</i> COB 912 ⁷⁾	1	0	64	36	52	18	−15.5°	30	+23°
	2	10	90	0	79	21	−15.4°	0	
<i>S. griseus</i> ATCC 10137 ¹³⁾	8	0	33	67	58	30	−15.9°	12	
<i>S. phaeofaciens</i> COB 153 ⁷⁾	1	8	37	55	68	9		23	+22.9°

are set out in Table I. The reactions were performed at least twice and the average values are reported.

Among the strains studied, eleven of streptomycetes (in the first part of Table I) produced polyether carboxylic ionophorous antibiotics. Reduction of α,β -unsaturated ketone systems is known to occur during their biosynthesis.¹⁶⁾ In all cases of 2-cyclohexen-1-one **1a**, the reaction had gone to completion after 48 hr. However, marked differences, in the amount of alcohol **3a** obtained were often observed between strains. At 50 mg per 100 ml of medium, the proportion of alcohol obtained ranged from 20% (*S. albus* LAB 3840) to over 85% (*S. aureofaciens* NRRL 8092). With *Beauveria sulfurescens* under the same conditions, the proportion of alcohol was about 50%.

Another antibiotic-producing strain was tested, namely *S. griseorubiginosus* LAB 3840, which produces cinerubins A and B. Their biosyntheses involve a polyketide intermediary and reduction steps. This strain gave results similar to the others.

Finally, some actinomycetes known for their oxidizing properties gave the same reaction. Thus *S. griseus* ATCC 10137, the oxidizing properties of which enable it to be used as a microbial model of drug metabolism,¹³⁾ was used.

In addition, strains able to degrade natural rubber were tried. Four of the 7 strains completely reduced 2-cyclohexen-1-one **1a** giving varying amounts of cyclohexanol **3a** (from 0 to 80%).

Results for the reduction of 2-methyl-2-cyclohexen-1-one **1b** are in Table I. The reduction of this ketone was slower than the reduction of the unmethylated ketone. More than 50% of the unconverted ethylenic ketone was often recovered. The best results were obtained with *N. asteroides* COB 906 (80% reduction after 48 hr). On the other hand, *S. albus* NRRL 1476 left the starting material intact. It can be noticed that generally when the reaction occurred, a large amount of saturated ketone **2b** was obtained.

With 2-methyl-2-cyclohexen-1-one, both reaction products had asymmetrical centers (one for the ketone and two for the alcohol). The stereochemistry of the products was obtained by comparing the observed optical rotations with those given in the literature, i.e.,

$$(S) (-) \text{ 2-methylcyclohexanone } [\alpha]_D^{20} = +16.8^{\circ(14)}$$

$$(1S,2R) (+) \text{ 2-methylcyclohexan-1-ol } [\alpha]_D = +24.3^{\circ(15)}$$

In Table I, one can observe that the stereochemistry

TABLE II. INFLUENCE OF SUBSTRATE CONCENTRATION ON THE REDUCTION RATE

Microorganisms
	1a	2a	3a
<i>Streptomyces albus</i> sub. sp. NRRL B1865			
40 mg	0	45	55
80 mg	72	23	7
<i>Nocardia asteroides</i> COB 906			
40 mg	5	90	5
80 mg	38	53	9
<i>Streptomyces griseolus</i> COB 912			
40 mg	10	90	0
80 mg	63	37	0

TABLE III. INFLUENCE OF AERATION ON THE REACTION RATE

Aeration rate (ml/liter/mn)
	1b	2b	3b
300	31%	14%	55%
100	37%	16%	48%
10	50%	13%	37%

of the reaction products was always the same (when this determination could be performed), *i.e.* (2*R*) for the saturated ketone **2b** and (1*S*,2*R*) for the saturated alcohol **3b**. These configurations are the same as those observed with *Beauveria sulfurescens*³⁾ and other fungi.¹⁾

In some cases the amount of substrate was varied. In Table II are set out the results of reactions with 40 mg or 80 mg of substrate for 100 ml of culture medium.

The increase of substrate concentration always lead to an important decrease of the reaction rate.

Finally, the influence of aeration rate on the reduction of 2-methyl-2-cyclohexen-1-one **1b** by the strain *S. albus* NRRL B1477 was studied in a fermentor. The results are given in Table III.

Thus, unlike *B. sulfurescens*,³⁾ these strains gave a higher yield at high aeration rates.

CONCLUSION

Evidently, the reduction of the carbon-carbon double bond of α,β -unsaturated cyclohexenones is common to those strains of actinomycetes possessing widely differing biochemical properties. No strains of actinomycetes were tested which failed to effect this reduction with unsubstituted cyclohexenone.

REFERENCES

- 1) A. Kergomard, M. F. Renard and H. Veschambre, *Agric. Biol. Chem.*, **46**, 97 (1982).
- 2) M. Desrut, A. Kergomard, M. F. Renard and H. Veschambre, *Biochem. Biophys. Res. Commun.*, **110**, 908 (1983).
- 3) A. Kergomard, M. F. Renard and H. Veschambre, *J. Org. Chem.*, **47**, 792 (1982).
- 4) P. Gachon, A. Kergomard, T. Staron and C. Esteve, *J. Antibiot.*, **28**, 345 (1975).
- 5) C. Delhomme, A. Kergomard, G. Kergomard and T. Staron, *J. Antibiot.*, **29**, 692 (1976).
- 6) L. David, M. Duteurtre, A. Kergomard, G. Kergomard, E. Scanzi and T. Staron, *J. Antibiot.*, **33**, 49 (1980).
- 7) M. Borel, Thèse Docteur-Ingénieur, Clermont, 1980.
- 8) M. O. Chaney, P. Y. Demarco, N. D. Jones and J. C. Occolowitz, *J. Am. Chem. Soc.*, **96**, 1932 (1974).
- 9) A. Agtarap, J. W. Chamberlin, M. Pinkerton and C. Steinrauf, *J. Am. Chem. Soc.*, **89**, 5737 (1967).
- 10) C. M. Liu and J. Westley, European Patent Appl. 9,636 (CI.CO7D493/20), 16 April 1980.
- 11) J. W. Westley, R. H. Evans, C. M. Liu, T. Hermann and J. F. Blount, *J. Am. Chem. Soc.*, **100**, 6784 (1978).
- 12) J. L. Occolowitz, D. H. Berg, M. Debono and R. L. Hamill, *Biomed. Mass. Spectros.*, **3**, 272 (1976).
- 13) J. P. Rosazza and R. V. Smith, "Advances in Applied Microbiology," Vol. 25, Academic Press, New York, 1979, p. 169.
- 14) J. Barry, A. Horeau and H. Kagan, *Bull. Soc. Chim. (France)*, 989 (1970).
- 15) R. Backstrom and B. Sjöberg, *Ark. Kemi*, **26**, 549 (1967).
- 16) J. W. Westley, "Polyethers Antibiotics Biosynthesis," in "Antibiotics," Vol. IV, Biosynthesis, ed. by J. W. Corcoran, Springer Verlag, Berlin, 1981.