Notes on Duality Theorems

Ryûki MATSUDA*

§ 0.

In § 1, we improve [2]. In § 2, we give a counter example for the Tannaka duality of topological groups. Results of these sections are after Roeder [3]. In § 3, we supplement [1]. The author thanks Mr. Hirabuki for his encouragement.

§ 1.

1. The case [2], 3. If only G_2 satisfies the Tannaka duality, θ is a homeomorphism. Conversely, if θ is a homeomorphism for $G_1=Z$ (the rational integers), G_2 satisfies the Tannaka duality.

2. The case [2], 5. If only G_2 satisfies the generalized Tannaka duality, F and Ψ_0 are canonically homeomorphic. Conversely, if F and Ψ_0 are canonically homeomorphic for $G_1=Z$, G_2 satisfies the generalized Tannaka duality.

3. The case [2], 7. If only g_2 satisfies the Tannaka duality, $\theta: F \rightarrow \Psi$ is a bijection. Conversely, if θ is a bijection for $g_1=K$, g_2 satisfies the Tannaka duality.

The same holds for the generalized Tannaka duality.

4. The case [2], 9. If only G_2 satisfies the Tannaka duality, $\theta: F \rightarrow \Psi$ is bijective.

θ is injective for any G_1 , if and only if G_2 is linear.

The same holds for the generalized Tannaka duality.

§ 2.

1. Let G be a topological group. For any positive integer n , we define nG the intersection of kernels of all continuous n -dimensional complex representations of G . Let V be a neighbourhood of 1 in G .

Received August 15, 1971.

*Department of Mathematics, Faculty of Science, Ibaraki University, Mito, Ibaraki, Japan.

If G satisfies the generalized Tannaka duality [4], then we have $nG \subset V$ for all sufficiently large n .

2. We see, using the fact of 1, that the example by Roeder ([3], 4) is a counter example for the generalized Tannaka duality (accordingly for the Tannaka duality [5]).

§ 3.

It is proved in [1] that the universal domain of characteristic 2 does not satisfy the Tannaka duality as Lie algebra. Let p be any positive prime number. We will prove moreover that *any field K of characteristic p does not satisfy the Tannaka duality nor the generalized Tannaka duality.*

PROOF. Let $A \in \mathfrak{gl}(n, K)$. We get a representation ρ_A of the Lie algebra K by setting $\rho_A(x) = xA$ for $x \in K$. All representations of K are of this form. We denote $K^\times$ the set of representations of K . We get a special representation ζ of $K^\times$ by setting $\zeta(\rho_A) = A^p$ for $\rho_A \in K^\times$. Now assume that the canonical map $\Phi: K \rightarrow K_0^{\times \times}$ to the set of special representations of $K^\times$ is bijective. We have $\Phi(x) = \zeta$ for some $x \in K$. Setting

$$M = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix},$$

we get the contradiction:

$$\begin{bmatrix} x & x \\ 0 & x \end{bmatrix} = xM = \rho_M(x) = \Phi(x)(\rho_M) = \zeta(\rho_M) = M^p = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

References

- [1] R. Matsuda, A counter example for the duality of Lie algebras, Bull. Fac. Sci., Ibaraki Univ., Math., **2-2**(1970), 1-3.
- [2] R. Matsuda and K. Sato, A topic on duality theorems, Bull. Fac. Sci., Ibaraki Univ., Math., **2-1**(1969), 15-18.
- [3] D. Roeder, A characterization of unitary duality, Trans. Amer. Math. Soc., **148** (1970), 129-135.
- [4] K. Sugita and M. Sugiura, On a certain type of duality theorem, Sci. Papers of the College of Gen. Edu., Univ. of Tokyo, **18**(1968), 115-124.
- [5] M. Sugiura, Some Remarks on duality theorems of Lie groups, Proc. Japan Acad., **43**(1967), 927-931.