1P069**Observation of GroEL-GroES action by high speed atomic force microscopy**○Daisuke Yamamoto^{1,2}, Masaaki Taniguchi¹, Toshio Ando^{1,2}¹Dept. of Physics, Kanazawa Univ., ²JST/CREST

Chaperonin GroEL guides a misfolded protein to its native state in the presence of co-chaperonin GroES and ATP. GroEL has a double ring structure, and GroES binds to the end faces of GroEL. It is widely believed that GroES cannot simultaneously bind to both ends of GroEL ring during the chaperonin reaction cycle. However, this alternate GroES binding has not directly been evidenced. To elucidate this negatively cooperative binding action directly using a high-speed atomic force microscope (high-speed AFM), we first prepared a sample system in which GroEL was lying on its side on a substratum. Because GroEL attaches onto mica in an end-up orientation, an appropriate substrate was needed. For this purpose, we performed two-dimensional crystallization of streptavidin on supported phospholipid bilayer. Over a wide area the obtained streptavidin crystal was flat enough for AFM observation. For the modification of GroEL with biotin, a GroEL mutant was expressed; Asp at the equatorial domain was replaced with Cys. With these preparations it became possible to observe GroEL from its side by high-speed AFM. GroEL alone looked like a barrel, while GroEL associated with one GroES looked like a bullet. In the presence of GroES and ATP, dynamic changes in the GroEL appearance were successfully captured on video. We will report the details of these studies.

1P071**The allosteric transition of GroEL in the presence of ADP**

○Kazunobu Takahashi, Hokuto Yamada, Kosuke Maki, Kunihiro Kuwajima

Department of Physics, School of Science, University of Tokyo

The chaperonin GroEL from *Escherichia coli* is a tetradecameric protein of 14 identical 57 kDa subunits arranged in two heptameric rings stacked back-to-back with a central cavity. The ATP-dependent control of the chaperone functions of GroEL is underpinned by two levels of allostery, one within each ring and the other between the rings. To simplify the complexity of this system, we have used a single ring variant of GroEL (SR1) in addition to wild-type GroEL. We investigated the ATP-induced allosteric transitions of single-tryptophan variants (Y485W-SR1 and Y485W-GroEL) of SR1 and wild-type GroEL, respectively, by stopped-flow fluorescence measurement. In each case, the ATP concentration dependence of the rate constant of allosteric transition exhibits a multiple sigmoidal behavior. To explain this behavior, we have assumed that GroEL has another second ATP binding site distinct from the well-known first ATP binding sites. This assumption leads to the idea that blocking ATP binding to the second nucleotide binding site brings about changes in the ATP concentration dependence of the rate constant of allosteric transition. We thus investigated the ATP-induced allosteric transition of Y485W-SR1 and Y485W-GroEL in the presence of excessive amount of ADP as an inhibitor of ATP. Based on these results, we will discuss the possibility of the existence of the second nucleotide binding sites and the effect of inter-ring interactions.

1P070**Single-molecule imaging of chaperonin GroEL-GroES dynamics using high-speed atomic force microscopy**○Masaaki Taniguchi¹, Daisuke Yamamoto^{1,2}, Noriyuki Kodera¹, Toshio Ando^{1,2}¹Department of Physics, Faculty of Science, Kanazawa University, ²JST/CREST

High-speed atomic force microscopy (high-speed AFM) makes it possible to observe directly the dynamics of protein-protein interactions at single molecule level. The double-ring chaperonin GroEL and co-chaperonin GroES are molecular machines that facilitate protein folding by undergoing ATP-dependent structural movements. To analyze the dynamic molecular processes contained in a chaperonin cycle, we attempted high-speed AFM imaging of GroEL. GroEL is adsorbed onto mica in an upright manner, which allows us to study height changes accompanied by its conformational change. We detected significant height differences (1.1 nm) between neighboring GroEL molecules in a low [ATP]. When GroES was added to the buffer solution, GroES were repeatedly observed to associate with and dissociate from GroEL. The height difference between GroEL alone and the GroEL-GroES complex was 3.4 nm. The lifetime distribution of GroEL-GroES complex revealed that approximately two reaction steps go through prior to their dissociation. Using UV-flash photolysis of caged ATP, we also detected a rapid height increase in GroEL alone, which was followed by a slow height decrease characterized with a single rate constant. In addition, we observed rapid GroES binding to GroEL immediately after UV irradiation, which was followed by slow dissociation. Thus, we succeeded in sketching dynamic processes of the GroES encapsulation reaction through direct visualization of conformational changes in individual GroEL molecules and their interaction with GroES.

1P072**Flexible C-terminal region of GroEL contributes to maintaining the environment of central cavity for proper protein folding**

○Koudai Machida, Akane Okada, Kunihiro Hongo, Tomohiro Mizobata, Yasushi Kawata

Dept. Biotech., Fac. Eng. and Grad. Sch. of Biomed. Sci., Tottori University

In order to understand the functional role of the C-terminal region of GroEL, we generated and characterized C-terminal deletion mutants (dC541, dC531 and dC525). First, we monitored the fluorescent resonance transfer from the rhodanese bound on apical domain to the C-terminal amino acid of GroEL labeled with FITC and TMR-5MA, respectively. FRET efficiencies for all truncated mutants were almost the same, indicating that the C-terminal region of the wild-type GroEL is not protruding into the central cavity, but forming a wall at the bottom of the cavity. ANS fluorescence intensity increased in proportion to the truncated length of amino acid residues, indicating that the property of the C-terminal region inside the cavity was changed to more hydrophobic by the truncation. ATPase activities of all truncated mutants were suppressed by approximately 50% in the presence of GroES. Therefore, interactions of these mutants with GroES were similar to wild-type GroEL. Furthermore, all truncated mutants showed similar ATPase activities to wild-type GroEL in the presence of rhodanese. However, the refolding abilities for several substrates decreased in proportion to the truncated length of amino acid residues. Furthermore, the refolding abilities for rhodanese, which is encapsulated by SR1 or truncated SR1 mutants, were the same as the double-ring GroEL. These results suggested clearly that the flexible C-terminal region of GroEL contributes to maintaining the environment of central cavity for proper protein folding.