1P077**Isothermal titration calorimetric and fluorescence quenching studies on the phloem exudate lectin from pumpkin (*Cucurbita maxima*)**

○A. Narahari, Musti J. Swamy

School of Chemistry, Univ. of Hyderabad, Hyderabad-500046

The chitoooligosaccharide-specific lectin from pumpkin (*Cucurbita maxima*) phloem exudate (PPL) has been purified to homogeneity by affinity chromatography on chitin. Binding of chitoooligosaccharides [(GlcNAc)_{3,5}] to PPL has been studied by isothermal titration calorimetry (ITC). Thermodynamic parameters obtained show that the dimeric PPL binds to two ligand molecules of chitotriose, chitotetraose and chitopentaose with comparable affinity at 298.15 K. The binding reactions were essentially enthalpy driven with the enthalpy of binding for the different chitoooligosaccharides ranging between -16.52 and -21.39 Kcal.mol⁻¹, whereas the entropic contribution to the binding reaction is negative, with the value of entropy of binding being -31.9 to -52.7 cal.mol⁻¹.K⁻¹. The accessibility and exposure of tryptophan residues in PPL were investigated by fluorescence quenching using a neutral quencher (acrylamide) and two ionic quenchers (I⁻ and Cs⁺). The emission λ_{max} of native PPL is seen at 338 nm, indicating that the Trp residues are significantly exposed to the aqueous environment. However, quenching with acrylamide was considerably higher than that with I⁻ and Cs⁺, clearly indicating that some of the Trp residues are not accessible to the ionic quenchers, which cannot penetrate into the protein interior. Further, among the two ionic quenchers the extent of quenching is more with iodide as compared to Cs⁺, suggesting the presence of positively charged residues near one or more Trp residues.

1P079**The interaction mechanism between the RING-H2 finger domain of ubiquitin ligase and the ubiquitin conjugating enzyme**

○Etsuko Katoh, Shizue katoh, Koji Kikuchi, Yuki Tsunoda, Cui Hong, Eiichi Minami

National Institute of Agrobiological Sciences

Ubiquitination requires three sequential enzymatic reactions: (i) an activating enzyme (E1) forms a thiol ester between one of its cysteines and the ubiquitin Gly76 carboxyl; (ii) a conjugating enzyme (E2) transiently carries the ubiquitin (as a thiol ester); and (iii) a ligase (E3) helps to transfer the activated ubiquitin from E2 to the substrate Lys. Generally, eukaryotic cells contain a single type of E1, multiple types of E2s, and many different E3s. In this presentation, we report experimental results such as 1) determination of solution structure of E3 RING-H2 finger domain and NMR based modeling structure of E2, 2) NMR titration experiments, 3) in vitro ubiquitination assay and 4) mutagenesis experiments of E3 RING-H2 finger domain and E2. From these experimental results, hydrophobic and basic residues (V136, W165, L166 and R176) of E3 RING-H2 finger domain are very important to bind E2. Especially, Trp(W) most important residue for E3 activity (= E2 binding) because many E3 RING finger protein has W residue at same position. On the other hand, the chemical shifts of hydrophobic and acidic residues (F62, A96, L97, T98, I99 and E90) of E2 show large chemical shift change. All of these residues are located in flexible loops and it is shown that these loops are recognition site of E3 RING finger domain. These results show that the complex formation between E2 and E3 RING finger domain occurred based on hydrophobic and electrostatic interaction.

1P078**Characterization of the interaction between Pex3p and Pex19p involved in Peroxisomal membrane protein trafficking**○Yasuhiko Sato¹, Hiroyuki Shibata², Hiroaki Nakano¹, Tsuneo Imanaka³, Hiroaki Kato¹¹Grad. Sch. Pharm, Kyoto Univ., ²Natl. Cardiovas. Ctr. Res. Inst., ³Dept. of Biol. Chem., Fac.of Pharm. Sci., Univ. of Toyama

Peroxisomes are discrete, single membrane-bound organelles and their biogenesis requires essential factors, peroxins (Pex). All peroxisomal proteins are synthesized in the cytoplasm and their translocation processes are proceeded by several Pex proteins. Pex3p and Pex19p contribute to the translocation of peroxisomal membrane proteins (PMPs). Pex3p is thought to localize on the peroxisomal membrane and recognizes a Pex19p carrying its cargo PMPs. Thus, the interaction between Pex3p and Pex19p is thought to play an important role in the PMPs trafficking mechanisms. To elucidate the molecular mechanisms of their interaction, we characterized their fluorescence properties. We performed intrinsic Trp fluorescence titrations of Pex3p with Trp-less Pex19 mutant, Pex19pW202F. Pex3p has two Trp residues, Trp104 and Trp224. The maximum wavelength of fluorescence spectrum for Pex3p was 340 nm. By addition of Pex19pW202F, the maximum wavelength was blue-shifted to 320 nm. Since the large 20-nm blue shift was observed, we expected that either or both of two Trp residues of Pex3p could contribute directly to the binding to Pex19p. Pull-down assays with using Pex3p in which these Trp were replaced with Phe or Ala, demonstrated that Pex3p (34-373) W104A did not bind to Pex19p, whereas W104F and W224A did. Thus we suggest that the side chain of Trp104 in Pex3p is directly participated in the interaction with Pex19p.

1P080**Bacterial Unsaturated Glycoside Hydrolases as Virulent Factors: Novel Catalytic Reaction Mechanism**○Takafumi Itoh¹, Akihito Ochiai², Bunzo Mikami¹, Wataru Hashimoto², Kousaku Murata²¹Div. of Appl. Life Sci., Grad. Sch. of Agriculture, Kyoto Univ., ²Div. of Food Sci. and Biotech., Grad. Sch. of Agriculture, Kyoto Univ.

Bacillus sp. GL1 unsaturated glucuronyl hydrolase (UGL), a member of glycoside hydrolase family GH-88, is a bacterial enzyme that degrades mammalian glycosaminoglycans and bacterial biofilms. The enzyme acts on unsaturated glucuronate saccharides with a α -glycoside bond produced by lyases responsible for bacterial invasion upon host cells and releases Δ GlcA. In this study, we detailed crystal structures of inactive UGL (D88N) complexed with three substrates, identified the active site residues, and postulated a novel reaction mechanism catalyzed by UGL that triggers the hydration of the vinyl ether group in Δ GlcA, but not of glycoside bond. In the proposed mechanism, Asp-149 acts as a general acid and base catalyst. On the other hand, YteR is derived from *Bacillus subtilis* strain 168 and has an overall structure similar to that of UGL, although it exhibits little amino acid sequence identity with UGL. YteR was found to constitute a novel enzyme, unsaturated galacturonyl hydrolase. This enzyme acts specifically on unsaturated rhamnogalacturonan (RG; Δ GalA-Rha) with a β -glycoside bond obtained from plant cell wall RG type-I treated with lyases and releases Δ GalA. We also clarify the active site residues and substrate recognition of YteR, based on a crystal structure of inactive mutant YteR (D143N) in complex with the substrate. These findings provide molecular insights not only into enzyme reaction mechanism but also into pathogenic bacterial systems for degradation of mammalian and plant cell surfaces.