1P261

Fluctuation analysis of microtubule sliding over kinesin thick filaments in vitro

○Daigo Hirata¹, Toshiki Taba¹, Masaki Edamatsu², Yoko Y. Toyoshima², Akira Yamada³, Yasuhiro Imafuku¹, Katsuhisa Tawada¹

¹Department of Biology, Kyushu University, ²Department of Life Sciences, Graduate School of Arts and Sciences, University of Tokyo, ³Biological ICT Group, National Institute of Information and Communications Technology

Kinesin is a motor protein that moves along a microtubule (MT) toward the plus end. A number of kinesin molecules are involved in intracellular transportations. In the present study, we investigated whether kinesin molecules cooperate one another or not to generate the MT movement. To test the cooperation, we measured the fluctuation of MT movement over a kinesin thick filament and its dependence on the microtubule length. For the fluctuation analysis, we used "kinesin" thick filaments, which are copolymers of catchin molecules and kinesin-myosin rod chimera molecules. These two molecules reassemble to form a straight and bipolar filament, in which kinesin motor domains are uniformly orientated (Taba et al., unpublished). Thus, we can avoid possible effects of the random orientation of the motor domain on the fluctuation analysis (Sekimoto & Tawada, 1995).

According to the probability theory, the effective diffusion coefficient, a measure of fluctuation, is proportional to the inverse of the microtubule length if the actions of kinesin motors are stochastically independent and random. However, we found that the effective diffusion coefficient does not depend on the microtubule length for a kinesin chimera containing intact neck region. This finding suggests the existence of cooperation between kinesin motors generating the MT movement. We are planning to study other kinesin chimeras containing truncated neck regions as well as the dependence of the diffusion coefficient on the sliding direction along kinesin thick filament.

1P263

Mutational studies of GTP-binding motif of FlhF in regulation of the number and location of flagella in *Vibrio alginolyticus*

○Akiko Kusumoto¹, Hiroyuki Terashima¹, Akari Shinohara¹, Toshiharu Yakushi^{1,2}, Seiji Kojima¹, Michio Homma¹

¹Dept. of Biological Science, Nagoya Univ., ²Fac. of Agriculture, Shinshu Univ.

Precise regulation of the number and the location of the bacterial flagella is critical for efficient motility. In a mono-polar-flagellated bacterium, *Vibrio alginolyticus*, flagellar number is regulated positively by FlhF and negatively by FlhG. FlhF is also involved in the location of flagella in the cell. FlhF has GTP-binding motif and shows similarity to *E. coli* signal recognition particle receptor, FtsY. Little is known about FlhF role in regulation of the number and the location of flagella. The aim of this study is to clarify the relationship between GTP-binding motif and FlhF function. A series of Ala replacements in the GTP-binding motif of FlhF was generated and each mutant protein was N-terminally fused to GFP to examine their localization in $\Delta flhF$ cells that have no flagellum. Expression of FlhF(WT) in $\Delta flhF$ cells caused production of flagella at the pole. FlhF(WT) was found throughout cytoplasm and some of them were localized at the flagellated pole. FlhF(T306A) failed to produce any flagellum in $\Delta flhF$ cells and was seen throughout cytoplasm. FlhF(G380A) produced more flagella at the pole in $\Delta flhF$ cells than that of FlhF(WT). FlhF(G380A) was intensely localized at the flagellated pole. These results suggest that GTPase activity is involved in the regulation of flagellar number and that polar localization of FlhF is required for production of flagella at the pole.

1P262

Improvement and analysis of ATP synthase crystals

○Yasuo Shirakihara¹, Aya Shiratori¹, Satoshi Murakami², Toshiharu Suzuki³, Masasuke Yoshida³

¹National Institute of Genetics, ²Osaka University, ³Tokyo Institute of technology

ATP synthase is responsible for ATP production in living cells, and is a membrane protein located in the energy conversion membrane. ATP synthase consists of a channel Fo portion and a large soluble catalytic F1 portion. The unique rotational catalysis mechanism of F1 includes rotation of the rod-like subunit.

In order to understand the working mechanism of ATP synthase, we previously started its crystallographic study using the thermophilic synthase. We had obtained 7Å diffraction patterns. We have been doing further studies to get better diffracting crystals, and also have been doing analyses of the diffraction patterns having a high mosaicity feature. This report describes those studies.

Firstly, we have continued intensive examination of the protein preparation by changing detergent species, PS3 culture batches, kinds of nucleotide present in the extraction step and various purification parameters. The results will be given in our poster with a special reference to the features distinct between decyl-maltoside and dodecyl-maltoside, both of which had been previously assumed to be similar to each other.

Secondly, we made further efforts to analyze the diffraction patterns obtained until last year. The analysis had been hampered by incorrect beam position parameters supplied and a high mosaicity of crystals. Data were recollected in a way that should allow diffraction analysis, and an analysis is going with our refined procedures.

1P264

Control of kinesin-microtubule motility by caged peptides derived from the kinesin C-terminus domain

○Akiko Nomura^{1,2}, Taro Q. P. Uyeda¹, Noboru Yumoto¹, Yoshiro Tatsu¹

¹AIST, ²RIKEN

The kinesin-microtubules system may be ideally suited for developing nanoscale biodevices because kinesin is able to transport intracellular materials (e.g., vesicles, chromosomes) along cytoskeletal networks composed of microtubule filaments. To design a nanoscale biodevice that can be controlled by an external stimulus, we have introduced photochemical switches into the kinesin-microtubule in vitro motility system. In this study, we examined the inhibitory activities of peptides derived from the kinesin C-terminus domain on the motility. The inhibitory effects of each peptide against the kinesin motility were evaluated by monitoring the motility of the fluorescent-labeled microtubules using fluorescence microscopy. Amino acid residues within the C-terminal region that are essential for the inhibitory effects were identified by alanine scanning mutagenesis. Based on the obtained results, one or two caging groups were introduced into the amino acids that are critical for the inhibition of the microtubule motility. The caged peptides only slightly affected the microtubule motility in the absence of UV-irradiation, but effectively inhibited the motility after the UV-irradiation. We demonstrated that the caged peptides from the kinesin C-terminus are promising tools for the development of photo-controlled nanoscale biodevices based on the kinesin-microtubule system.