1P541**Multiple domain interactions between the single cell adhesion molecule, nectin and cadherin, revealed by high sensitive force measurements**

○Yoshikazu Tsukasaki¹, Kazuo Kitamura², Kazuya Shimizu³, Atsuko H. Iwane¹, Yoshimi Takai⁴, Toshio Yanagida¹

¹Dept. of Nanobiology, Osaka Univ., ²Dept. of Neuroscience, Osaka Univ., ³Dept. of Internal Medicine, Kobe Univ., ⁴Dept. of Molecular Biology and Biochemistry, Osaka Univ.

Nectins and cadherins are cell adhesion molecules (CAMs) with key roles in the organization of various types of cell-cell adhesions. In this study, single molecule intermolecular force microscopy (IFM), with the force sensitivity at sub-piconewtons, was used to characterize the physical properties of the extracellular trans-interaction between paired nectins and between paired cadherins. Three different bond states between paired nectins and four different bond states between paired cadherins were identified based on bond strength where each bond state had unique bond lengths and lifetimes. Furthermore, as a result of analysis for the bond strength distribution, the bond lengths and lifetimes were not constant changing as a function of loading rate. These results lead to a model in which nectins and cadherins interact in multiple zipper-like manners across antiparallel aligned domains. Additionally, these results suggest that the multiple and cooperative zipper-like interactions allow nectins and cadherins to adaptively respond to dynamic exploratory recognition processes between cells.

1P543**Single-molecule structural and functional analyses of nuclear pore complex**

Shotaro Otsuka, Hirohide Takahashi, ○Shige H. Yoshimura
Graduate School of Biostudies, Kyoto University

Macromolecular traffic between the nucleus and the cytoplasm is governed by the interaction between cargo molecules and the nuclear pore complex (NPC). We recently developed a new system to investigate the molecular mechanism of the nuclear transport at single-molecule level by combining fluorescence microscopy and atomic force microscopy (AFM). The obtained results are the followings. (i) The nuclear envelope (NE) containing functional NPCs are reconstituted around a bead or on a glass substrate coated with small G-protein, Ran, by using *Xenopus* egg extract. Fluorescently-labeled nucleoporins (Nups), which were expressed in and purified from insect cells, were successfully incorporated into this reconstitution system to fluorescently label the NPC. (ii) The NE was isolated from *Xenopus* oocyte and spread onto a glass substrate by microsurgery. The AFM imaging of this isolated NE in a physiological buffer revealed that the cytoplasmic face and the nucleoplasmic face of the nuclear envelope and NPC have different structures. The structure of the NPC was largely affected by the presence of the cargo complex (importin α , importin β and NLS proteins) in the cytoplasm. (iii) By using a newly-developed glutathione-coupled cantilever, we demonstrated that the unbinding force between importin α and importin β was around 50 pN. The results of these single-molecule measurements will be reported in detail.

1P542**Single molecule imaging of chaperonin functions using zero-mode waveguides**

○Taro Ueno¹, Takashi Tanii², Naonobu Shimamoto², Takeo Miyake², Hironori Sonobe², Iwao Odumari², Takashi Funatsu¹

¹Graduate School of Pharmaceutical Sciences, The University of Tokyo, ²Graduate School of Science and Engineering, Waseda University

In this study, we aim to investigate chaperonin functions. Chaperonin GroEL, which is composed of two rings, mediates protein folding while co-chaperonin GroES binds to each ring of GroEL alternatively. We have analyzed GroEL-GroES interactions at a single molecule level using a total internal reflection fluorescence microscopy (TIRFM), and revealed the important intermediate state in the chaperonin reaction cycle at 10 nM GroES concentration. However, it is still unclear whether there is other intermediate state such as GroES-GroEL-GroES complex because GroES exists at about 5 μ M concentration in *E.coli*. In order to analyze chaperonin functions at the physiological condition, the illumination volume of 0.2 fL in TIRFM should be reduced to less than 1 aL. Arrays of zero-mode waveguides (ZMW), that is, nano-holes arrays lined up on a metal-clad quartz glass can reduce the illumination volume by three orders of magnitude than that by TIRFM. After IC5- and biotin- labeled GroEL was immobilized via biotinylated BSA and streptavidin on ZMW, the solution containing 200 nM Cy3-GroES, 2 mM ATP, denatured protein as substrate was introduced on the quartz slide. On the position of the IC5-GroEL preassigned, Cy3-GroES appeared and disappeared again and again. The histogram of durations of on-time, the time from binding to release, indicated that GroES released from GroEL after two sequential steps, which agreed with previous studies. Next, we are analyzing about GroES-GroEL-GroES complex.

1P544**Direct observation of unfolding-refolding dynamics of single cytochrome c proteins with milliseconds time resolution**

○Kenji Okamoto, Masayoshi Nishiyama, Masahide Terazima
Dept. of Chemistry, Grad. Sch. of Science, Kyoto Univ.

Protein folding process has been extensively investigated and is to date known to proceed by way of at least a few intermediate states even in the case of quite small proteins like cytochrome c (cyt c). Accurate understanding of those intermediate states is necessarily significant. We investigate the conformational dynamics of cyt c using a single molecule technique, which can unveil information lost in ensemble averaging. Cyt c molecules can take several kinds of structure depending on the solvent condition. We investigated the conformational change of cyt c under the various solvent conditions. Especially under the equilibrium condition, molecules fluctuate between a compact (native or molten-globule) and a unfolded structure and intermediate states in unfolding-refolding process are expected to be observed. Structure change of dye-labeled cyt c can be detected through the fluorescence intensity change, since cyt c has a heme, which can quench fluorescence emission of a dye in its vicinity with fluorescence resonance energy transfer (FRET), and the FRET efficiency depends on the heme-dye distance. We introduce a novel single molecule fluorescence detection system, which is based on information theory and can resolve state transitions from the fluorescence signal acquired by a photon counting detector with high time resolution like milliseconds. Our technique realizes direct observation of transient intermediate states in fast conformational dynamics of single protein molecules.