2P061

ATP-dependent interactions between *Escherichia coli* MinC/MinD complex and phospholipid membrane

○Takashi Okuno¹, Toshihiko Sakurai², Mayumi Maruyama¹, Shun Hirota^{1,3}, Noriaki Funasaki¹

¹Dept. of Phys. Chem., Kyoto Pharm. Univ., ²Dept. of Biotech., Faculty of Eng., Tottori Univ., ³PRESTO, JST

Escherichia coli divide in the center of the cell, producing two daughter cells with equal size. Division of cells at inappropriate site gives rise to chromosomeless minicells. The selection of the division site must be highly regulated. Division of cells in *E. coli* is initiated by formation of a FtsZ (tubulin-like protein) ring at the middle of the cell. The position of the FtsZ ring is regulated by min operon gene products (MinC, MinD, and MinE). MinC is a cell division inhibitor which prevents formation of a FtsZ ring. Walker-type MinD ATPase and MinE control the localization of MinC in the cells. Cytological studies showed that the MinC/MinD complex localizes into a horseshoe structure at the polar membrane and undergoes a pole-to-pole oscillation in living cells. ATP-dependent binding of MinD to the inner membrane surface must be a key step in the oscillatory dynamics of the Min system. To investigate the molecular mechanism of MinD ATPase, we prepared MinD mutants lacking or defecting ATP hydrolysis activity by introducing mutations to Lys residues in the ATP binding pocket. The MinD mutants lack membrane binding activity, but have affinity for ATP as same as that of wild type MinD. These results indicated that the Lys residues play important roles in the membrane binding steps. We also have purified the MinC protein with a His-tag without losing its activity. We will discuss on the molecular properties of the MinC/MinD interaction *in vitro*.

2P063

Expression and purification of bacteriophage T5 A1 and A2 proteins are related with viral DNA injection

○Tomoko Nakao^{1,2}, Laure Plancon¹, Lucienne Letellier¹

¹Institut de Biochimie et Biophysique Moléculaire et Cellulaire, Université de Paris-Sud, ²Department of Biomolecular Engineering, Graduate School of Bioscience and Biotechnology, Tokyo Institute of Technology

Bacteriophage T5 transfers its DNA in the *E. coli* cytoplasm through bacterial membranes like other phages. However T5 is known to have a particular DNA injection process in two steps. In the first step, called first step transfer (FST), 8% of DNA is injected in the bacteria. This DNA codes for pre-early proteins. Then there is a pause of a few minutes for the DNA being transcribed and translated. After the pause, the rest of the DNA or second step-transfer (SST) (92%) is injected into the bacteria. After completion of DNA injection, the early proteins appear, and at last the synthesis of the late proteins begins and continues until lyses. It is suggested that A1 and A2, synthesized during the pause between the two steps, was required to trigger the SST. Both proteins were purified from T5 phage infected cells. But details of their functions are not clear.

In order to investigate the role of pre-early proteins of bacteriophage T5, we cloned and expressed these two proteins, A1 and A2, in *E. coli*. Owing to the toxicity to hosts, it was impossible to purify A1 for biochemical analysis. On the other hand, we succeeded in purifying A2 from soluble cytoplasmic fraction by Ni-chelating and ion exchange chromatographies. Protein cross-linking analysis and size exclusion chromatography experiments indicated that purified A2 is composed of homodimer. And this dimer is stable even in high concentration of urea and heat treatment. The nucleotide sequence specific interactions between A2 and DNA have not been observed.

2P062

Expression and functional analysis of lipocalin super-family proteins in the olfactory epithelium of the newt

○Mandula Glaqin, Tsukasa Takahashi, Ken Sawada, Tatsuo Iwasa

Dept. of Mat. Sci. and Eng., Muroran IT

Olfactory sensation starts by binding of odorants to the olfactory receptor. The olfactory epithelium is covered with mucus secreted by Bowman's gland and odorants are usually hydrophobic molecules. Thus, the odorants are expected to be carried to olfactory cilium by an odorant-binding protein, a member of lipocalin super-family. We found two cDNA clones at high frequency from a cDNA library of the olfactory epithelium of *Cynopus pyrrhogaster*. The full-length cDNA clones were obtained and the deduced amino acid sequences of the clones showed a high homology to olfactory specific protein (OSP) and lipocalin, respectively. We named them as Cp-lip and Cp-OSP, respectively. The molecular phylogenetic tree showed that they belong to the lipocalin super-family, a group of small proteins (ca. 20 kDa) with a well conserved β -barrel structure. *In situ* hybridization analysis revealed that both proteins were expressed in Bowman's gland, but distribution of the cells which expressed the each protein was different in the nasal organ. The results suggested that odorants may be discriminated by lipocalins in a mucus layer before they bind to the odorant receptor. To test the possibility, we expressed both proteins in *E. coli*. After solubilization with 8M Urea solution, Cp-lip was purified by His-tag affinity column. We measure conformational stability and a ligand-binding character of the expressed proteins and will discuss these properties of both proteins based on their primary structural difference.

2P064

Dual Functional Property of non-heme iron containing ferritin Pfr in *Helicobacter pylori* under acid stress

○I-Ju Yeh¹, Shyh-Horng Chiou²

¹Dept. of Biochemistry, National Taiwan University, ²Dept. of Biochemistry, National Taiwan University

H. pylori is a spiral, slow-growing Gram-negative microaerophilic bacterium. It is unique and distinctive among various bacterial pathogens for its ability to persist in the extreme acidic environment of the human stomach. To identify *H. pylori* genes in response to low pH, we have used a proteomic approach to study the protein expression of *H. pylori* under neutral (pH 7) and acidic (pH 5) environments. Global protein-expression profiles were analyzed by high-resolution two-dimensional gel electrophoresis (2-DE) followed by LC-MS/MS and mass comparison. The results revealed that more than ten proteins were differentially expressed under different pH conditions, with decreased amounts in seven proteins and increased in six proteins. Time-course RT-PCR study reveals that mRNA expression level of a non-heme iron containing ferritin Pfr is enhanced 1.2 folds and another electron-acceptor protein flavodoxin is decreased 2 folds at pH 5 as compared with that at pH 7. Moreover, non-heme iron containing ferritin Pfr was found to lose its ferrioxdase activity at pH 5, and it was found to bind DNA binding more strongly at pH 5 than at pH 7 as revealed by gel-retardation assay. Fluorescence spectra revealed the tertiary structure difference between pH 5 and pH 7. Therefore in addition to the predominant role of urease in the protection of *H. pylori* under low pH, non-heme iron containing ferritin Pfr may have a dual function for *H. pylori* to adapt to the acidic environment of gastric lumen.