2P068 イヌミルクリゾチームの物性解析に向けた安定化変異体の構築
○野中 康宏¹、秋枝 大介¹、神谷 昌克^{1,2}、相沢 智康^{1,2}、出村 誠^{1,2}、河野 敬
—^{1,2}
2aF02 ¹北大院・理・生命理学、²北大院・生命科学
Construction of mutant canine milk lysozyme stabilized against degradation for
physical analysis
Yasuhiro Nonaka (1), Daisuke Akieda (1), Masakatsu Kamiya (1,2), Tomoyasu
Aizawa (1,2), Makoto Demura (1,2) and Keiichi Kawano (1,2). (1: Div. Biological
Sciences, Grad. Sch. of Science, Hokkaido Univ.; 2: Div. Life Science, Grad. Sch. of
Life Science, Hokkaido Univ.)

Canine milk lysozyme (CML) is a model protein that forms a partially unfolded, molten globule state at equilibrium. However, charge heterogeneity is easily generated in the recombinant sample after expression and purification. We confirmed that this heterogeneity was caused by spontaneous asparaginyl deamidation. Asparaginyl deamidation is a common non-enzymatic modification of proteins and peptides, transforming an asparaginyl residue into an aspartate or isoaspartate residue. CML can easily undergo asparaginyl deamidation at multiple sites under mild conditions. Substitution of the labile asparaginyl residues to glutaminy residues prevented deamidation effectively. We constructed the quintuple mutant CML that has no deamidation site. The overall conformation of this mutant was equivalent to that of wild type in the native state. The most interesting property of CML, molten globule formation was also observed as well as wild type. Stabilization of the CML against deamidation improves the yield of pure recombinant protein and the reproducibility of experiments. In addition, recombinant CML expressed in *E. coli* has an extra N-terminal methionine residue (Met 0). The presence of Met 0 in CML decreases the thermal stability, solubility, and perhaps metal binding affinity. We obtained recombinant CML without Met 0 from methylotrophic yeast expression system. We analyzed the effect of the extra methionine residue on the structure and stability of CML.

2P070 サルモネラ菌べん毛蛋白質輸送装置サブユニット FlhAC 温度感受性変異体の物理化学的測定
○島田 賢史¹、西條 由見子²、古川 進朗²、南野 徹^{1,2}、今田 勝巳^{1,2}、難波 啓
—^{1,2}
¹阪大院 生命機能、²Dynamic NanoMachine Project, ICORP, JST
Physicochemical analysis of TS-mutants of FlhAC, a subunit of the flagellar protein
export apparatus of Salmonella
Masafumi Shimada (1), (2), Yumiko Saijo-Hamano (2), Yukio Furukawa (2),
Tohru Minamino (1), (2), Katsumi Imada (1), (2), Keiichi Namba (1), (2), (1: Grad Sc
Frontier Biosci, Osaka Univ; 2: Dynamic NanoMachine Project, ICORP, JST)

Mutants that show different behaviors from wild type upon temperature raise are known as "Temperature sensitive mutant (TS-mutant)." Because TS-mutants can do on/off control of functions by temperature, they have been used for function analysis. Although it has been assumed that this phenotype is caused by structural instability derived from mutation, its detail analysis such as a thermal transition analysis of purified mutant proteins, has never been done. We carried out such analysis on FlhAC, the C-terminal domain of a protein component of the flagellar protein export apparatus. Salmonella flagellum axial complex is divided into 3 parts: the basal body, the hook, and the filament. All the flagellar axial proteins are translocated into the central channel of the flagellum for self-assembly. Protein translocation is mediated by the flagellar protein export apparatus, which is believed to be located on the cytoplasmic side of the basal body. This export apparatus is composed of six transmembrane proteins and three cytoplasmic proteins. The transmembrane protein FlhA (75kDa) is the largest component and has two domains: N-terminal hydrophobic part that contains eight transmembrane segments (FlhATM), and C-terminal hydrophilic part that forms a large cytoplasmic domain (FlhAC). We carried out a thermal transition analysis by Far-UV-CD measurement of wt-FlhAC, a TS-mutant of FlhAC (G368C) and its suppressor mutants and obtained clear evidence of different transition behaviors corresponding to their phenotypes upon temperature raise.

2P069 ジスルフィド結合を導入することによって高度に安定化した低温ショック相同蛋白質の熱力学的評価
○長谷川 葵¹、飯塚 一成¹、林(岩崎) 容子²、城所 俊一¹
2aF03 ¹長岡技科大・生物系、²共和化工株式会社・環境微生物学研究所
Thermodynamic evaluation of the highly stabilized cold shock protein homologue by
introducing a disulfide bond
Aoi Hasegawa (1), Kazumari Iizuka (1), Yoko Hayashi-Iwasaki (2), Shun-ichi
Kidokoro (1) (1: Department of Bioengineering, Nagaoka University of
Technology; 2: Institute of Environmental Microbiology, Kyowa Kako Co., Ltd)

The aim of this study is to elucidate the stabilization mechanism of the highly stabilized cold shock protein mutant based on the homologue from extreme thermophile, thermophilus HB8, by introducing a disulfide bond considering the three-dimensional estimated structure, and by evaluating the thermodynamic stability precisely with differential scanning calorimetry (DSC). The pair of the mutation sites to be displaced by cysteine were determined using the following three criteria: (1) the mutation sites were largely separated along the amino-acid sequence, (2) the side-chain of the mutation sites were estimated to be near enough to make a disulfide bond in the native three-dimensional structure, and (3) the side chain of the sites were exposed to the solvent, and finally A11C/T40C mutant was selected as the most hopeful candidate. The mutant gene by two-step PCR with the wild type template containing His-tag was introduced to *E. coli* host, BL21 to produce the mutant. The expressed proteins were once denatured, purified by Ni-NTA column, refolded by dialysis with multi-step dilution of the denaturant, and finally processed by TEV protease to remove the His-tag. From the CD measurement, the mutant was found to have the same three-dimensional structure as that of the wild type. The transition temperature evaluated by DSC at pH 3.5 was higher than that of the wild type by more than 20 K.

2P071 トランス-シクロヘキサレンジアミンによるリボヌクレアーゼ A とリゾチームの加熱失活抑制: 静電力和疎水性力の役割
○平野 篤¹、浜田 寛之¹、白木 賢太郎¹
¹筑波大院・数理工学
Trans-cyclohexanediamine prevents thermal inactivation of ribonuclease A and
lysozyme: Roles of electrostatic and hydrophobic forces.
Atsushi Hirano (1), Hiroyuki Hamada (1) and Kentaro Shiraki (1). (1: Inst. Appl.
Phys., Univ. Tsukuba)

Protein tends to form inactive aggregates in solution. The protein aggregates cause many problems for biotechnology and pharmaceutical industries. We show that trans-cyclohexanediamine effectively prevents thermal inactivation of ribonuclease A (RNaseA) and hen egg lysozyme. At neutral pH (< 8), heat-induced inactivation of RNaseA and lysozyme was improved by the addition of trans-1,2-cyclohexanediamine (12CHDA) better than trans-1,4-cyclohexanediamine (14CHDA) in solution, while at alkaline pH (> 8), the inactivation of RNaseA and lysozyme was improved more effectively by 14CHDA than 12CHDA. The results indicate that the aggregation of RNaseA and lysozyme was affected by the electrostatic repulsions in the presence of trans-cyclohexanediamine molecules. Moreover, hydrophobic amino acids could be dissolved better in 12CHDA solution than in 14CHDA solution. The results indicate that 12CHDA strongly interacts with hydrophobic residues, leading to prevent aggregation of RNaseA and lysozyme at neutral pHs. Increase in the repulsive force between aggregation-prone protein molecules may depend on both factors, hydrophobic interaction between protein molecules and electrostatic repulsion of the additives on protein.