- 3P124 分子動力学シミュレーションと FT-IR を用いたアモルファストレハロースの水素結合特性の解析
 ○河崎 直子¹、朱 波²、古木 隆生¹、櫻井 実¹
¹東工大院・バイオセンター、²東工大院・生命理工
 Analysis of Hydrogen Bonds in the Glassy States of Trehalose and Neotrehalose Using Molecular Dynamics Simulation and FT-IR Spectroscopy
 Naoko Kawasaki (1), Bo Zhu (2), Takao Furuki (1) and Minoru Sakurai (1). (1: Center for Biological Resources and Informatics, Tokyo Institute of Technology; 2: Dept. of Biomolecular Engineering, Graduate School of Bioscience and Biotechnology, Tokyo Institute of Technology)

Trehalose is known to be a good protectant against various environmental stresses in many organisms. For instance, larvae of *Polypedilum vanderplanki* accumulate a large amount of trehalose when they are desiccated and their metabolic activity becomes undetectable, and this state is maintained as a result of vitrification of the accumulated trehalose. From DSC measurements, it was found that trehalose has the highest glass transition temperature (T_g) among the naturally occurring gluco-disaccharides. In this study, to understand why trehalose has such excellent properties as a desiccation protectant, we studied its glassy state at atomic level using molecular dynamics simulations and compared the results with those for neotrehalose. The simulations revealed that the number of intermolecular hydrogen bonds (n_{HB}) of trehalose is smaller than that of neotrehalose in the glassy state and the slopes of n_{HB} against temperature obviously change at T_g . To confirm these tendencies, temperature scanning FT-IR measurements were done for dehydrated amorphous sugars, and the temperature-dependent shifts of O-H stretching vibration bands were examined. The results (n_{HB} , T_g , etc.) obtained from these measurements were consistent with the tendencies found from the above MD simulations. From these results, it was confirmed that the higher T_g of trehalose, compared with neotrehalose, is not ascribed to the contribution of intermolecular hydrogen bonds, but to other properties deduced from the α,α -1,1 glycosidic linkage of this sugar.

- 3P126 ナノポアにおけるイオン透過機構の理論的解析
 ○炭竈 享司¹、斉藤 真司²、大峰 巖¹
¹名大院理、²分子研
 Theoretical Analysis of Ion Permeation Mechanism in Nanopore
 Takashi Sumikama(1), Shinji Saito(2) and Iwao Ohmine(1).(1: Dept of Chemistry, Faculty of Science, Nagoya University; 2: Dept of Theoretical and Computational Molecular Science, Institute for Molecular Science)

Transport of ions through ion channel has been extensively investigated since the determination of the X ray structure of the K⁺ channel. However, the dynamics of ions had not been understood well, and the energetics of ion permeation is not studied so far. We mimicked the ion channel by the anion-doped carbon nanotube, this approximation made it possible to observe the full dynamics of the ions in the model channel. The analyses of the dynamics indicate the main rate-limiting step is how an ion enters the channel, i.e., how an ion replaces the position originally located by a water molecule, which may imply that the reaction of the ion permeation is entropy bottleneck. In order to clarify this point, we evaluated the energy, free energy, and entropy profile along the ion permeation reaction coordinate, and compared this mutual importance. The result shows that the component of the free energy barrier located at the entrance of the channel consists of the entropy, and free energy minima are also comprised of the entropy. On the other hand, the energy term just makes the barrier of about 2 kcal/mol, alters slightly the shape of the free energy profile. That is, the entire process of ion permeation in this model channel is dominated by the entropy.

- 3P125 水溶液中におけるイノシトール異性体と β -アミロイド間相互作用に関する分子動力学シミュレーション
 張本 乾一¹、○朱雀 史朗¹、櫻井 実¹
¹東工大バイオセンター
 Molecular dynamics study of the interaction between inositol isomers and β -amyloid in aqueous solution
 Kenichi Harimoto, Shiro Sujaku and Minoru Sakurai. (Center for Biological Resources and Informatics, Tokyo Institute of technology)

Recently much evidence has been accumulated that some of inositol isomers have the ability of suppressing the aggregation of β beta;-amyloid (A β) in aqueous solution and hence they exhibit some therapeutic effect on Alzheimer disease. However, the functional mechanism of inositol remains unclear. Considering that sugar and polyol such as inositol affect hydrophobic interactions of biomolecules, it is hypothesized that inositol affects the aggregation properties of A β by modifying water structure. To confirm this, we performed molecular dynamics (MD) simulations for aqueous solutions of pure inositol and inositol/A β mixtures. Here, we focused our attention on two isomers of inositol, scyllo- and myo-inositol. The present simulations revealed the following points: 1) scyllo-inositol molecules are excluded selectively from the hydration shells around the hydrophobic parts of A β , while myo-inositol molecules are distributed almost uniformly around the whole hydration shell of A β , 2) consequently a larger number of water molecules exist in the hydration shell of A β in the scyllo-inositol system than in the myo-inositol one, 3) the water molecules surrounding a scyllo-inositol molecule are regularly ordered to fit the water structure, while those around a myo-inositol molecule is less ordered. In conclusion, by addition of scyllo-inositol, the degree of preferential hydration around A β molecules is enhanced, leading to the suppression of their amyloid formation or modification of their aggregation form.

- 3P127 3次元RISM理論による溶媒効果を取り入れた分子動力学法の開発
 ○宮田 竜彦¹、平田 文男¹
¹分子研・理論
 Combination of Molecular Dynamics Method and 3d-RISM Theory for Conformational Sampling of Flexible Molecules in Solution
 Tatsuhiko Miyata and Fumio Hirata. (Department of Theoretical and Computational Molecular Science, Institute for Molecular Science)

We have developed an algorithm for sampling the conformational space of large flexible molecules in solution, which combines the molecular dynamics (MD) method and the 3-dimensional reference interaction site model (3d-RISM) theory. The latter is a statistical mechanical theory describing the 3-dimensional solvation structure around solute molecules as well as the solvation free energy. The solvent-induced force acting on solute atoms was evaluated as the gradient of the solvation free energy with respect to the solute-atom coordinates. In order to enhance the speed of computation, we have applied a multiple timestep algorithm based on the RESPA (Reversible System Propagator Algorithm) to the combined MD/3d-RISM method. By virtue of the multiple timestep algorithm, one can choose a longer timestep for renewing the gradient of solvation free energy compared with that of the conformational update. We applied the present method to acetylacetone in aqueous solution, as a simple model of flexible solute. Acetylacetone possesses an intramolecular hydrogen-bonding capability, and it is known that hydration effect generally reduces the barrier height in conformational space around intramolecular hydrogen bonds. The present method sampled larger conformational space of acetylacetone than the MD simulation in vacuum. This result qualitatively agrees with the above general observation on the reduction of barrier heights around hydrogen-bonding in solution.