3P308 src 遺伝子産物のミリストイル基の部位特異的修飾法を用いた機能解析
○近藤 晶子¹、萩原 英雄¹、谷川 瑞穂¹、中村 政志¹、大野 敏²、横川 隆志²、
光岡 有美子²、細谷 孝充³、平松 俊行³、鈴木 正昭⁴、橋本 敬一郎¹、西
川 一八²、林 宣宏¹
¹藤田保健衛生大・総医研、²岐阜大・工・生命工、³東工大・院・生命理工、⁴岐阜大・
院医・再生医科学
**Functional analyses of N-terminal myristoylation of Src using site-directed-labeling
method**
Akiko Kondow (1), Hideo Hagihara (1), Mizuho Yagawa (1), Masashi Nakamura
(1), Satoshi Ohno (2), Takashi Yokogawa (2), Yumiko Mitsuoka (2), Takamitsu
Hosoya(3), Toshiyuki Hiramatsu (3), Masaaki Suzuki (4), Keiichiro Hashimoto
(1), Kazuya Nishikawa (2) and Nubuhiro Hayashi (1) (1: Inst. for Comprehen.
Med. Sci., Fujita Hlth. Univ; 2: Fac. of Eng., Gifu Univ; 3: Grad. Sc. of Biosci. and
Biotech, Tokyo Inst. of Tech; 4: Grad. Sch. of Med, Gifu Univ)

The lipid modification, N-myristoylation, of v-Src has been required for its ability to induce tumour although the kinase activity is unaffected in its N-myristoylation-deficient mutant. A function of the N-myristoylation is to anchor a protein to lipid membrane via weak hydrophobic interaction. Therefore, it has been hypothesized that signaling mediated by Src is regulated by its subcellular localization. One candidate protein regulating the subcellular localization of Src is calmodulin, which interacts with N-myristoyl moiety of Src. In the present study, we aimed to observe real-time membrane localization of Src *in vivo*, and to investigate role of calmodulin in subcellular localization of Src by introducing fluorescent-labeled Src protein in living cells. We chose a site-specific-labeling method for introduction of single fluorescence dye per protein to avoid disturbance of the weak hydrophobic interaction by labeling target proteins with large fluorescent protein or excess hydrophobic fluorescence dye. We then investigate some methods to introduce the labeled protein into cells.

3P310 GFP の相互作用による蛍光変化を用いたタンパク質構造変化の検出
○岩井 草介¹、上田 太郎¹
¹産総研・セルエンジニアリング
Detection of conformational changes of proteins using GFP proximity imaging method
Sosuke Iwai and Taro Q.P. Uyeda (Research Institute of Cell Engineering,
National Institute of Advanced Industrial Science and Technology)

Conformational changes of proteins such as myosin motor have been examined using GFP-based fluorescence resonance energy transfer method. GFP proximity imaging (PRIM) is another imaging technique that utilizes fluorescence changes of GFP molecules that are brought to proximity (Ward et al., 1982; De Angelis et al., 1998). Here we have examined the possibility that PRIM can be used to detect conformational changes of proteins. To this end, two GFP molecules were linked in tandem by polypeptide linkers and the fluorescence spectra of the proteins were measured. We found that a combination of a wild-type and a circularly permuted GFP molecules with 11-amino acid liker gave a larger fluorescence change. The fluorescence excitation ratio of this PRIM pair at 390 and 488 nm was 0.18. To examine whether the PRIM pair of GFP molecules can be used to detect conformational changes of myosin motor, the light chain binding domain of Dictyostelium myosin II was replaced by the PRIM pair of GFP. We are currently studying the fluorescence changes of the GFP molecules induced by conformational changes of the myosin motor domain during the ATP-hydrolysis cycle.

3P309 細胞サイズベシクル内における転写反応の実時間解析
○辻 明彦¹
¹科学技術振興機構・ICORP・時空間秩序プロジェクト
Analysis of Transcription Reactions in a Cell-Sized Vesicle
Akihiko Tsuji, Spatio-Temporal Order Project, ICORP, JST

Regulation of gene expression is one of most essential functions for living organisms. The ON/OFF regulation of expression of an individual gene in a cell has been thought to be performed through binding of DNA binding proteins to the corresponding specific sequences on DNA and dissociation from those, and to be performed very strictly. However, in a cell, these are the reactions between one molecule (a gene on DNA) and several to several hundreds of molecules (DNA binding proteins), which is thought to be under thermodynamic fluctuation. Therefore, we should revisit the characteristics and robustness of the ON/OFF regulation of transcriptional reactions of an individual gene, by directly measuring transcriptional synthesis of a particular RNA from a single genome DNA. To address this subject, we take an approach to reconstitute cell-sized vesicles which contain plasmid/genome DNA and transcriptional and translational components, and analyze the transcription reactions and dynamics of the mRNA in real time. We prepared cell-sized vesicles encapsulating DNA and transcription components, and then the synthesis of a particular mRNA was monitored under fluorescence microscope. The synthesized RNA molecules were labeled with two fluorescent oligonucleotide probes that hybridize to an adjacent location on the target RNA, resulting in FRET. Results about kinetics of transcriptional reactions and localization of mRNA will be discussed.

3P311 全反射蛍光顕微鏡を用いたリアルタイム多点型蛍光相関分光法の開発
○大杉 友¹、金城 政孝¹
¹北大院・生命科学学院・細胞機能科学
**Multipoint fluorescence correlation spectroscopy for analysis of molecular dynamics in
live cells by total internal reflection microscopy**
Yu Ohsugi(1) and Masataka Kinjo(1). (1:Lab of Molecular Cell Dynamics,
Graduate School of Life Science, Hokkaido Univ)

Intracellular signaling pathways are triggered by the binding of a ligand to its receptor on the cell surface to regulate cell activities. Hence, elucidation of the molecular interactions on the plasma membrane is especially necessary for understanding of cellular functions and regulations. Recently, we have reported the application of TIR-FCS (Total Internal Reflection Fluorescence Correlation Spectroscopy) to cell biology. By using TIR-FCS, we measured membrane-binding fluorescent proteins in living cells. However, because a conventional FCS or TIR-FCS monitors at a single point, it lacks quantitative data in two-dimensional area and heterogeneous structure, such as plasma membranes. Thus, FCS measurements in a two-dimensional plane will be fascinating methods. Although our future goal is to develop FCS imaging in two dimensions, in this research we demonstrated a novel real-time multipoint FCS system combined with TIR-FM and multichannel correlation system.TIR-FCS employs an evanescent field as an excitation light, which is very thin and wide field. Thus, we set up a multipoint FCS system by using TIR-FCS system and putting seven optical fibers at image plane and detectors at the back of optical fibers.With this multipoint TIR-FCS, we measured fluctuation of fluorescence from fluorophores in solution or in live cells at seven points. Furthermore, we will report the analysis results of Kv4.2 potassium channnels in living cells by using multipoint TIR-FCS.