an actin filament.

1P-140 GFPを用いた蛍光ストレーンセンサーの開発とミオシン-アクチン相互作用の可視化

Visualizing myosin-actin interaction with a genetically encoded fluorescent strain sensor.

Sosuke Iwai and Taro Q. P. Uyeda. (Research Institute for Cell Engineering, National Institute of Advanced Industrial Science and Technology (AIST))

Many proteins have been shown to undergo conformational changes in response to externally applied force *in vitro*, but whether the force-induced protein conformational changes occur *in vivo* remains unclear. To reveal the force-induced conformational changes, or strains, within proteins in living cells, we have developed a genetically encoded fluorescent "strain sensor", by combining the proximity imaging (PRIM) technique (De Angelis *et al.*, 1998) and myosin-actin as a model system. PRIM depends on direct contact between two GFP molecules, which can lead to spectral changes. The developed PRIM-based strain sensor module (PriSSM) consists of the tandem fusion of a normal and circularly permuted GFP. To apply strain to PriSSM, it was inserted between the two motor domains of *Dictyostelium* myosin II. In the absence of strain, the two GFP moieties in PriSSM are in contact, whereas when the motor domains are bound to F-actin, PriSSM has a strained conformation, leading to the loss of contact and a concomitant spectral change. Using the sensor system, we found that the position of the lever arm in the rigor state was affected by mutations within the motor domain. Moreover, the sensor was used to visualize the interaction between myosin II and F-actin in *Dictyostelium* cells. We are currently examining the nucleotide-dependent lever arm positions in the myosin motor domain attached to F-actin, by modifying the sensor system.

1P-141 糸状仮足伸長時におけるミオシンXの『釣りかぎ』の役割

Interaction between myosin-X and integrin- β acts as a crampon during filopodia protrusion.

Tomonobu M. Watanabe (1)(2)(3), Hiroshi Tokuo(3), Kosuke Gonda(4), Hideo Higuchi(4) and Mitsuho Ikebe (3) (1:Immunology Frontier Research Center ,Osaka University. 2: Precursory Research for Embryonic Science and Technology, JST, 3: University of Massachusetts Medical School, 4: Biomedical Engineering Research Organization, Tohoku University)

The filopodia are finger-like cytoplasmic projections, which extend from the leading edge of migrating cells. Unconventional myosin-X has been shown to be involved in the protrusion of filopodia. It was previously shown that myosin X interacts with an adhesion factor, integrin- β , via the FERM domain in its tail. Here we clarify the mechanism by which the interaction of myosin-X with integrin- β induces the formation of filopodia. We found that the filopodia induced by myosin X underwent the cycle of elongation and retraction. Filopodia elongated 2.4 μ m during one protrusion cycle from the extension to retraction, and the following elongation started during the slow retraction period with certain probability. Unlike the full length myosin X, myosin-X without FERM could localize at te tip but it could neither stay at the tips of filopodia nor induce long filopodia. The deletion of FERM domain decreased the probability of the phased-extension. Furthermore, the adhesion of the filopodia on the substrate activated the extension, but not the retraction. These results suggest that myosin X moved to the tip of filopodia stays at the tips via binding to integrin- β , and this is critical to promote the initiation of the protrusion that is achieved by utilizing the adhesion of the membrane to the substrate. We propose the model that myosin-X plays a role of "crampon" in the protrusion of filopodia.

1P-142 リン酸結合蛋白を用いた F_1 -ATPase からのリン酸放出における β Glu190 の役割の解明

The role of β Glu190 in phosphate release from F_1 -ATPase revealed by phosphate binding protein

Shingo Harada, Takayuki Nishizaka and Tomoko Maseike. (Dept. Phys., Fac. Sci., Gakushuin Univ.)

F_1 -ATPase, the catalytic part of F_0F_1 -ATP synthase, is a rotary molecular motor in which rotation of the central γ subunit within the $\alpha_3\beta_3$ cylinder is powered by ATP hydrolysis in the β subunits.

To study the mechanism of P_i release from F_1 -ATPase, we measured P_i release from $\alpha_3\beta_3\gamma$ subcomplex of F_1 -ATPase from thermophilic *Bacillus* PS3 which has a mutated catalytic residue β Glu190 under uni-site conditions. Measurements were carried out by the phosphate binding protein method using stopped flow apparatus in which increase in fluorescence intensity of coumarin-labeled phosphate binding protein (MDCC-PBP) upon binding of P_i from F_1 -ATPase was monitored. Meanwhile, we also measured the rate of ATP cleavage to determine the rate constants of each chemical step separately under uni-site conditions.

The rate constant of P_i release from $\alpha_3\beta_3(E190D)\gamma$ was 0.10 ± 0.01 s^{-1} , which was 30 times smaller than that from $\alpha_3\beta_3(R333W)\gamma$ (2.8 ± 0.1 s^{-1}) (Maseike *et al.* 2002). It indicates that the catalytic residue β Glu190 has an essential role in P_i release from F_1 -ATPase. Comparing crystal structures of various nucleotide binding states, a role of β Glu190 in P_i release from F_1 -ATPase would be to control P_i release from F_1 -ATPase by changing electrostatic interactions between β Glu190 and P_i .

To study how P_i release drives mechanical rotation under multi-site catalysis conditions, we are developing the system to observe P_i release from single F_1 -ATPase molecules each enclosed in femtoliter chambers in array under the optical microscope.

1P-143 ダイニン・微小管複合体の構造学的研究

Structural analysis of outer arm dynein molecules bound to microtubules

Hironori Ueno(1), Takuo Yasunaga(2), Chikako Shingyoji(3) and Keiko Hirose(1)(1:Research Institute for Cell Engineering, AIST; 2:Department of Biochemical Engineering and Science, Kyushu Institute of Technology; 3:Department of Biological Sciences, Graduate School of Science, Univ of Tokyo)

Dynein is a molecular motor that hydrolyzes ATP and moves along a microtubule (MT). The predominant model for dynein motility implies that the stalks of dynein, while bound to MTs, rotate toward the MT plus end (Burgess *et al.*, 2003). However, stalks bound to MTs have been difficult to observe. To study the structures of dynein bound to MTs, we used flagellar outer arm dynein-MT complex in which dynein molecules regularly crossbridged MTs. We observed sliding of MTs from the dynein-MT complex, indicating that the dynein molecules in our complex are able to move MTs. In order to see the stalks clearly, the complex was observed by cryo-positive stain electron microscopy, which we have recently developed, and the resulting images were classified using hierarchical cluster analysis. The class-averaged images showed that the majority of dynein images in the no nucleotide state (with apyrase) exhibited one ring-like structure per dimeric molecule, indicating superimposition of the two heads. When ATP and vanadate were added (the ADP/Vi state), many of the images showed two rings in an 8 shape, with one of the rings staying close to the MT to which the tail attaches, whereas the other ring shifted closer to the stalk-side MT. Some of the second head in the ATP/Vi state were also shifted toward the MT minus end. The stalk angles relative to the MTs did not change significantly between the two nucleotide states. Our results disagree with models in which the stalk acts as a lever arm to amplify structural changes.

1P-144 腸内連鎖球菌V型ATPaseの外周ストークのサブユニット構成とその溶液構造

Stoichiometry and solution structure of the peripheral stalk subunits of *Enterococcus hirae* V-ATPase

Misaki Yamamoto (1,2), Satoru Unzai (3), Shinya Saijo (4), Kazuki Ito (5), Kenji Mizutani (2), Yoshimi Kakinuma (6), Shigeyuki Yokoyama (7), Ichiro Yamato (1), So Iwata (2,7,8), and Takeshi Murata (2,7,8). (1: Dept Biol Sci Tech, Tokyo Univ of Science; 2: Grad Sch of Med, Kyoto Univ; 3: Protein Design Lab, Yokohama City Univ; 4: JASRI, SPring-8; 5: Harima Inst, RIKEN; 6: Faculty Agri, Ehime Univ; 7: SSBC, Yokohama Inst, RIKEN; 8: JST, ERATO, Iwata Project)

The vacuolar ATPases (V-ATPases) function as ATP-dependent proton pumps in the membranes of acidic organelles, in plasma membranes of eukaryotic cells. This acidification participates in concentration of neurotransmitters, processing of secretory proteins, and endocytosis. The V-ATPase contains a globular catalytic domain, V_1 , which hydrolyzes ATP, attached by central and peripheral stalks to an integral membrane domain, V_0 , which pumps ions across the membrane. The subunit arrangement and stoichiometry of the peripheral stalk are still unclear.

V-ATPases are also found in bacteria. We have identified a variant of V-ATPase in a fermentative bacterium *Enterococcus hirae*. *E.hirae* Na^+ -ATPase is a homolog of eukaryotic V-ATPase, with a simpler subunit composition. The peripheral stalk of the complex is likely to be composed of NtpE and NtpF with the N-terminal hydrophilic domain of NtpI (NtpI^{Nterm}).

In this report, we purified NtpE, NtpF, and NtpI^{Nterm} individually, and examined property of purified NtpE-F-I^{Nterm} complex by analytical ultracentrifugation (AUC) and small-angle x-ray scattering (SAXS). The ternary complex appeared to be monodisperse, and the molecular masses estimated from AUC and SAXS indicated that it is formed with a 1:1:1 stoichiometry. A low resolution structure model of the NtpE-F-I^{Nterm} complex produced from the SAXS data showed an elongated "L" shape.

1P-145 粗視化モデルによる双頭キネシンのステップ機構の解明

Simulation study on stepping mechanism of Conventional Kinesin by coarse-grained model

Ryo Kanada (1), Kei-ichi Okazaki (1) and Shoji Takada (1). (1: Dept of Biophysics, Graduate School of Science, Kyoto University)

Conventional kinesin is a motor protein that moves processively and unidirectionally along a microtubule using the chemical energy produced by ATP hydrolysis. By a variety of experimental studies using optical trapping bead assays, kinesin, a dimeric motor protein with two head domains is supposed to make stepping motions in hand-over-hand manner (Howard *et al.*, 20001). In this manner, there are mainly three stages: (i) both of head are bound to MT, (ii) rear head detaches from MT, while front head still remains to attach to MT, (iii) the detaching head finds new binding site on MT and bind to MT. This fashion suggests that by repeating the above three stages kinesin moves its heads alternatively such as human walking. However, still many remain to be answered.

Detaching process: In stage (ii), why is the rear head rather than the front head likely to detach from MT? What is the mechanism to regulate the detaching process?

Attaching process: How can the detaching head find the proper site on MT? What is the mechanism to avoid miss binding?

In this study, to reveal the above question we try to reproduce the whole of stepping process by molecular dynamics simulation based on Go-like model. In particular, to discuss how the mechanical stepping couples to the chemical reaction, we will apply multiple-basin energy landscape (Okazaki *et al.*, 2006) to conventional kinesin. About the detail result of research we are going to report in poster presentation.