2P-304 生体膜による非平衡開放系の効果を取り入れた細胞の生命機能モデリング

Modeling of a living cell taking account of the effect of nonequilibrium open system by biomembrane

Masahiro Takinoue (1), Yue Ma (1, 2), and Kenichi Yoshikawa (1, 2). (1: Dept. Phys., Kyoto Univ.; 2: ICORP, JST)

Rhythmic phenomena are essential for life, and ubiquitously appear in living systems, for example, a cell cycle and a circadian rhythm. Therefore, construction of mathematical models for biological rhythmic phenomena is essential to understand dynamical aspects of life. Usually, in order to describe rhythmic phenomena in living systems, negative feedback regulation models of gene expression network are used. The descriptions by gene regulation network models reflect that a behavior of a living cell is fatefully controlled by genomic information. However, this type of models lacks the viewpoint of life as a nonequilibrium open system; that is, although a living cell sustains biochemical reactions using influx and efflux of energy and matter through biomembrane, the effects of those flux are not taken into account in the current models. In this study, we propose a novel mathematical model for rhythmic phenomena in a living cell and demonstrate that the flux of energy and matter through biomembrane plays an indispensable role in generating and sustaining rhythmic phenomena in a living cell.

2P-305 培養心筋細胞のペーシングにおける刺激波形の効果

Effect of waveform in electrical pacing of cultured cardiac cells

Akira Ozeki and Takahiro Harada. (Department of Human and Artificial Intelligent Systems, University of Fukui)

Nonlinear oscillations are ubiquitous in physiological systems, including our heartbeat. Entrainment by external stimuli is one of the most important properties of nonlinear oscillators. Rhythmic contractions of cardiac muscle cells have been extensively studied from this aspect. In the former studies, pulse-wise stimuli have been widely utilized, although it remains unclear whether another waveform of stimulus is more effective for entrainment. In this presentation, we report on the effect of waveform on entrainment of rhythmic contractions of cultured cardiac cells. In the experiment, we cultured rat ventricular muscle cells on a glass electrode, and field stimuli possessing several waveforms were applied. As a consequence, we found that pulse-wise stimulus with a narrow width (0.5% of the whole cycle) has the potential to entrain the rhythmic contraction with much less power compared to other waveforms. In addition, according to the theory of phase oscillators, it is generally proved that the most effective waveform of stimulus is the same as that of the phase response curve (PRC) of the oscillator. Hence, our experimental finding suggests that the PRC of cultured cardiac cells should possess a pulse-wise waveform with a narrow width. Now, we attempt to examine this prediction by separately measuring the PRC of cultured cardiac cells. We expect that this study leads to a better understanding of the properties of cardiac cells from nonlinear dynamics aspect.

2P-306 藍色細菌時計タンパク質の ESR 解析

The ESR analysis of the cyanobacterial circadian clock protein

Risa Mutoh (1)(2), Hiroyuki Mino (2), Reiko Murakami (1) and Masahiro Ishiura (1). (1: Center for Gene Research, Nagoya Univ.; 2: Graduate School of Science, Nagoya Univ.)

Circadian rhythms, biological oscillations with a period of 24-hour in various metabolic and behavioral activities, are observed ubiquitously from prokaryotes to eucaryotes including *Drosophila*, *Neurospora*, *Arabidopsis*, mice, and human. Cyanobacterial circadian clock proteins, KaiA, KaiB, and KaiC, play essential roles in generating circadian oscillations. KaiA, KaiB, and KaiC interact each other, and the stoichiometry of Kai protein complexes may change during circadian cycle.

In this study, we performed ESR analysis to clarify the structural changes of Kai proteins during circadian cycle. We constructed a plasmid expressing a Cys-substitution mutant of *Thermosynechococcus elongatus* KaiB, expressed the mutant in *Escherichia coli*, and purified it. We confirmed that the KaiB mutation did not affect *in vitro* KaiC phosphorylation rhythm in the presence of KaiA, KaiB, KaiC and ATP. Then, we labeled the mutant KaiB with spin labels and measured the ESR spectra of labeled KaiB. The signal intensity from the labeled KaiB increased in the presence of three Kai proteins and ATP, suggesting that structural changes of the spin-labeled KaiB might occur during circadian cycles. Similar increases were also observed in the presence of KaiA, but not observed in the presence of KaiC.

2P-307 *Synechocystis* sp. strain PCC6803 は生物時計たんぱく質 KaiA, KaiB および KaiC に加え、KaiB および KaiC ホモログを 2 つずつ有している。

Synechocystis sp. strain PCC6803 has the *kaiAB1C1* circadian clock gene and clock gene homologues, *kaiC2B2* and *kaiB3* and *kaiC3* genes.

Reiko Murakami (1), Kentaro Ishii (1)(2), Kiyoshi Onai (1), Megumi Morishita (1), Masahiro Ishiura (1)(1: Center for Gene Research, Nagoya University; 2: Department Biology, Graduate School of Science, Osaka University)

Circadian rhythms, 24-h biological oscillations of metabolic and behavioral activities observed ubiquitously in prokaryotes and eukaryotes, are endogenously regulated by circadian clocks. Cyanobacteria are the simplest organisms that exhibit circadian rhythms. We previously cloned and analyzed the *kaiABC* circadian clock gene cluster in the cyanobacterium *Synechococcus* sp. strain PCC

7942, which is essential for the generation of circadian rhythm. The cyanobacterium *Synechocystis* sp. strain PCC6803 (*Synechocystis*) has *kaiAB1C1* circadian clock gene cluster and additional clock gene homologues, *kaiC2B2* cluster, and *kaiB3* and *kaiC3* genes. In this study, we carried out biochemical analysis to characterize functional and structural difference among *Synechocystis* 3 KaiB proteins.

2P-308 シアノバクテリア時計関連蛋白質 SasA の生化学的構造解析

Biochemical and structural analysis of the cyanobacterial clock-related protein SasA

Jonathan Valencia Swain (1), Kyouhei Bitou (1), Reiko Murakami (1), Katsumi Imada (2), Masahiro Ishiura (1). (1: Center for Gene Research, Nagoya University; 2: Graduate School of Frontier Biosciences, Osaka University)

Circadian clocks generate internal biological rhythms with a period of about 24 hours. Various organisms are known to live by these processes, which can regulate from gene expression to even behavior. Cyanobacteria are the simplest organisms that have been found to have a circadian clock.

Circadian systems are composed of the sensory input system (for environmental input), the oscillator (entrained by the input), and the output system. In the cyanobacterium *Synechococcus* sp. strain PCC 7942 (*Synechococcus*) the oscillator is composed of the three clock proteins KaiA, KaiB and KaiC. Its oscillation is transduced to genome-wide transcription cycles mainly by the output pathway staged by the protein SasA (a histidine kinase) and its cognate response regulator, RpaA. In this pathway KaiC regulates the autophosphorylation of SasA, and SasA then transfers its phosphate group to RpaA.

The current structural, biophysical and biochemical analysis for this system is still lacking. Hence, in this study we searched for crystallization conditions for *Synechococcus* and *Thermosynechococcus elongatus* SasA, aimed towards a structural analysis by X-ray crystallography. We also analyzed the biochemical interaction between SasA and its pathway partners.

2P-309 EMSA による Pex 結合に重要な特定配列の同定

Identification of the specific nucleotide sequence critical for Pex binding by EMSA

Hiroshi Satoh (1)(3), Risa Mutoh (2)(3), Reiko Murakami (3) and Masahiro Ishiura (3). (1: School of Science, Nagoya Univ.; 2: Graduate School of Science, Nagoya Univ.; 3: Center for Gene Research, Nagoya Univ.)

Various organisms from prokaryotes to eucaryotes exhibit rhythms in various biochemical, physiological and behavioral activities with a period of about 24 hours, called circadian rhythms. The rhythms derive from an internal oscillator named a circadian clock. Cyanobacteria are the simplest organisms that have the circadian clock. The cyanobacterial circadian system is composed of three clock proteins, KaiA, KaiB and KaiC, and some clock-related proteins. Pex is a clock-related protein, which is involved in the clock light input pathway. Pex binds to an upstream region of the *kaiA* clock gene and suppresses its expression. *pex*-deficient Cells showed a shortened period phenotype.

Previously, we identified a minimal 25-bp sequence essential to Pex binding in the *kaiA* upstream region. In this study, to determine which base pairs are critical for Pex binding, we performed electrophoretic mobility shift assay (EMSA) using the minimal 25-bp sequence with nucleotide substitution mutations at various positions.

2P-310 F_1 -ATPase 1 分子の揺動散逸定理の破れの測定

Measurement of Violation of Fluctuation Dissipation Theorem of Single Molecule F_1 -ATPase

Shoichi Toyabe(1), Takahiro Watanabe-Nakayama(2), Tetsuaki Okamoto(1), Hiroshi Takekuni(1), Seishi Kudo(3), Shigeru Sugiyama(4) and Eiuro Muneaki(5). (1: Dept. of Physics, Graduate School of Science & Technology, Chuo Univ.; 2: Graduate School of Bioscience & Technology, Tokyo Institute of Technology; 3: Faculty of Engineering, Toin Univ. of Yokohama; 4: National Food Research Institute)

F_1 -ATPase is a rotary molecular motor, which rotates in one direction with hydrolyzing ATPs. F_1 motor works in nonequilibrium steady states (NESS). Although the fluctuation dissipation theorem (FDT) holds in equilibrium states, it generally violates in nonequilibrium states. Thus, the extent of FDT violation is recognized to characterize nonequilibrium states. Recently, the quantification of the FDT violation in NESS has been attracting many researches. Among those, the Harada-Sasa equality is especially a promising one. This equality connects the extent of the FDT violation to the energy dissipation rate and enables us to calculate the energy dissipation rate from readily obtainable quantities in experiments, such as the correlation function and response function of the velocity. This helps us to understand the energy flow through molecular motors.

In this presentation, we measured the FDT violation of a single F_1 motor. The frequent response function was obtained by applying a periodic torque utilizing an electrorotation method. We found that FDT violated in low frequencies but held in high frequencies. Thus, we specified the frequency range in which F_1 motor is in nonequilibrium states. Also, using Harada-Sasa equality, we found that 60-70% of the input energy of ATP hydrolysis was dissipated in the rotational freedom. This value was independent of ATP and ADP concentration. This is the first experiment to measure the FDT violation of molecular motors in a single molecule level.

2P-311 チャコウラナメクジ嗅覚系の電氣的振動パターンの解析

Analysis of electrical oscillation patterns of the olfactory center of land slugs