energetics in hydration phenomena. We tried to reproduce dynamical and physical properties of hydrating water molecules around several types of solutes by using classical MD simulations with widely used typical water models. After that, the hydrating water molecules around methane-like molecule with a moderate negative charge showed increased mobility compared with bulk water although the dynamical properties of hydrating water were difficult to quantitatively reproduce even for simple solutes such as I^- and K^+ ions unfortunately. On the other hand, a recent simulation based on a quantum chemical Car-Parrinello MD showed a weakened electric dipole and increased mobility of hydrating water molecules around alkaline metal ions (Na^+ , K^+) (J. Chem. Phys. 126: 034501, '07) although the results was quantitatively insufficient. From these, it is necessary to develop improved MD method to reproduce real hydrating water properties. Recent approaches will be presented. Finally, peculiar phenomena such as increased mobility of hydrating water around motor protein (Biophys J 85: 3154, '03) may be explained by calculating the amount of specific hydrating water in a reaction such as ATP hydrolysis.

3P-120 拡張 scaled particle 理論によって計算した動径分布関数

Radial distribution function calculated by using extended scaled particle theory

Daiki Mitarai (1) and Masayuki Irida (1)(2). (1: Graduate school of Computer Science and Systems Engineering, Kyushu Institute of Technology; 2: Department of Bioscience and Bioinformatics, Faculty of Computer Science and Systems Engineering, Kyushu Institute of Technology)

Macromolecular crowding effects have been studied intensively as an attractive force between proteins caused by a cell environment, which has a high density for large bio-molecules. This attractive force is driven by an entropy change of the surrounding bio-molecules. One of the simplest model systems for macromolecular crowding effect is hard-sphere system. In this study, radial distribution functions of hard-spheres have been calculated by using extended scaled particle theory (XSPT). Radial distribution function is derived from the work creating the cavity through the cavity function, $y(r)$. Works for creating a dumbbell-shaped cavity composed of two hard spheres in the hard-sphere solvent can be given by XSPT. We also calculated radial distribution functions by using fundamental measure theory (FMT) in the framework of the cavity function for a dumbbell-shaped cavity. Both XSPT and FMT can give works for a non-spherical cavity. However, the expressions from the two theories are different in spite of the identical expression in the case of a spherical cavity in the solvent. Our calculated results show that values of radial distribution functions from Monte Carlo simulation are sandwiched by those from XSPT and FMT. This result showed that two theories give analytical expressions for upper and lower limits for an effective potential for two spherical molecules under macromolecular crowding effect.

3P-121 周波数変調原子間力顕微鏡によるモデル生体膜上に形成された水和層の分子分解能観察

Molecular-scale imaging of hydration layers formed on a model biological membrane by frequency modulation atomic force microscopy

Takeshi Fukuma (1, 2). (1: Frontier Science Organization, Kanazawa University; 2: PRESTO, JST)

The interactions between water and biological molecules have the potential to influence the structure, dynamics, and function of biological systems. However, the molecular-scale investigation of hydration phenomena has been impeded by the lack of a method able to visualize molecular-scale structuring of water under physiological environments. Recently, we have made a great progress in the resolution and sensitivity of the liquid-environment atomic force microscopy using frequency modulation detection technique (FM-AFM). The method has enabled atomic-resolution imaging with pico-Newton order loading forces under physiological environments (i.e., in liquid). In this study, we have investigated the structuring of water at the interface of supported dipalmitoylphosphatidylcholine (DPPC) bilayers in phosphate buffer solution using FM-AFM. The force versus distance curves measured between the bilayer and the AFM tip show oscillatory force profiles with a peak spacing of 0.28 nm, indicative of the existence of up to two hydration layers next to the membrane surface. Furthermore, FM-AFM imaging at the water/lipid interface visualizes individual hydration layers in three dimensions, with molecular-scale corrugations corresponding to the lipid headgroups. The results demonstrate that the intrinsic hydration layers are stable enough to present multiple energy barriers to approaching nanoscale objects, such as proteins and solvated ions, and are expected to affect membrane permeability and transport.

3P-122 蛋白質の大局的構造変化に伴う水和体積の変化の解析

Analysis of hydration volume changes with global conformation changes of proteins

Yasutaka Seki (1), Jumpei Fujii (1) and Kunitugu Soda (1). (1: Dept. Bioeng., Nagaoka Univ. of Technol.)

The partial molecular volume V_p of a protein is a thermodynamic quantity that provides information on its 3D structure and hydration. Accurate estimation of the contribution of hydration water to V_p of proteins with a variety of conformations is essential for analyzing protein structures in various conformational states based on the experimental values of V_p . The aim of this paper is to evaluate changes in hydration contribution to V_p due to global conformation changes of proteins. We examined the spatial distribution of water around various solutes by MD simulation to find that the effect of hydration water to V_p consists of three contributions below: First, the low-density region around the molecular surface of

solute contributes to V_p positively. Second, the high-density region near the center of the first hydration layer contributes negatively. Third, a small positive contribution comes from the low-density region between the first and second hydration layers. The first contribution depends strongly on the polarity and shape of the molecular surface. The peak height and bottom depth responsible for the respective 2nd and 3rd contributions do not depend on surface characteristics, though both positions of the peak and bottom vary with the type of the surface atom. In addition, these contributions of hydration water were found similar for both proteins and amino acids. Based on these results, changes of V_p with the folding and unfolding of proteins will be discussed at the meeting.

3P-123 逆ミセル中タンパク質の赤外分光 I : テラヘルツ領域

Infrared spectroscopic studies on a protein in a reverse micelle I: THz spectral region

Hiroshi Murakami(1,4), Yuji Toyota (1,2), Takaki Nishi (1,3). (1: Kansai Photon Science Institute, Japan Atomic Energy Agency; 2: Dept Applied Physics, Graduated School of Engineering, Osaka City Univ; 3: Division of Chemistry and Material Technology, Graduated School of Science and Technology, Kyoto Institute of Technology; 4: Res. Unit Quantum Beam Life Sci., Japan Atomic Energy Agency)

It is believed that low-frequency vibrational motions of proteins play an important role on their functions. THz spectroscopy as well as inelastic neutron scattering measurements is a suitable method to investigate such vibrational motions. Most of the studies using THz spectroscopy and inelastic neutron scattering measurements have been done for lyophilized proteins with a certain hydration level. Background signal due to solvent waters is huge compared with signal due to protein molecules in protein aqueous solutions for such experimental methods. This is the reason why lyophilized proteins are used as a sample. It is important to investigate low-frequency vibrations of a protein in water because proteins work in an aqueous solution. A protein in a reversed micelle was employed as a sample for THz time domain spectroscopy using a femtosecond laser. The protein aqueous solution is in the nanometer-sized reversed micelle, while the solvent is non-polar one whose THz absorption is much weaker than that due to solvent waters. Therefore, content of water in the sample is very little. A surfactant, sodium bis(2-ethylhexyl) sulfosuccinate (AOT) was dissolved in a non-polar solvent of isooctane, and aqueous solution of myoglobin was injected into the solution. It has been found that the absorption spectrum due to the protein is obtained from its dependence on protein concentration. We will discuss the results in detail in the presentation.

3P-124 逆ミセル中タンパク質の赤外分光 II : 近赤外領域

Infrared spectroscopic studies on a protein in a reverse micelle II: NIR spectral region

Takaki nishi(1,2), Yuji Toyota(1,3), Hiroshi Murakami(1,4), Nobuyuki Ichinose(2). (1: Kansai Photon Science Institute, Japan Atomic Energy Agency; 2: Division of Chemistry and Material Technology, Graduated School of Science and Technology, Kyoto Institute of Technology; 3: Dept Applied Physics, Graduated School of Engineering, Osaka City Univ; 4: Res. Unit Quantum Beam Life Sci., Japan Atomic Energy Agency)

IR spectroscopy of amide bands is a suitable method to study hydrogen bonds in proteins. However, there are few studies on amide A band around 3300 cm^{-1} for aqueous solution of proteins, because the O-H stretching vibration of solvent waters shows huge absorption in the spectral range. The purpose of this study is to measure amide A band of a protein in a reverse micelle and to investigate how the IR spectrum depends on the micelle size. In the reverse micelle, protein aqueous solution is in the nanometer-sized reverse micelle, and so content of water in the sample is very little. The size of a reverse micelle is variable by changing a water-surfactant molar ratio, and affects hydrogen bonds of water in a reverse micelle. It is interesting to know how the spectrum of the amide A band, which sensitive to change in hydrogen bonds of proteins, depends on the micelle size. Aqueous solution of myoglobin was injected into a sodium bis(2-ethylhexyl) sulfosuccinate (AOT)/isooctane solution. The concentrations of water and AOT were constant, while that of myoglobin was varied. It has been found that the difference spectrum between the absorption spectra of the sample with different protein concentrations shows a peak around 3300 cm^{-1} without change in the spectral shape, and that the amplitude of it is proportional to the protein concentration under a constant water-surfactant molar ratio. We will discuss the results including the dependence of the spectrum on the water-surfactant molar ratio in the presentation.

3P-125 生体分子のモード選択性の解明

Elucidation of mode-selectivity of biomolecules

Hiroshi Teramoto (1), Tamiki Komatsuzaki (1) (1: Institute for electronic science, Hokkaido University)

Chemical reaction has following two aspects; selectivity and stochasticity. The conventional chemical reaction theory assumes stochasticity. It means that the outcome of the chemical reaction does not depend on the detailed initial condition of the reactant and the resultant distribution of the products only depends on the width of the bottleneck which mediates between the reactant and product. Many chemists believe that molecular dynamics is highly chaotic and the memory of the detailed initial condition immediately wears away as the system evolves. However, in 2002, B. C. Dian et al. (B. C. Dian et al. Science 296, 2369-(2002)) showed that there exists mode selectivity, which means that, depending on which vibrational mode they excite, the resultant product distribution can be significantly different in case of the structural transition reaction of NATMA, which consists of 36 atoms. This system has much more complicated potential