Hyonchol Kim (1), Hiroyuki Takei (2), Kenji Yasuda (3) ((1) Kanagawa Academy of Science and Technology; (2) Kanagawa Academy of Science and Technology; Department of Life Sciences, Faculty of Life Sciences, Toyo University; (3) Kanagawa Academy of Science and Technology; Department of Biomedical Information, Division of Biosystems, Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University)

Monitoring functions and variations of expression profiles in a cellular system is a key to understanding the mechanisms of living organisms, especially to identifying those phenotypical characteristics. The *in situ* labeling for expressed biomolecules is one of useful methods to visualize amounts and spatial distributions of such molecules, and detailed measurement in single cell, electron microscopy (EM) is well used with gold nano-particle labels. At the simultaneous labeling of several molecules, different sizes of particles are used; that yielded derogation of quantitative capability because of the difference of labeling efficiency depended on the sizes of particles. Moreover, size variation of practicable particles is not so many; therefore, new labeling method should be developed for exhaustive study of expressed biomolecules in a cell.

We tried to establish a new labeling technology in the EM, which develops conventional labeling of the EM to "multi-color" one. We prepared nano-particles with different metals in the same size and used it as "multi-color" label. To evaluate labeling ability of these particles, probe DNAs were immobilized on the particles and target DNAs on a substrate were labeled. Kinds of the labeled particles were identified by measuring backscattered electron using a field emission scanning electron microscope. Our method would achieve simultaneous measurement of exhaustive molecular expression in a single cell.

2TA3-09 細胞内の局所的な温度変化の測定とその解析

2P-258 Measurement and analysis of local temperature change inside living cells.

Kotaro Oyama (1), Madoka Suzuki (2), Vadim Tseeb (3), Fumichika Ono (4), Yusuke Seto (4), Masahiro Motoyoshi (1), Kaoru Iwai (5), Shin'ichi Ishiwata (3) ((1) Pure and Applied Physics, Graduate School of Advanced Science and Engineering, Waseda University; (2) COBRI, Comprehensive Research Organization, Waseda University; (3) Department of Physics, Faculty of Science and Engineering, Waseda University; (4) Bioscience and Biomedical Engineering, Graduate School of Advanced Science and Engineering, Waseda University; (5) Department of Chemistry, Faculty of Science, Nara Women's University)

Activity of proteins such as Ca^{2+} -ATPase is thermosensitive and thermogenic. Feedback loop between their activity and local temperature change may play an important role in biological systems. But there have been no experimental evidence that a cell uses intracellular temperature change effectively for intracellular processes. Recently, a few techniques have been developed to measure temperature change inside single living cells. We and other groups, using fluorescent thermosensitive polymers, have shown the intracellular temperature elevation of about 1°C that was induced by chemical stimulations such as ionomycin (Suzuki, M. et al., 2007, *Biophys. J.*, **92**: L46-L48) and FCCP (Gota, C. et al., 2009, *JACS*, **131**: 2766-2767). However, single cells cannot produce enough amount of heat to explain these phenomena from the viewpoint of calorimetric studies with cell suspension.

In this study, we tried to explain this gap by measuring and changing local temperature inside HeLa cells. Aims of this study are (i) the detection of localized heat sources in a cell which could rise temperature heterogeneously during thermogenesis induced by ionomycin, and (ii) the determination of cellular thermophysical coefficients. Our results showed that intracellular temperature elevation induced by ionomycin was relatively uniform. We found that the thermal diffusivity of a living cell was less than that of water. The value, however, was insufficient to bridge the gap. Therefore it is suggested that there are other missing viewpoints to connect the data from single cells and those from cell suspensions.

2TA4-01 機能的細菌べん毛モーターにおけるモーター構成素子の交換割合の見積り

Turnover of subunit-exchange in functioning bacterial flagellar motor

Hajime Fukuoka (1), Shun Terasawa (2), Hiroto Takahashi (1), Yuichi Inoue (1), Akihiko Ishijima (1) ((1) Tohoku Univ.; IMRAM; (2) Tohoku Univ.; Grad. Sch. Life Sci.)

Bacterial flagellum is a supramolecular complex and consists of a basal body (rotary motor), a helical filament, and a hook. The flagellar motor is driven by the electrochemical potential of ions, and the interaction between stators and a rotor in flagellar motor is thought to generate the motor rotation. Recently, it was shown that stator parts are exchanged in a functioning motor and that the assembly of stator into the motor requires coupling ion for the motor rotation. In our study, we focused on the dynamics and the exchange of rotor components in a functioning motor. We constructed the expression systems of GFP-FlhN, FlhM-GFP, and GFP-FlhG as GFP-fusions of rotor component. In tethered cells producing each GFP-fusions, fluorescent spot was localized at the rotational center. Each GFP fusion was probably incorporated into flagellar motor as a rotor component. To observe the exchange of rotor components, we carried out Fluorescent Recovery After Photobleaching experiment using evanescent illumination to the tethered cell. In a tethered cell producing GFP-FlhN or FlhM-GFP, the recovery of fluorescence at the rotational center was observed after the complete-photobleaching. On the other hand, in a cell producing GFP-FlhG, the

recovery of fluorescence was not observed. From the complete- and partial-photobleaching procedures for the cells producing of GFP-FlhN, the rate constant of FlhN-dissociation was calculated as 0.0007 s⁻¹. Thus, FlhN and FlhM, at least, could be exchanged in a functioning flagellar motor, and FlhN subunits would be exchanged in tens of minutes. These results would mean that a bacterial flagellar motor is not a stable structure and exchanges their component flexibly even in functioning state.

2TA4-02 高温におけるキメラべん毛モーターの回転トルク応答

2P-132 Torque response of bacterial flagellar motor induced by transient heating.

Yuichi Inoue (1), Hajime Fukuoka (1), Akihiko Ishijima (1) ((1) Tohoku University)

We have demonstrated the torque-speed curve of Na⁺-driven chimeric flagellar motor in *E. coli* (Inoue et al., 2008). The chimeric motor produced almost constant torque at constant temperature, as reported for WT motor in *E. coli*. We found that transient heating over 40°C can induce torque change in a reversible fashion. To understand the mechanism of the thermal response of torque, we examined effect of high temperature on the chimeric motor using back-focal-plane interferometry.

Heating at 40-50°C for 2-5 min induced reduction of motor torque (reducing phase) and cooling to room temperature induced resurrection of torque (resurrection phase) mostly in several minutes. In the resurrection phase, torque recovered in a stepwise manner. Histograms of resurrection phases measured with 0.35-1.0 μm bead showed several peaks up to 11 with unitary torque of 190-210 pN nm. The results are consistent with the report showing that the motor can be equipped with ~11 torque generating units (Reid et al., 2006). Reducing phase of torque also showed similar peaks in the histograms with unitary torque of 180-200 pN nm.

These results suggest that active number of torque generating units in a motor can be controlled by manipulating temperature. Thermal response of torque may be a key to understand the dynamics of motor components and the mechanism of torque generation.

2TA4-03 F₁-ATPase のストールトルク

Stall torque of F₁-ATPase

Shoichi Toyabe (1), Takahiro Watanabe-Nakayama (2), Tetsuaki Okamoto (1), Seishi Kudo (3), Eiro Muneyuki (1) ((1) Faculty of Engineering, Chuo University; (2) Grad. School of Bioscience and Biotechnology, Tokyo Institute of Technology; (3) Faculty of Engineering, Toin University of Yokohama)

F₁-ATPase is a rotary molecular motor; hydrolyzing ATP to ADP and phosphate, the central γ subunit rotates unidirectionally. On the other hand in cells, it works as a part of F₀F₁-ATP synthase and rather synthesizes ATP from ADP and phosphate with rotating in the opposite direction of hydrolytic rotations. Thus, it is crucial to study the rotations in the synthetic direction to reveal the nature of F₁-ATPase. Here, we applied an external torque to F₁-ATPase in the synthetic direction using an electrorotation method. We found that F₁-ATPase shows stepwise rotations in the synthetic direction at a sufficiently strong torque in the presence of ATP, ADP, and P_i. The torque necessary to rotate the wild type F₁-ATPase in the synthetic direction times 120° was almost equal to the free energy change of a single ATP hydrolysis under various conditions. On the other hand, smaller torque was enough to rotate mutated F₁-ATPase (βG181A, T165S, Y341W) in the synthetic direction.

2TA4-04 チューブリンC末端の伸長した構造が弱結合に果たす役割

2P-134 The role of extended C-terminal tail of tubulin on weak-binding interaction between motor proteins and microtubules

Itsushi Minoura (1), Seichi Uchimura (1), Masashi Degawa (1), Etsuko Muto (1) ((1) RIKEN Brain Science Institute)

The mechanism of weak binding between motor proteins and microtubules (MTs) is not fully understood in spite of previous kinetic, structural and motility studies. To get a comprehensive picture of the weak-binding interaction, we combined two methods: simulation of the weak-binding interaction using positively charged nanoparticles (made of polyacrylamide gel, ~60 nm in diameter) and MTs, and in vitro motility assay using mutant MTs. The charged particles show one-dimensional (1D) Brownian motion on MTs in a charge-dependent manner. Analysis of this interaction using mutant MTs revealed that the C-terminal tail (CTT) of tubulin is not essential for 1D Brownian motion of the charged particles, but the duration of Brownian motion was significantly reduced by the deletion of the CTTs. There are two possible interpretations for this observation: 1) 1D Brownian motion of the charged particles was prolonged merely due to the negative charges on the CTTs, and 2) prolonged interaction of the charged particles with MTs is a result of both charge effect and the extended structure of the CTTs. According to the latter interpretation, the structure of CTTs extending into the solution may prevent the charged particles from firmly attaching to the surface of the MTs. To confirm kinds of these interpretations is true, we are studying the motility of the particle/kinesin along those mutant MTs having the