(DPAMPS) by pulse field gradient spin echo (PFG-SE) NMR measurements. And we found out the relation between the DPAMPS and concentrations of PAMPS solutions. NMR measurements were carried out using ECA-600 (JEOL) at 30 ± 0.05 degrees. NM-60THAT/FG2 probe for solutions were used. And all experimental data were measured by BMS-005J micro cells (SHIGEMI). T1 in DW was measured by using inversion recovery measurements. We found that value of T1 was about 3.58[s]. Proton diffusion coefficients in DW (Dw) estimated by Stejskal-Tanner's equation were 2.33×10^{-9} [m²/s]. Values of DPAMPS increased with increasing concentrations of PAMPS solutions. The maximum value of DPAMPS was about 4 percent higher than Dw at PAMPS concentration of 2.14[mg/ml]. Then DPAMPS decreased monotonically in the concentration range from 2.14 to 29.9 [mg/ml]. As a result, we found higher proton diffusion coefficients in PAMPS solution than that in bulk water in the concentration range from 0.58 to 14.4 [mg/ml].

1P-112 リン酸中和反応のエントロピー変化に対する誘電水和解析および ATP 加水分解系への応用

Dielectric hydration analysis of the negative entropy change for the neutralization of sodium dihydrogenphosphate solutions and application for ATP hydrolysis system

George Mogami (1), Takashi Miyazaki (2), Makoto Suzuki (1) ((1) Dept. of Materials Processing, Graduate School of Engineering, Tohoku University; (2) Industrial Technology Center of Wakayama Prefecture; (3) CREST, Japan Science and Technology Agency)

Of various unsettled issues of fundamental importance in biological science is the molecular origin of a large negative Gibbs energy change ($\Delta_{\text{hyd}}G^\circ < 0$) for hydrolysis of ATP into ADP and inorganic phosphate (Pi). George et al. (*BBA* 223:1, '70) proposed that the very difference in the hydration energy between the reactants (ATP + H₂O) and products (ADP + Pi) would principally contribute to the $\Delta_{\text{hyd}}G^\circ$. The neutralization of Pi coincides with ATP hydrolysis reaction, in which the negative entropy change ($\Delta_{\text{neutral}}S^\circ < 0$) has not been yet explained. The thermodynamics of this phosphate buffer reaction was reexamined using hydration analysis of solute ions by microwave dielectric spectroscopy. The high-resolution complex dielectric spectra of NaH₂PO₄, Na₂HPO₄ and NaOH were measured. Using a novel microwave technique the dielectric relaxation properties and size of the hydration layer around solutes were determined from those spectra, expressed as a series of Debye relaxation functions, and the spectrum of bulk water. For all three solutes both hyper-mobile water, having a higher relaxation frequency f_c than that of bulk water f_{cw} , as well as constrained water, with $f_c < f_{\text{cw}}$, were observed around the solutes. After the reaction the dispersion intensity of the constrained water decreased by 2.9, corresponding to a positive entropy change. On the other hand, a marked reduction (-7.8) in the dispersion intensity of hyper-mobile water was also observed. These results deepened the understanding of the negative entropy change associated with this reaction. This analysis applied to ATP hydrolysis system.

1P-113 Brownian motor in muscle mechanics

Lorenzo Marcucci (1), Tetsuya Shimokawa (1), Mitsuhiro Iwaki (1), Toshio Yanagida (1) ((1) Osaka University, Graduate school of Frontier Biosciences, Soft Biosystem Group)

Recent experiments performed on single myosin head in our laboratory have shown interesting and unusual features of the actomyosin interaction, leading to the new possible interpretations of the Brownian Search, to find the optimal position of the strongly attached heads, and of the Strain Sensor mechanism, to switch from a weak bound state to a strong bound state.

This report investigated the possible consequences of such mechanisms on the macroscopic behaviour of the skeletal muscle perturbed from the isometric contraction condition.

The investigation is performed by using numerical simulation techniques from the stochastic description of the whole sarcomere view as a set of myosin heads interacting in parallel with an actin filament. Both Brownian Search and Strain Sensor mechanisms can be described in a natural way in the Brownian Ratchet theory in order to provide a theoretical model to explain "smart" actions performed by the cross bridge in an uncorrelated noisy environment.

The hypotheses can be confronted with the classical hypotheses of Huxley Huxley '54 and Huxley and Simmons '71 models, giving an interpretation to their structural basis.

1P-114 アクチン-ミオシン硬直複合体の高分解能 3次元像の獲得

Obtainment of high resolution 3D map of actin-myosin rigor complex

Yoshihiro Tsukada (1), Takeyuki Wakabayashi (2), Takuo Yasunaga (1) ((1) Dept Bioscience and Bioinformatics, Computer Science and Systems Engineering, Kyushu Institute of Technology Graduate School; (2) Dept Biosciences, School of Science and Engineering, Teikyo University)

Myosin and actin are understood to generate muscle contraction with ATP hydrolysis. Myosin exhibits the activity even in the absence of actin, but the interaction with actin promotes the activity. Some types of the structures of myosin without actin have been solved by X-ray crystallography. The detailed

observation of the actin-myosin complexes, however, has not been reported yet and so the interaction mechanism is still unclear; high resolution EM maps in the complex state are required for structural biological discussions at the amino acid level. Previous studies using helical symmetry could not show the structure at the high resolution beyond 15 Å. We, here, combined electron cryo-microscopy with the image processing called helical particle analysis and obtained the three-dimension map of the actin-myosin rigor complex from the same data set as that of the 3D reconstruction with helical symmetry. The map has wholly high resolution beyond 10 Å by FSC 0.16 and partly showed more highly resolved structure so as to discriminate the pitch of α -helix and individual β -sheet of SH3 domain. In the map, α -helices exposed in the upper 50K domain are detectable position and changed in comparison of that of myosin only by X-ray crystallography. To obtain higher resolution and correct the map, we are currently increasing filament data and improving the image processing. Improved maps will demonstrate some unexplained points of actin-myosin interaction mechanism.

1P-115 アクチンフィラメントの滑り運動と形態の歪みの関係

Relationship between the sliding movement of actin filaments and their morphological distortions

Kouji Oota (1), Shigeru Sakurazawa (1) ((1) Graduate School of Systems Information Science, Future University-Hakodate)

An actin filament sliding in an in vitro motility assay system ruptures frequently. The rupture could be interpreted as a sign of deterioration of a protein's biological activity. On the other hand, the frequency of the rupture of the sliding actin filaments depends on ATP concentration. We found that the frequency tended to become high at low ATP concentration condition. And also, as the result of image analysis, local distortion could be seen around the rupture point right before the rupture. From these facts, two suggestions follow: ATP hydrolysis of myosin causes local stress and structural distortion in an actin filament. Secondly, smooth movement of an actin filament at high ATP concentration is realized as a result of temporal and directional coordination of these local stresses. We thus assume that some fluctuations occur at the onset of sliding movements. To verify our assumption, we observed the actin filaments which were starting to move right after adding ATP under microscope. From the result, we found that the actin filament fluctuated and was distorted before starting to move. It might be thought that there is some relationship between sliding movements and the coordination of their distortions caused by ATP hydrolysis, which was made to avoid the rupture of a filament structure.

1P-116 骨格筋活性化におけるトロポニン構造変化と細胞内カルシウム濃度の同時測定

Simultaneous measurements of the troponin structural change and the intracellular Ca²⁺ concentration during skeletal muscle activation

Tatsuhito Matsuo (1), Naoto Yagi (1) ((1) SPring-8/JASRI; (2) Department of Engineering Science, Osaka University)

It is well known that skeletal muscle contraction is regulated by the intracellular Ca²⁺ concentration ([Ca]), especially by the Ca²⁺ binding and dissociation process on troponin molecules. However, the detailed relationship between the structural change of troponin and the Ca²⁺ binding-dissociation process is not clear. To study it, time-resolved x-ray diffraction experiments combined with the fluo3 (a Ca²⁺-indicator) fluorescence measurement were carried out at BL40XU in SPring-8 using overstretched frog skeletal muscles as specimens with the temperature 16°C. X-ray patterns and fluo3 signals were recorded at a time resolution of 1msec and 0.1msec, respectively. The structural change of troponin can be monitored through the meridional x-ray reflection intensity at 1/38.5nm⁻¹ (I_{38.5}) and the change of [Ca], can be monitored by the fluo3 signal, which was later converted to the change of the fraction of troponin that bound Ca²⁺ ([CaTn]) using published rate constants between troponin and Ca²⁺. I_{38.5} began to increase at 4 msec after the stimulus and reached the peak at 10 msec. Then it started to decrease with a half-time of 25 msec. On the other hand, [CaTn] began to increase at 1 msec and reached the peak at 5 msec, then decreased with a half-time of 40 msec. These results indicate that troponin molecules on the thin filament undergo the structural change only after most of them have bound Ca²⁺, and return to the resting conformation even if some still have bound Ca²⁺, showing the high cooperativity in the regulatory system.

1P-117 アクチンのヌクレオチド結合領域のゆらぎ

Conformational fluctuation in the nucleotide binding region of actin

Atsushi Ooi (1), Tsuyoshi Okagaki (1) ((1) Graduate School of Bioresources, Mie University)

Actin binds one nucleotide and one divalent cation in a cleft between two domains. Both nucleotide and cation bound to actin are freely exchanged with nucleotide or cation in solution. In most crystal structures of actin, the bound nucleotide is deeply buried within the cleft. Therefore it is thought that some conformation change to enable exchange is necessary. In order to understand molecular basis of the conformational change, we have investigated structural