chemical fixation with paraformaldehyde (PFA) and/or glutaraldehyde (GA), which suppresses the diffusion of membrane molecules, blocking their antibody-induced crosslinking and aggregation. However, several recent studies suggested that standard chemical fixation protocols often fail to block or sometimes even enhance antibody-induced clustering of putative raft-associated molecules, confusing raft researchers. Here, using single fluorescent-molecule tracking, we found that, even under extensive fixation conditions with 4%PFA + 0.1% GA for 30 min at 24 °C, GPI-anchored proteins and a phospholipid (Cy3-DOPE) could not be effectively immobilized whereas more than 90% of Lyn, H-Ras, and transferrin receptor (a transmembrane protein) were immobilized. An increase of the GA concentration to 0.2% was required to fix > 90% GPI-anchored proteins. Without GA, only about 61% H-Ras and 32% transferrin receptor were immobilized, whereas more than 90% of Lyn was immobilized. These results indicate that, in interpreting immunolocalization data, the presence of large fractions of mobile molecules, particularly GPI-anchored proteins and lipids, has to be considered, especially when the specimen is fixed without GA, which tends to abolish the antigen sites and to give higher fluorescent background.

1P-151 神経—免疫相互作用における接着分子 CADM1 の機能解析

Cell Adhesion Molecule 1 (CADM1) Works in Two Ways in Neuro-Immune Interaction

Tadahide Furuno (1), Keisuke Okamoto (2), Miho Sekimura (2), Akihiko Ito (3), Naohide Hirashima (2), Mamoru Nakanishi (1) ((1) School of Pharmacy, Aichi Gakuin University; (2) Graduate School of Pharmaceutical Sciences, Nagoya City University; (3) Institute of Medical Science, The University of Tokyo)

Cell adhesion molecules are important to communicate between nerve and immune cells (mast cells). However, the adhesion molecules involved in their communication has not been fully identified yet. We previously showed that cell adhesion molecule 1 (CADM1) expressed on both mast cells and sympathetic superior cervical ganglia (SCG) promoted their attachment and communication via the homophilic binding in an *in vitro* coculture approach. Here we have studied the role of CADM1 in the interaction between sensory dorsal root ganglia (DRG) and mast cells. DRG hardly expressed CADM1, but CADM1 on bone marrow-derived mast cells (BMMCs) was localized in the contact region. An IC-2 mast cell line, which lacked CADM1, was defective in attachment to DRG, while transfection with CADM1 recovered to the similar level to BMMCs. The efficiency in calcium response of attached CADM1-transfected IC-2 cells (IC-2^{CADM1} cells) after specific activation of DRG was significantly higher than that of IC-2 cells and was as similar as BMMCs. Then, we searched the binding partner molecule of CADM1 on DRG, and found that nectin-3 was expressed on DRG and was colocalized with CADM1 on BMMCs in the attached area. A neutralizing anti-nectin-3 antibody inhibited both the attachment and calcium response of BMMCs and IC-2^{CADM1} cells. These indicated that CADM1 on mast cells promoted the attachment and functional interaction to DRG via the heterophilic binding to nectin-3 on neurites. Thus, we conclude that CADM1 works in two ways in nerve-mast cell communication.

1P-152 アクチン調節タンパク質 zyxin のストレスファイバーへの張力依存的会合

Tension-dependent association of the actin-regulatory protein zyxin with stress fibers

Hiroaki Hirata (1), Hitoshi Tatsumi (2), Masahiro Sokabe (3) ((1) Cell Mechanosensing Project, SORST, Japan Science and Technology Agency; (2) Dept. of Physiology, Graduate School of Medicine, Nagoya Univ.; (3) Cell Mechanosensing Project, SORST, Japan Science and Technology Agency; Dept. of Physiology, Graduate School of Medicine, Nagoya Univ.)

Adherent cells including fibroblasts and endothelial cells adhere to substrate through focal adhesions (FAs) in which termini of actin stress fibers (SFs) are anchored. In response to changes in the tension along substrate-FA-SF, cells alter their organizations of FAs and SFs. We recently reported that the tension enhances local actin polymerization at FAs, and that the tension-dependent accumulation of the actin-regulatory protein zyxin at FAs is required for the actin polymerization. On the other hand, the tension has been reported to thicken SFs, while the molecular mechanism underlying this phenomenon remains unrevealed. As actin molecules steadily incorporate into and dissociate from SFs, the tension-dependent thickening of SFs may be caused by facilitated actin incorporation into SFs. Thus, we hypothesized that zyxin is involved in this process, and investigated in this study whether tension facilitates the accumulation of zyxin along SFs. In control fibroblasts, zyxin localized along SFs in a sarcomeric manner. When the substrate where cells adhered was uniaxially stretched by 150% for 5 min, large patches of zyxin accumulations on SFs appeared. In contrast, when tension in SFs was reduced by inhibiting the actomyosin interaction with the myosin II inhibitor blebbistatin, zyxin accumulation on SFs was disappeared accompanied by SF thinning. These results suggest that tension in SFs is crucial for the regulation of the zyxin localization on SFs, which implies that zyxin is involved in the tension-dependent regulation of actin polymerization along SFs. Based on this and our previous studies, we propose that zyxin acts as a common regulator of tension-dependent actin polymerization at FAs and along SFs.

1P-153 アクチンフィラメントの運動自由度の制限はコフィリンの結合頻度を低下させる

Restricted motion of actin filament reduces binding of cofilin to the filament

Kimihide Hayakawa (1), Yusuke Hunato (2), Hitoshi Tatsumi (2), Masahiro Sokabe (2) ((1) Cell Mechanosensing Project, SORST, JST; (2) Dept. of Physiology, Nagoya University Graduate School of Medicine)

A number of reports show that mechanical tension affects the dynamics of the actin cytoskeleton, however, little is known about the underlying biophysical mechanisms. We have demonstrated that tension in a single actin filament inhibits the severing of actin by cofilin using an *in vitro* assay system. This suggests that mechanical status of actin filament affects the binding and/or severing activity of cofilin. EM studies show that the binding of cofilin to the actin filament tilts the actin subunit and alters the pitch of twist in the large fragment of the filament. These studies suggest the cooperative binding of cofilin to the filament. The increase in the tilt angle by cofilin binding is in the range of the natural fluctuation of the tilt angle of pure actin filaments, implying that cofilin preferentially bind to the actin subunit in twisted states, and it is plausible that tension in the actin filament restricts the fluctuation of the tilt angle of actin subunits and reduces the susceptibility of actin to cofilin binding. To test a hypothesis that the fluctuation of actin filaments affects the susceptibility of actin to cofilin binding we tethered actin filaments on a cover-slip and cut the filaments by a glass capillary to create filaments with free end, and examined the binding of IAF-cofilin to these actin filaments by TIRF microscopy. The rate of binding of cofilin to the actin filaments near the cutting wedge was increased and the duration of binding was prolonged, suggesting that restrictions of the fluctuation of actin filaments by the tethering to the cover-slip decreased the susceptibility of actin to cofilin binding.

1P-154 異なったキネティクスを持った経路が乳腺における ATP 放出に関わっている

Multi Pathways with Different Kinetics Contribute to ATP Releases in Mammary Epithelial Cells

Kishio Furuya (1), Kyoko Harada (1), Masahiro Sokabe (2) ((1) JST, Cell Mechanosensing; (2) Graduate School of Medicine, Nagoya University)

ATP is now recognized as a ubiquitous extracellular messenger in most cells and tissues. Mammary epithelial cells are highly sensitive to mechanical stimulations and release ATP. The released ATP surrounds mammary alveoli and acts as autocrine and paracrine mediators on the secretory and myo- epithelial cells. However, it is not yet clear how ATP is released. Here, we investigate the mechanism of ATP release using a real-time ATP bioluminescence imaging system under microscope. ATP releases were observed by mechanical stimulations, by a hypotonic stress, by ligand stimulations, and also spontaneously that were enhanced in Calcium-free solution. These ATP releases possess different kinetics and pharmacological properties, indicating contribution of different pathways in the cells. Multiple pathways may work concomitantly to achieve mechano-purinergic coupling in mammary glands.

1P-155 細胞骨格に作用する力が細胞骨格の調節タンパクの結合性に与える影響の熱力学的検討

Thermodynamic investigation of modulation of binding affinity between cytoskeleton and its regulatory proteins in response to mechanical stresses

Yasuhiro Inoue (1), Taiji Adachi (1), Masaki Hojo (2) ((1) Kyoto University; RIKEN; (2) Kyoto University)

In dynamic cellular behaviors including migration, growth, and differentiation, cell shape and cytoskeleton structure are changed by mechanical stresses which are either transmitted to cells from the extracellular matrix or are generated within the contractile cytoskeleton of individual cell. These mechanical stresses are transduced to chemical signals leading to modulations of signaling pathways. Thus, the mechanical stress plays crucial roles in controlling the dynamic cellular behaviors while the mechanism of the transduction from mechanical stresses to chemical signals remains elusive. To better understand the dynamic cellular behaviors, it is crucial to elucidate how cytoskeleton structure adapts in response to mechanical stresses. In particular, because binding affinity is an integral of molecular interactions and entity of signaling pathways, it is important to understand how mechanical stresses modulate binding affinity between cytoskeletal proteins and their regulatory proteins. In this study, to relate stress-induced conformational changes of actin filament (F-actin), of which mechanics have a central role in cytoskeletal structure, with binding affinity of actin-regulatory proteins, we introduce a thermodynamic model expressing binding affinity as chemical potential of the regulatory protein in the F-actin. Investigating binding affinity in terms of mechanical stresses arisen from tensile, torsion and bending of the F-actin, binding affinity is likely changed in response to these mechanical stresses in a certain condition.