various pH conditions. Flickery gating was observed in the negative potentials. Kinetics and open channel noise analyses of the single-channel current were performed. Based on our present data, we will discuss how the KcsA channel regulates ion conduction using two separate gates.

1P-194 サブミリ秒時間分解能での KcsA カリウムイオンチャンネル
開閉構造変化計測
Conformational Changes of Single-Molecular KcsA Potassium
Channels during Gating Recorded in a Sub-Millisecond Time
Resolution

Hirofumi Shimizu (1), Masayuki Iwamoto (1), Takashi Konno (1), Reo Sakakura (2), Yoshimitsu Aoki (2), Shigetoshi Oiki (1) ((1) *Department of Molecular Physiology and Biophysics, University of Fukui Faculty of Medical Sciences*; (2) *Department of Electronics and Electrical Engineering, Faculty of Science and Technology, Keio University*)

We had succeeded in recording global twisting motions of KcsA potassium channels during gating in a single molecule with Diffracted X-ray Tracking (DXT) method (1). The channels were fixed on a glass plate and a gold nanocrystal was attached to the other side as a probe. Synchrotron white X-ray was irradiated and a diffraction spot from the attached nanocrystal was recorded at video rate, which can be translated into the motions of channel. While this method was sensitive to detect motions of the channels (~0.1 degree), its time resolution was shortened to reveal the whole gating motion (30 frames per second). To overcome this problem we installed a new detecting system which was composed of x-ray image intensifier and high-speed camera. With this system the twisting motions during gating were successfully recorded at 5000 frames per second (200 μ s/frame) which is equivalent to the sampling rate in single channel current recordings. In this presentation we will discuss about the contribution of this development to revealing the gating motions in detail.

(1) H. Shimizu, M. Iwamoto, T. Konno, A. Nihei, Y.C. Sasaki, S. Oiki, *Cell* 137, 67-78, 2008

1P-195 T細胞認識における PUFAs の影響
Effects of disruption of lipid rafts by PUFAs on T cell recognition

Keiko Uda (1), Satoshi Komaniwa (1), Arisa Yano (1), Toshihiro Komatsu (1), Akifumi Hagi (2) ((1) *Department of Immunology, Kochi Medical School*; (2) *Otsuka Pharmaceutical Factory, Inc.*)

see ITA3-08

1P-196 FPGA を用いた 1 分子生体シミュレーター
Single Molecule Bio-Simulator on FPGA

Michihiro Hori (1), Masahiro Ueda (1) ((1) *Osaka University*)

Progressing in single molecule imaging technology, stochastic nature of signaling molecules has been revealed in living cells. This leads to a fundamental question on the intracellular signaling processes: how does the signaling system in living cells reliably operate under thermal and stochastic fluctuations for flexible and robust responses in fluctuating environments? For understanding the design principle and computational principle of the system, it is essential that the relationships between cell's behaviors on the macroscopic level and stochastic properties of single molecules on the microscopic level are clarified. To carry out the simulation of the information processing in a cell including stochastic behaviors of single molecules, we have developed a calculation method by utilizing field programmable gate arrays (FPGAs). FPGAs are the LSI devices which can change digital circuits easily by computer programming. FPGAs are more flexible than application-specific integrated circuits (ASICs), and FPGAs can process information faster than software-based calculation. In this study, we propose a method for FPGA implementation, in which stochastic nature of biochemical reactions and diffusion of single molecules can be included in the simulation of the information processing in a cell. We applied this method to the stochastic simulation of chemotactic signaling system.

1P-197 細胞内 1 分子解析に基づいた走化性応答における受容体-G 蛋白
質間相互作用のモデル
Two state model of G protein-coupled chemoattractant receptors for
directional sensing in chemotaxis

Yukihiro Miyanaga (1), Michihiro Hori (1), Tatsuo Shibata (2), Mashahiro Ueda (1) ((1) *Graduate School of Frontier Biosciences, Osaka University*; (2) *Department of mathematical and life science, Hiroshima university*)

Transmembrane signaling for directional sensing in chemotaxis of eukaryotic amoeboid cells is mediated by heterotrimeric G protein-coupled receptors. Early activation events of G proteins by the receptors have been explained by a two state model, in which the receptors have both high and low affinity for

chemoattractants, and the receptor with high affinity promotes activation of heterotrimeric G protein. We used single-molecule imaging techniques to reveal the intracellular dynamics of G proteins in living *Dictyostelium* cells, which provided direct measurement of the mobility on membrane. We found that receptor stimulation induced a slow down of G α subunits in their lateral mobility, attaining a half-maximum response at about 320 nM of a chemoattractant. The mobility shift was probably due to a complex formation between the receptor and G α subunit. This dose-dependence of the response corresponds to low affinity of receptor but is different from that of G protein activation that was obtained previously to be a half-maximum at about 10 nM, roughly high affinity of the receptor. To explain these observations, we extended the two state model, in which the receptor with high affinity serves as an activator for heterotrimeric G protein in an enzymatic manner while the receptor with low affinity serves as a binding site for the activated G protein, which have important implications in the mechanism underlying the generation of a spatiotemporal pattern of active G proteins on membrane for directional sensing in chemotaxis.

1P-198 トロンボポエチンシグナルの初期段階は脂質ラフトによって制御
されている
Initial thrombopoietin signaling is regulated by lipid rafts

Akihiko Sakamoto (1), Takashi Kato (2), Takashi Funatsu (3) ((1) *Graduate School of Pharmaceutical Sciences, The University of Tokyo*; *Center for Medical System Innovation, The University of Tokyo*; (2) *Faculty of Education and Integrated Arts and Sciences, Waseda University*; (3) *Graduate School of Pharmaceutical Sciences, The University of Tokyo*; *Center for NanoBio Integration, The University of Tokyo*; CREST, Japan Science and Technology Agency)

Thrombopoietin (TPO) is a cytokine that primarily regulates megakaryocyte maturation and platelet production. Binding of TPO to its receptor, MPL, causes a conformational change of MPL, which allows cross-phosphorylation of Janus kinase 2 (JAK2). Phosphorylated JAK2 then phosphorylates MPL and several downstream effectors including signal transducers and activators of transcription, mitogen-activated protein kinase, and phosphoinositide 3-kinase. Activation of these pathways promotes proliferation and differentiation of megakaryocytes. However, how TPO signaling is regulated has not been clarified. Here, we show that initial TPO signaling is regulated by the distribution of MPL on the cell surface. Fluorescently labeled TPO gradually clustered on the cell surface of MPL-expressing cells, followed by internalization. Cholesterol depletion by methyl- β -cyclodextrin inhibited both clustering and internalization of TPO. Furthermore, phosphorylation of JAK2 and MPL were dependent on clustering of TPO rather than its internalization. Cell fractionation by sucrose density gradients revealed that TPO induced a translocation of MPL and JAK2 from lipid raft fractions to nonraft fractions. These data suggest that the distribution of MPL on the cell surface is a key regulator of initial TPO signaling. We are now studying (i) a molecular mechanism for TPO-dependent translocation of MPL; and (ii) the behaviors of MPL on the cell surface by single molecule imaging.

1P-199 コントラスト変調と異常分散を用いた膜蛋白質結晶中の脂質二重
膜の可視化
Visualization of lipid bilayer in the crystal of the membrane protein
using x-ray contrast variation and anomalous scattering

Yoshiyuki Norimatsu (1), Junko Tsueda (1), Ayami Hirata (1), Shioh Iwasawa (1), Chikashi Toyoshima (1) ((1) *Institute of Molecular and Cellular Biosciences*)

see ITA3-07

1P-200 小脳核グルタミン酸作動性シナプスにおけるモノアミンを介した
修飾作用
Monoaminergic cross-talk action at the mossy fiber-deep cerebellar
nuclei synapses

Fumihito Saitow (1), Hidenori Suzuki (1) ((1) *Dept. Pharmacology, Nippon Medical School*)

Activity of the deep cerebellar nuclei (DCN) takes an important role in outputting processed information from the cerebellum. Modulatory effects of synaptic transmission are important for regulating the information flow. As reported previously, one of monoamine, 5-HT shows the inhibitory effects on both glutamatergic and GABAergic transmissions through the presynaptic mechanisms. In this study, we examined a role of other monoamines, noradrenaline (NA) and dopamine (DA) on the synaptic transmission in the DCN using rat cerebellar slices. We recorded stimulation-evoked EPSCs from DCN neurons using whole-cell patch-clamp recording. Bath application of 10 μ M NA or DA decreased the amplitude of glutamatergic EPSCs but not GABAergic IPSCs. Biophysical and pharmacological studies revealed that presynaptic α_2 -adrenergic (α_2 -AR) and D2-like receptors (D2R) were responsible for the inhibitory effects on the EPSC amplitude, respectively. Although ~40 % of synapses did not show the inhibitory effects by the application of a D2R agonist quinpirole, application of DA

decreased the amplitude of EPSCs in all cases. Regarding the extent of inhibitory effect, there was strong correlation between NA- and DA-induced inhibitory actions. Moreover, an α_2 -AR antagonist RS79948 partly inhibited the DA-induced inhibitory action. These results suggested that 1) the mossy fiber terminals innervating the DCN had heterogeneity in the expression of D2R; 2) DA affected not only D2R but also α_2 -AR at the mossy fiber terminals.

1P-201 固体 NMR 分光法による SWAP-70 PH ドメインの脂質膜界面における構造転移および機能の解析

A solid state NMR study of the structure and function of the SWAP-70 PH domain induced at the membrane interface.

Naomi Tokuda (1), Hitoshi Yagisawa (1), Yasuhisa Fukui (2), Satoru Tuzi (1) ((1) *Grad. Schl. Life Sci., Univ. Hyogo.*; (2) *Hoshi. Univ.*)

The pleckstrin homology (PH) domain is a rather small lipid-binding domain consists of 100-120 amino acid residues frequently found in the proteins included in the important cellular functions such as the cellular signal transduction. A core structure of the PH domain is a β -sandwich consists of seven anti-parallel β -strands and a C-terminal α -helix located at one of the splayed corners of the β -sandwich. The unoccupied splayed corner of the β -sandwich includes a specific binding site for phosphoinositide. One of the well-known functions of the PH domain is to mediate membrane localization of the protein through stereoselective recognition of headgroup of phosphoinositides. In this study, we investigated structural alteration the switch associated protein (SWAP)-70 PH domain at the membrane surface by using solid-state NMR spectroscopy. It is reported that SWAP-70 is translocated from the cytoplasm to the plasma membrane as a result of a specific binding of the PH domain to PtdIns(3,4,5)P₃, and subsequently transported from the membrane to the nucleus by Importin- α depend on the nuclear localization signal (NLS) involved in the C-terminal α -helix of the PH domain. In the nucleus, SWAP-70 is reported to contribute to the class-switching of B lymphocyte. The solid-state ¹³C NMR spectra of the membrane associated SWAP-70 PH domain revealed that a conformational transition of the C-terminal α -helix from the α -helical structure to the random coil is induced at the membrane surface. This conformational transition of the C-terminal α -helix would facilitate the recognition of the PH domain by Importin- α and subsequent nuclear localization as a result of a conformational alteration of the NLS from the α -helix to the water-exposed extended structure.

1P-202 サリニバクター-SRI シグナル伝達系を用いた、走光性大腸菌作成に向けた試み

Towards engineering of phototactic *Escherichia coli* by using sensory rhodopsin I and Che proteins from *Salinibacter ruber*

Daisuke Suzuki (1), Michio Homma (1), Yuki Sudo (1) ((1) *Division of Biological Science, Graduate School of Science, Nagoya University*)

Microbial organisms utilize light not only as energy sources but also as signals by which rhodopsins work as photoreceptors. Sensory rhodopsin I (SRI) is known as a dual photoreceptor that regulates both negative and positive phototaxis in the archaeon, such as *Halobacterium salinarum* [1]. SRI forms 2:2 complexes with a methyl-accepting transducer protein (HtrI) and transmits light signals to a two-component signal transduction cascade mediated by Che proteins, which controls the flagella motor motion. To understand the signal-relay mechanism, we tried to engineer *E. coli* having the activity of the phototaxis. Recently we characterized the first eubacterial SRI protein from *Salinibacter ruber* and named it SrSRI [3-5]. This is an advantage for the engineering because *E. coli* is eubacterium as well as *S. ruber*. In fact, SRI from *H. salinarum* does not work in *E. coli*. The genome sequence of *S. ruber* also predicts that the bacterium has HtrI gene under the control of the same promoter as SrSRI and one set of Che genes. We cloned these genes and succeeded in expressing them as a recombinant protein in *Escherichia coli* [5]. Interestingly we found that SrHtrI works as a receptor for thermo-taxis in *E. coli*, indicating that SrHtrI transfer the external signal to cytoplasm Che proteins from *E. coli*.

[1] Spudich, J.L. and Bogomolni, R.A. (1984) *Nature* [2] Kitajima-Ihara, T. et al. (2008) *J. Biol. Chem.* [3] Suzuki, D. et al. (2008) *Biochemistry* [4] Suzuki, D. et al. (2009) *J. Mol. Biol.* revised. [5] Sudo, Y. et al. in preparation.

1P-203 蛙坐骨神経の複合活動電位に及ぼすビスフェノールAの抑制作用 Inhibitory action of bisphenol A on compound action potentials in frog sciatic nerves

Kotaro Mizuta (1), Tsugumi Fujita (1), Hiroki Yamagata (1), Eiichi Kumamoto (1) ((1) *Department of Physiology, Faculty of Medicine, Saga University*)

Bisphenol A (BPA) is widely used as a monomer to produce polycarbonate plastics while being one of endocrine-disrupting chemicals owing to its toxicological actions on humans. Although BPA is well-known to have an estrogenic action, it has not been fully examined yet how BPA affects the nervous system. We investigated the effect of BPA on fast-conducting and Na⁺-channel

blocker tetrodotoxin (1 μ M)-sensitive compound action potentials (CAPs) recorded from the frog sciatic nerve by using the air-gap method. Treatment of the sciatic nerve with BPA (0.5 mM) for 20 min resulted in a decrease in CAP peak amplitude to about 40% of control. The CAP amplitude did not recover to control in nerves returned to drug-free solution for up to 1 hr, and remained to be about 65% of control. This reduction of CAP peak amplitude produced by BPA was dose-dependent, and a half-maximal inhibitory concentration (IC₅₀) value for BPA was 0.31 mM (Hill coefficient: 1.5). When compared with the actions of local anesthetics on CAPs, the efficacy of BPA in inhibiting CAPs was larger than those of procaine, lidocaine and cocaine (IC₅₀ = 2.3, 0.74 and 0.80 mM, respectively), smaller than that of tetracaine (0.013 mM) and similar to that of ropivacaine (0.34 mM). These results indicate that BPA has an ability to inhibit nerve conduction with an efficacy comparable to those of local anesthetics.

1P-204 ラット脊髄後角における TRPA1 チャネルの局所麻酔薬によるシナプス前性の活性化

Presynaptic activation by local anesthetics of TRPA1 channels in the rat spinal dorsal horn

Lian-Hua Piao (1), Tsugumi Fujita (1), Chang-Yu Jiang (1), Kotaro Mizuta (1), Hai-Yuan Yue (1), Masanari Inoue (1), Eiichi Kumamoto (1) ((1) *Dept. Physiol., Fac. Med., Saga Univ.*)

Although local anesthetics are recently reported to activate TRPV1 channels which are cloned and expressed in dorsal root ganglion neurons, this has not been examined yet in the central nervous system. We investigated the effects of lidocaine, procaine, ropivacaine and tetracaine on glutamatergic spontaneous excitatory synaptic transmission in substantia gelatinosa (SG; lamina II of Rexed) neurons of adult rat spinal cord slices by using the whole-cell patch-clamp technique. Bath-applied lidocaine dose-dependently and reversibly increased the frequency but not the amplitude of spontaneous excitatory postsynaptic current (EPSC) recorded at -70 mV in SG neurons. Lidocaine activity was unaffected by the voltage-gated Na⁺-channel blocker tetrodotoxin and the TRPV1 antagonist capsazepine, but was inhibited by the non-selective TRP antagonist ruthenium red. In the same neuron, the TRPA1 agonist allyl isothiocyanate and lidocaine both increased spontaneous EPSC frequency. Like lidocaine, tetracaine also increased spontaneous EPSC frequency in SG neurons. In contrast, procaine and ropivacaine did not increase spontaneous EPSC frequency. It is concluded that some kinds of local anesthetics activate TRPA1 in the SG to increase the spontaneous release of L-glutamate from nerve terminals.

1P-205 フルボキサミンの光異性化の理論的研究

Theoretical study of photoisomerization of fluvoxamine

Kei Odai (1), Tohru Sugimoto (2), Etsuro Ito (3) ((1) *Dept. of informatics and Media Technology, Shohoku Coll., Japan.*; (2) *Dept. of Humanities, Coll. of Engin., Kanto Gakuin Univ., Japan.*; (3) *Lab. of Functional Bio., Kagawa School of Pharmaceutical Sci., Tokushima Bunri Univ. Japan.*)

Fluvoxamine, a selective serotonin re-uptake inhibitor (SSRI), is used as antidepressant/anxiolytic. Recently, photoisomerization of fluvoxamine has been reported to occurs from trans to cis isomer, with respect to the C=N double bond in its structure. Two chains of fluvoxamine implies the existence of many conformations, since there are 8 single bonds in the chains excepting two terminal single bonds with methyl group. The purpose of the present study is to disclose the photoisomerization paths and their activation energies by using a MNDO-AM1 molecular orbital method. The trans and cis isomers of fluvoxamine are constructed by twisting those 8 single bonds by the amount of 0, 120 or 240 degrees, and a total of 663 trans and 1796 cis isomers are obtained, excepting the isomers including steric hindrance. Their optimized structures and molecular vibrations are calculated to find that 191 trans and 1091 cis isomers among those isomers are in their ground states. The transition states of these ground states and then saddle point of the trans-cis isomerization of fluvoxamine are calculated, and our preliminary result shows that the activation energy of the trans-cis isomerization path of fluvoxamine is 2.9 eV (420 nm).

1P-206 モルモット盲腸筋層間神経叢培養グリア細胞の受容体、イオンチャネル及び膜電位に対する神経伝達物質の作用

Neurotransmitters act on receptors, ion channels, and membrane potentials in cultured glial cells from the myenteric plexus of the guinea-pig cecum

Kazuo Suzuki (1), Kazunori Ohta (1), Tomoharu Kimura (1), Syoichi Siga (1), Tomo Inoue (1), Daisuke Matsumoto (1), Mitsuyoshi Iwata (1), Masashi Sato (1), Katsuhiko Ikeda (1) ((1) *Department of Biomedical Engineering, Tokai University*)

The electrophysiological properties and the role of the glial cells obtained from an explant culture of the myenteric plexus of the guinea-pig cecum were studied using the patch-clamp technique. The myenteric glial cells were identified using fluorescent antibodies, anti-S100 and anti-GFAP. A polygonal shaped-cell was used for the patch-clamp recording. The cell membrane of the glial cells located in the cecum myenteric plexus had high- and low-conductance Cl⁻ channels which were insensitive to Ca²⁺ increases. Adrenalin or forskolin application evoked the