363 was expressed in E.coli BL21(DE3) and purified by Co-NTA column. Microtubule dependant ATPase assay was observed to have a rather low activity in contrast to conventional kinesins while its affinity for microtubule is similar. For the kinetic characterization of K23, we have employed fluorescent ATP analogue 2'-(3')-O- (N-methylanthraniloyl)-ATP (Mant-ATP). Fluorescence intensity of Mant-ATP did not alter significantly during hydrolysis by K23. On the other hand, we have successfully observed significant FRET between Mant-ATP and tryptophan in K23. Kinesin K23 has 3 tryptophans in its motor domain. After performing a structural comparison with conventional kinesins we learned that there is a possibility that the first Trp, located at the 21st amino acid is the one that works as a donor for FRET predominantly.

2P-113 Stopped Flow FRET を用いたミオシンクロスブリッチサイクルの構造解析

Analysis of Conformation of Skeletal Muscle Myosin Cross-bridge Cycle using Stopped Flow FRET

Masafumi Yamada (1), Koichiro Ishiyama (1), Eisaku Katayama (2), Yoshitaka Kimori (3), Shinsaku Maruta (1) ((1) Dept. of Bioinfo., Fac. of Eng., Soka Univ. : (2) Div. of Biomolecular Imaging, Inst. of Med. Sci., The Univ. of Tokyo: (3) Dept. of Ultrastr. Res., Nat'l Inst. of Neurosci., Nat'l Ctr of Neurol. and Psych.)

Previously biochemical studies have demonstrated that the highly reactive cysteine residues SH1 and SH2 can be crosslinked by variety of bifunctional reagents with different spans (3-14 Å) in the presence of nucleotides, suggesting that the region is highly flexible. The SH1-SH2 region is believed to play a key role in the conformational changes that occur in the myosin head during the force generation coupled to ATP hydrolysis. We have previously shown that the HMM, which SH1-SH2 was crosslinked by p-Phenylene-dimaleimide (pPDM) in the presence of ADP, have a novel conformation using quick freeze deep etch electron microscopy (QFDE-EM). We have also demonstrated that conformational states of myosin motor domain corresponding to several chemical steps of ATP hydrolysis, including S1•ADP•pPDM complex, can be monitored by measuring the FRET using fluorescent ATP analogue NBD-ATP. However, the conformational change of S1 interacting with actin has not been clarified yet. In the present study, we analyzed the conformation of the myosin interacting with actin using the stopped flow FRET between the fluorescent probes, BD and NBD which were incorporated into ATP binding site and the A1 essential light chain (ELC) respectively. The fluorescence intensity of donor BD at the ATP and ADP state in the presence of actin was higher than that in the absence of actin. These results suggested that the conformation of S1 was apparently different from when it binds to actin.

2P-114 アゾベンゼン誘導体によるキネシンの機能の光制御

Photo-regulation of kinesin function using azobenzene derivative

Takeshi Itaba (1), Masafumi Yamada (1), Shinsaku Maruta (1) ((1) Division of Bioengineering, Graduate School of Engineering, Soka University)

Azobenzene is a photochromic molecule that undergoes rapid and reversible isomerization between the cis- and trans-forms in response to ultraviolet (UV) and visible (VIS) light irradiation, respectively. Previously, we have introduced the sulfhydryl-reactive azobenzene derivative 4-phenylazophenyl maleimide (PAM) into the functional region of kinesin to reversibly regulate the ATPase activity of kinesin by photoirradiation. The five kinesin motor domain mutants, A247C, L249C, A252C, G272C and S275C, which contained a single reactive cysteine residue in loops L11 and L12 were prepared. The PAM-modified S275C and L249C mutants exhibited reversible alterations in ATPase activity accompanied by cis-trans isomerization upon UV and VIS light irradiation. In the present study, we prepared newly the six mutants (A21C, G26C, S66C, R16C, R25C, M96C) that have single cysteine at near the ATP binding site in order to regulate kinesin activity more effectively than previous mutants. The PAM-modified A21C, R25C, G26C and S66C mutants exhibited reversible alteration in ATPase activity accompanied by cis- trans isomerization upon UV-VIS light irradiation. However, regulatory efficiencies of the mutants were not significantly. Therefore, we prepared additionally four mutants (E200C, S202C, S203C, R204C) that have a single reactive cysteine residue in switch 1. Photoregulation of the ATPase activities of the mutants modified with PAM were also studied.

2P-115 キネシンにおける特有のループ L5 の機能的役割

The functional role of the unique loop L5 of kinesin

Kumiko Ishikawa (1), Keiko Tanaka (1), Shinsaku Maruta (1) ((1) Division of Bioinformatics, Graduate school of engineering, Soka University)

Recent crystallographic studies have shown that kinesin has striking structural similarity to the core of the catalytic domain of myosin. On the other hand, kinesin has several unique loops, L5 and L12. L5 is located near the ATP-binding site, and L12 in the microtubule-binding site, which are not present in myosin. The precise function of these loops remains unclear. However, it is known that Eg5 has elongated L5, to which monastrol binds and inhibits ATPase activity. We have previously shown that rice kinesin K16 has shortened L5 and exhibits unique enzymatic characteristics comparing with conventional kinesin. Therefore, the

loop may determine the characteristic properties of kinesin. In this study, we examined how loop L5 is related to the function of kinesin. First, we replaced the loop L5 of rice kinesin K16 and conventional kinesin KIF5A with that of Eg5. The engineered rice kinesin K16 that has elongated L5 showed no ATPase activity. On the other hand, the engineered KIF5A retained normal ATPase activity. However, the KIF5A mutant was not inhibited by monastrol. The results may suggest that these kinesins have different functions of L5.

2P-116 新規 ATP 誘導体の合成およびモータータンパク質との相互作用

Synthesis of novel fluorescent ATP analogue and interaction with nucleotide dependent motor proteins

Taro Kimura (1), Masafumi D. Yamada (1), Shinsaku Maruta (1) ((1) Division of Bioinformatics, Graduate School of Engineering, Soka University)

Several kind of fluorescent ATP analogues have been synthesized for the application to the kinetic studies of ATPase. However, some of the ATP analogues exist as a mixture of isomers. For instance, 2'-(3')-O-Mant-ATP have isomer of 2' and 3' in their ribose moiety and each isomer performs differently as substrate for the ATPases. In the present study, we have tried to synthesize novel fluorescent ATP analogues that have no isomer. The fluorescent ATP analogue 6-(N- (7-nitrobenz-2-oxa-1, 3-diazol-4-yl) amino) ethyl triphosphate (NBDTP) and N-methylanthraniloyl amino ethyl triphosphate (MANTTP) have been designed and synthesized, which are similar to non-nucleotide ATP analogue 2-[(4-azido-2-nitrophenyl) amino] ethyl triphosphate (NANTP). It is known that NANTP are good substrate for skeletal myosin and induces actin gliding in vitro motility assay. Excitation and emission maximums in the fluorescence spectrum of the ATP analogues were 474nm and 533nm for NBDTP, and 374nm and 430nm for MANTTP, respectively. The analogues showed ATP hydrolysis for conventional kinesin and skeletal myosin at the almost same level to that of ATP. Moreover, the analogues induced dissociation of acto-myosin. FRET between the ATP analogues and tryptophan in skeleton myosin were observed during ATP hydrolysis. Formation of skeletal myosin•ATP analogues•BeFn complexes was monitored with the fluorescence change of ATP analogues. The fluorescence changes of the ATP analogues during ATP hydrolysis have been also studied.

2P-117 X線小角散乱を用いたキネシンネックリンカーの構造変化の解析

Analysis of conformational change of kinesin neck-linker using small angle X-ray solution scattering

Keiko Tanaka (1), Yasunobu Sugimoto (2), Masafumi D. Yamada (1), Katsuzo Wakabayashi (2), Shinsaku Maruta (1) ((1) Division of Bioinformatics, Graduate school of Engineering, Soka University: (2) Graduate school of Engineering Science, Osaka University)

Recently, we have successfully dissolved the crystal structure of ADP bound K16 motor domain. The overall structure of the K16MD is similar to that of conventional kinesin motor domains, as expected from the high similarity of amino acid sequence. However, neck-linker region showed an ordered conformation in a position quite different from conventional kinesin. Previously, we have designed the K16 motor domain chimera protein fused with GFP at the neck-linker in order to monitor the conformational change of the neck-linker during ATP hydrolysis by small angle X-ray solution scattering. We determined the Radius gyration (Rg) values of K16-GFP in the presence or absence of nucleotides by X-ray solution scattering. The Rg of nucleotide-free K16-GFP was about 42 Å. In the presence of ADP and ATP, the Rg values were 38 Å and 39 Å, respectively. These results suggested that the neck-linker of nucleotide free K16 is in the docked conformation, on the other hand, the neck-linker of nucleotide bound state is in the novel conformation observed in crystal structure. In this study, conventional kinesin fused with GFP (KIF5A-GFP) was prepared and analyzed by small-angle X-ray scattering in order to compare its neck-linker conformation with K16-GFP. The conformational structure of KIF5A is well studied and expected to be good comparison for K16-GFP. KIF5A-GFP was expressed in E. coli. and purified with Co-chelate column. KIF5A-GFP showed normal microtubule dependent ATPase activity. Small-angle X-ray scattering analysis of the KIF5A-GFP has been performed and the Rg value was compared with that of K16-GFP.

2P-118 分子モーターの力発生におけるブラウンアン・ラチェット機構の寄与

Contribution of Brownian ratchet mechanism to the force-generation of molecular motors

Mitsunori Takano (1), Jun Narita (1) ((1) Dept. Phys, Waseda Univ.)

The force-generating mechanisms of molecular motors have been studied extensively but still remain controversial. For example, in the case of actomyosin, there has been a controversy between the so-called lever-arm mechanism and the Brownian ratchet mechanism. Although this controversy is not a mere superficial dispute but a truly essential controversy to understand how the molecular machines are designed to work robustly and efficiently under disturbing thermal noises, the study to settle this controversy seems to be stalled. One of the reasons of this stagnancy would be due to the technical difficulty to observe the molecular