Golgi cells and Lugaro cells. These two types have been characterized so far, but we have observed another type of interneurons; a globular-like neuron (GloLN), which has a small globular soma with several processes extending along the PC layer. Here, in order to identify interneurons underneath the PC layer by fluorescence microscopy we made cerebellar slices from a glutamic acid decarboxylase (GAD) 67/green fluorescent protein (GFP) knock-in mouse expresses GFP in GABAergic neurons. To establish electrophysiological criteria for the types of interneurons, we applied patch-clamp techniques and studied their electrophysiological properties, combined with pharmacological assays. We found that Golgi cells fired spontaneously at rest, but neither Lugaro cells nor GloLNs caused spontaneous action potential. Serotonin facilitated spontaneous firing in not only Lugaro cells but also Golgi cells and GloLNs. The frequency of spontaneous IPSCs (sIPSCs) recorded in GloLNs was much higher (~40 Hz) than those in Golgi (~2 Hz) and Lugaro cells (~20 Hz). Serotonin facilitated sIPSCs in Golgi cells but not in Lugaro cells or GloLNs. Membrane properties of GloLNs were different from Golgi and Lugaro cells, but similar to basket cells except for the membrane potential. Double-patch recordings demonstrated that sIPSCs in GloLNs and action potentials of PCs located close to the GloLNs were coincidentally detected, meaning that GloLNs could receive PC axon collateral inputs. These results suggest that GloLNs we observed here are likely to integrate signals from PCs.

3P-201 薬剤投与マウスの歩行指標の開発

New index for a behavioral differentiation across psychomotor stimulants

Hiroto Shoji (1), Yasuhito Nakatomi (5), Chihiro Yokoyama (3), Daiki Masaki (4), Kenji Fukui (4), Kazumitsu Hanai (6) ((1) *Kyoto Prefectural University of Medicine*; (2) *RESTO JST*; (3) *Functional Probe Research Lab, RIKEN Frontier Research System*; (4) *Dept Life-science, Kyoto Prefectural Univ of Medicine*; (5) *Dept Psychiatry, Graduate School of Medical Science, Kyoto Prefectural Univ of Medicine*; (6) *Dept Physics, Graduate School of Medical Science, Kyoto Prefectural Univ of Medicine*)

When mice are placed at an open-field, they explore mainly the peripheral zone of the open-field, which are called thigmotaxis. The thigmotaxis have been used as an index of anxiety in anxiogenic drug applied BALB/C mice. However, mice of another strains (C57BL/6 and DBA/2) walk mainly in center zone of the open-field after applying drug. That indicates the thigmotaxis cannot be adapted for the index for the anxiety in all the species. We have investigated the locomotor activity of cocaine-applied mice in an open-field (the circular field 60cm in diameter, with the wall 40cm high) test through the image analysis. Comparing the index of statistical physics concerning about the random walk behavior, we have investigated the characteristics of locomotor activity. We found plots of log(duration time of walking) vs log(numbers of walking event) are essentially linear with slopes which appear to depend on doses of applied cocaine. Then, we develop new index depending on obtained slopes. Our index can integrate locomotor behavior of all three strains (C57BL/6, DBA/2, BALB/c) without any contradiction.

3P-202 ヨーロッパモノアラガイの中枢神経系で見られる陰影反射の神経基盤

The neuronal substrates of predator induced stress on the shadow response in *Lymnaea*

Hiroshi Sunada (1), Manabu Sakakibara (1), Tetsuro Horikoshi (1) ((1) *Graduate School of Bioscience, Tokai University*)

In the pond snail *Lymnaea* a tactile or shadow stimulus induces an avoidance behavior, that is part of the whole body withdrawal (WBW) response. We tested at both the behavioral and electrophysiologic level whether WBW was altered in the presence of an ecologically relevant stress. Snails in the predator-exposed group had a significantly more pronounced and longer lasting WBW to the shadow stimulus compared to a control. Since RPeD11 is assumed to be the key interneuron controlling WBW, we focused on the activity of RPeD11. Presenting a shadow stimulus to the pneumostome area evoked an EPSP in RPeD11 in semi-intact preparations (without ocular photoreceptors). This EPSP is monosynaptic and is mediated by dermal photoreceptors in the pneumostome area. That is, the EPSP is present in hi Mg^{2+} /hi Ca^{2+} saline while it disappears in Ca^{2+} free saline. The amplitude and duration of the EPSP are significantly enhanced in the presence of the predator compared to naive preparations. These data strongly suggest that augmentation of sensory input to RPeD11 plays a key role in the enhanced WBW when predator is present.

3P-203 軟体動物腹足類における神経型 NO 合成酵素の同定

Gene Expression of Neuronal Nitric Oxide Synthase in the mollusk, *Lymnaea stagnalis*

Hisayo Sadamoto (1), Etsuro Ito (1) ((1) *Tokushima Bunri University*)

Nitric oxide is well known gaseous signaling molecule which works in numerous biological mechanisms, including retrograde transmission in the central nervous system (CNS). Generally, NOS proteins were classified into three groups, and each referred as inducible NOS (iNOS), endothelial NOS (eNOS) and

neuronal NOS (nNOS). The nNOS protein is different from other NOS proteins in possessing PSD/Disks-large/ZO-1 homologous (PDZ) domain at its N-terminus. The PDZ domain of nNOS is known to heterodimerize with other PDZ domain proteins in discrete subcellular regions, which is essential for anchoring nNOS at the synaptic sites. As well as in mammalian brain system, the regulatory mechanisms of NO have been investigated in the invertebrate CNS. In the pond snail *Lymnaea stagnalis*, previous reports showed the regulatory role of NO in feeding behavior and memory formation. More, histological analyses showed that the NADPH-diaphorase positive signal in the neuropil, indicating the subcellular localization of NOS proteins in the *Lymnaea* CNS. These findings were consistent with the idea that *Lymnaea* NOS proteins should be expressed as neuronal NOS, however, none of NOS related gene encoding the PDZ domain has been isolated. To start our study for analyzing the functional NOS involved in the synaptic regulatory mechanism, we first identified neuronal NOS gene with the PDZ domain. The deduced amino acid sequence (1645 AA) was highly identical with other species nNOS, and it was named as *Lymnaea* neuronal NOS (Lym nNOS). We also examined in situ hybridization for Lym nNOS mRNA and NADPH-d staining to examine the function of nNOS in the CNS.

3P-204 老化マウスの瞬目反射条件付け学習におけるニコチン性アセチルコリン受容体の関与

Contribution of nicotinic acetylcholine receptor to the classical eyeblink conditioning in aged mice

Takashi Kubota (1), Yutaka Kirino (1) ((1) *Tokushima Bunri University*)

Brain $\alpha 7$ nicotinic receptor ($\alpha 7$ nAChR) has become a therapeutic target for Alzheimer's disease (AD) based on their memory-enhancing and neuroprotective actions. In this study, we aim at revealing contribution of $\alpha 7$ nAChR on aging-related learning deficits using delay version of eyeblink classical conditioning. For conditioning in mice, 85 dB, 1 kHz tone was used as conditioned stimulus (CS) and 100 Hz electric stimulus to eyelid was as unconditioned stimulus (US). Conditioned response (CR) was evaluated by measuring myogenic potentials from eyelid. Interval from the CS onset to the US onset was fixed at 250 ms and the two stimuli terminated simultaneously. After seven days of conditioning sessions, 12-month-old (aged) mice showed a marked impairment in the percentage of conditioned responses (CR) compared with 8-week-old (young) mice. Treatment of selective $\alpha 7$ nAChR agonist, PNU282987, ameliorated the impairment of learning in aged mice. Central acetylcholinesterase inhibitor, Galanthamine, was more effective in the ameliorating learning deficit than PNU282987. It suggests that there may be another mechanism which contributes to the acquisition of delay CR without $\alpha 7$ nAChR signaling. The level of $\alpha 7$ nAChR expression was decreased in aged mice in the prefrontal cortex, hippocampus and cerebellum. Interestingly, it is known that cerebellum is essential to delay CR acquisition. These results suggest that impairment of acquisition of delay conditioning observed in aged mice caused by hypoactivity of $\alpha 7$ nAChR function.

3P-205 大腸菌発現系を用いた *salinarum* ハロロドプシンの機能的再構成

Functional Reconstitution of *salinarum* Halorhodopsin Expressed in *Escherichia coli*

Yasutaka Yamashita (1), Takashi Kikukawa (1), Kamiya Masakatsu (2), Tomoyasu Aizawa (2), Keiichi Kawano (1), Naoki Kamo (3), Makoto Demura (2) ((1) *Graduate School of Science, Hokkaido University*; (2) *Graduate School of Life Science, Hokkaido University*; (3) *College of Pharmaceutical Science, Matsuyama University*)

Halorhodopsin (HR) is a light-driven inward chloride pump expressed in the archaeal membranes. The representative HRs are from *Halobacterium salinarum* (HsHR) and *Natronomonas pharaonis* (NpHR). Despite of high sequence similarities, their characteristic differences were reported in chloride-dependent spectral shifts, other transportable anions, schemes and the chloride-concentration dependencies of the photocycles, and so on. In addition, the functional expression in *E. coli* membrane was reported for NpHR but not for HsHR. In this work, to facilitate the comparative studies between NpHR and HsHR, we established the procedures to prepare photoactive HsHR from *E. coli* cells and reconstitute purified HsHR functionally.

Expressed HsHR had no color in the *E. coli* membrane, but turned purple like in archaeal membrane after solubilization in the presence of all-*trans* retinal. The colored HsHR was highly purified and reconstituted in alectin liposomes. Light-induced outward directed transport of chloride by HR was evaluated by measuring passive proton efflux in the presence of uncoupler CCCP. By using this reconstitution system, the chloride-concentration dependence of the pumping activity was examined. From the relations between the pumping activity and the corresponding photocycle, we will discuss the chloride pumping mechanism of HsHR and the differences from NpHR.

3P-206 共鳴ラマン分光法によるファラオニスフォロドプシンN中間体の検出