initiation of polymerization and the strand displacement reaction by 3D^{pol}, which will be applied to the genomic diagnostics and the gene expression analysis. In this report, we evaluated the characteristics of the key enzyme 3D^{pol} synthesized in PURESYSYSTEM, Rabbit reticulocyte or insect cell-free protein synthesis system, regarding the effect of the post-translational modification.

1P101 FUNCTIONAL CHARACTERIZATION OF THE RECOMBINANT HUMAN TUMOR SUPPRESSOR 101F6 PROTEIN, A CYTOCHROME B561 HOMOLOGUE

Mariam Recuenco¹, Md. Motiur Rahman², Yoichi Sakamoto¹, Fusako Takeuchi³, Hiroshi Hori⁴, Sam-Yong Park⁵, Motonari Tsubaki¹, (¹Department of Chemistry, Graduate School of Science, Kobe University; ²Department of Molecular Science, Graduate School of Science and Technology, Kobe University; ³Department of Pharmacy, College of Pharmaceutical Sciences, Ritsumeikan University; ⁴Center for Quantum Science and Technology under Extreme Conditions, Osaka University; ⁵Protein Design Laboratory, Yokohama City University)

Human tumor suppressor 101F6 is a highly hydrophobic transmembrane protein and a member of the cytochrome b561 family. The 101F6 protein was expressed and purified from *Pichia pastoris* cells as a histidine-tagged protein with its molecular mass of ~26 kDa. MALDI-TOF-MS spectrometry confirmed the protein in its intact form. Heme content analysis by the pyridine hemochrome method indicated two hemes B per molecule. The midpoint potentials were found as +109 and +26 mV for the two hemes. These values were slightly lower than those reported for bovine adrenal chromaffin granule cytochrome b561 or plant *Zea mays* cytochrome b561. EPR spectra showed a highly anisotropic low-spin (HALS) signal at g_z=3.75 when measured at 5 K. At temperatures 15 K and 20 K, an EPR signal from a new HALS species with a different behavior in power saturation was overlapped at g_z=3.65. The EPR signal at g_z=3.16, previously seen in other cytochromes b561, was not observed. The lower homology in the amino acid sequence of 101F6 to other members of the cytochrome b561 family would account for the difference in the heme environments. Absence of the inhibition in the electron transfer activity from ascorbate upon DEPC-treatment of 101F6 protein was observed suggesting that 101F6 might accept reducing equivalent from ascorbate via a mechanism distinct from other cytochromes b561. It is very likely that these unique properties of 101F6 might be responsible for the function of this protein as a putative tumor suppressor during the development of human cancers.

1P102 T quaternary state of oxy haemoglobin in the presence of heterotropic effectors

Kenji Kanaori¹, Yusuke Tajiri¹, Antonio Tsuneshige², Takashi Yonetani³, (¹Kyoto Institute of Technology; ²Hosei University; ³University of Pennsylvania)

Oxygen affinity and its cooperative binding to hemoglobin (Hb) are related to change of the quaternary states; T (deoxy) and R (oxy) states. The binding of effectors to Hb decreased the oxygen affinity, and its cooperativity. In this study, in the presence of L35 and IHP, strong effectors, ¹H-NMR spectra of Hb were measured to clarify the correlation between the change of the oxygen affinities and the structural change of Hb by the effectors.

At pH 6.8, ¹H-NMR spectrum of deoxy-Hb in the presence of L35 and IHP showed four "T-state" signals in 10-14 ppm, originated from hydrogen bonds between Hb subunits, and hyper fine shifted (hfs) signals around 20 ppm, caused by high spin Fe(II) in the heme. With O₂ pressure increasing, the hfs signals disappeared, which means that O₂ binds to heme, but the four T-state signals were still observed under the O₂ pressure of 760 torr. In order to confirm whether O₂ is bound to the heme, visible spectrum of the NMR sample was measured, which coincides with that of oxy-Hb. Consequently, in the presence of the effectors, Hb formed T quaternary state at pH 6.8, though its heme takes 6-coordination with O₂. The pH dependence of the T (oxy) state was studied by measuring ¹H-NMR spectra between pH 6.8 and 8.0. The transition occurred at pH 7.4, and above pH 7.8, only two hydrogen bond signals remained like R (oxy). Hb in the presence of L35 and IHP takes T (oxy) state but not R (oxy) below pH 7.0, which related to the decrease of the oxygen affinity and its cooperativity.

1P103 ウシ心筋チトクロム酸化酵素の酸素還元中心への塩素イオン結合の可能性の X 線構造解析による検討

1G1255 An X-ray structural evaluation of the possibility of chloride binding to the O₂ reduction site of bovine heart cytochrome c oxidase

Tomitake Tsukihara^{1,2}, Michihiro suga², Kazumasa Muramoto², Kyoko Shinzawa-Itoh², Tomoko Maeda², Eiki Yamashita¹, Shinya Yoshikawa², (¹Institute for Protein Research, Osaka University; ²Department of Life Science, University of Hyogo)

The resting oxidized form of bovine heart cytochrome c oxidase (CcO) does not pump protons upon initial two electron reduction, while the fully oxi-

dized CcO under turnover conditions pumps protons upon receiving electrons. The OH⁻ binding to Fe_{a3}³⁺ in the oxidized form under turnover conditions has been established by a resonance Raman analysis. The electron density between Fe_{a3} and Cu_B in the resting oxidized form is consistent either to a bridging peroxide with the O-O distance of 1.7 Å or to one equivalent of chloride ion equally (50 % each) distributed between the two sites. In fact, the chloride binding to the resting oxidized form has been proposed by EXAFS. Establishment of the structure of the resting oxidized form would provide important clues for elucidation of the proton pump mechanism. Thus, for evaluation of the possibility of the chloride binding, an anomalous dispersion analysis for the resting oxidized bovine heart CcO was performed at 50K using low energy X-rays with the wavelength of 1.7470 Å, at which the anomalous dispersion of chloride is approximately as intense as those of S, Cu and Fe atoms. The anomalous map obtained shows clear peaks at the positions of the sulfur atoms of methionine and cysteine residues, Cu and Fe sites while no peak is detectable for the electron density between Fe_{a3} and Cu_B. These results clearly exclude the possibility that chloride binds to O₂ reduction site of the resting oxidized CcO, consistent with the chloride content analyses showing 0.15 Cl/CcO.

1P104 水素原子レベルでのプロトン輸送機構解明を目的としたチトクロム酸化酵素の高分解能 X 線結晶構造解析

High resolution X-ray structural analysis of cytochrome c oxidase to clarify the proton pumping mechanism at hydrogen atom level.

Naomine Yano¹, Kazumasa Muramoto¹, Tomoko Maeda¹, Kyoko Shinzawa-Itoh¹, Masao Mochizuki¹, Eiki Yamashita², Tomitake Tsukihara², Shinya Yoshikawa¹, (¹Department of Life Science, University of Hyogo; ²Institute for Protein Research, Osaka University)

Bovine heart Cytochrome c oxidase (CcO) in the mitochondrial respiratory chain is a terminal enzyme, which catalyzes the molecular oxygen reduction coupling the proton transfer across the inner membrane. To understand proton transfer mechanism, it is necessary to determine orientations and protonation states of amino acid residues (imidazole and carboxyl groups etc.) in the proton pathways. Therefore, we have analyzed the orientations of amino acids by the temperature factor refinement of each atom in the amino acid. The result indicates that if the temperature factor is lower than 30 Å², it is possible to determine the orientation of amino acids. If we can distinguish carboxyl group bond length and observe hydrogen atom electron density, it is possible to determine the protonation states of amino acids. For the purpose, we are currently carrying out the X-ray structural analysis of bovine CcO at 1.4 Å resolution. To improve data accuracy, the diffraction redundancy was increased by merging of many image data. Besides, we select good quality reflections judging from mosaicity and linear R-factor with the program Denzo and Scalepack.

1P105 Atomic Resolution X-ray crystallographic Analyses and

1G1325 Spectroscopic Studies of a Cytochrome c' from *Alcaligenes xylosoxidans*

Akiko Takashina¹, Masaki Unno², Takamitsu Kohzuma^{1,2}, (¹Institute of Applied Beam, Ibaraki University; ²Frontier Research Center for Applied Atomic Sciences, Ibaraki University)

Cytochrome c' (cyt c') are a class of c-type cytochrome, which have been found in several photosynthetic bacteria and denitrifying bacteria. Cyt c' has been known to be a typical four α-helices bundle structure, which attaches a heme group near the C-terminal region. The unusual magnetic properties of ferric cyt c' at neutral pH have been interpreted for the quantum mechanical admixture of an intermediate-spin (S = 3/2) and a high-spin (S = 5/2) state. The reduction potential of cyt c' from *Alcaligenes xylosoxidans* was reported to be 184 mV vs. NHE. Eady et al reported the NO and CO forms a stable 5-coordinate NO adduct and 6-coordinate CO adduct with cyt c', respectively. High quality crystal of cyt c' giving an atomic resolution data is required to add more precise knowledge to heme-protein chemistry and elucidate the small molecule recognition mechanisms of cyt c'. The optimized crystallization methods according to the crystal growth mechanism of *Alcaligenes xylosoxidans* cyt c' were reported by Kohzuma et al., and the high quality cyt c' crystals were obtained. The 0.92 - 1.11 Å resolution data were collected for the ferric, ferrous and NO-binding cyt c'. The atomic resolution structures of cyt c' from *Alcaligenes xylosoxidans* and spectroscopic properties will be presented and discussed.

1P106 Conformational Change of NOS Probed by Transpeptidase Mediated C-terminus Specific Fluorescence Labeling

Akihisa Kouzai, Shigetoshi Miura, (Department of biological Science and Technology, Tokyo University of Science, 2641 Yamazaki, Noda, Chiba, 278-8510, Japan)

Staphylococcus aureus expresses sortase (SrtA) that catalyzes the transpepti-