ing TM1 inner helices are tilted and crossed each other near the cytoplasmic side, forming the most constricted hydrophobic part of the pore called gate. Upon membrane stretch, the helices are dragged by lipids at Phe78 and tilted, accompanied by the outward sliding of the crossings. This led to a slight expansion of the gate associated with an exposure of oxygen atoms of the backbone to the inner surface of the gate. This allows water penetration in the gate and formation of hydrogen bonds between water and the exposed oxygen, which in turn weakened the hydrophobic interaction at the crossings, causing a further opening of the gate and water permeation.

2P118 化学ポテンシャルに基づいたバクテリオロドプシンのフォースカーブのシミュレーション Simulation of force-distance curve by chemical potential model for bacteriorhodopsin

Tatsuya Yamada, Shigeki Mitaku, (Department of applied physics Nagoya University)

Force-distance curve of membrane proteins by atomic force microscopy is due to the combination various interactions within membrane proteins and protein-lipid interactions. For example, the force-distance curve of bacteriorhodopsin has four significant peaks which should arise from some physical mechanism. For understanding the shape of the force-distance curves, we developed a novel method for Brownian dynamics simulation by chemical potential model. In this method we assumed that the chemical potential difference between dissociative amino acid side chain in water and in lipid membrane is the major factor for the peaks of force-distance curves of membrane proteins. We applied this simulation method to the force-distance curves of bacteriorhodopsin, and found that the force-distance curves by the simulation were very similar to experimental ones. Furthermore, we could calculate the force-distance curves of bacteriorhodopsin for drawing from loop segments other than the N- and C-terminal segments. We are planning to compare the force-distance curves of simulation and the corresponding experiment at various pH, which will be discussed at poster in detail.

2P119 Functionally rotating mechanism of a multidrug transporter studied by coarse-grained simulation

Xingqiu Yao¹, Hiroo Kenzaki¹, Shoji Takada^{1,2}, (¹Department of Biophysics, Kyoto University; ²CREST, Japan Science and Technology Agency)

Multidrug resistance has been a serious problem in the treatment of infectious diseases and cancers. The efflux system that is largely responsible for the drug resistance in *E. Coli* contains the transporter, AcrB. The structures of AcrB were solved first in 2002 as the symmetric homo-trimer¹, and then in 2006 as the asymmetric homo-trimer², latter of which suggested the functionally rotating mechanism. Due to the difficulty of *in vitro* experiment on AcrB, direct proof of such a mechanism is still lacking. In this sense, computational study of this protein becomes primarily important. Here, by molecular simulations, we have investigated several key questions about AcrB: What causes the symmetry-breaking that distinguishes the 2006's structure from that of 2002? How is the energy input (proton gradient) utilized by the protein to power the functional rotation coupled with drug export? By what uptake channel do drugs enter the binding pocket of the transporter from the outer leaflet of the membrane? In particular, we employed the multiple-basin model³ for the study, which is a residue-level coarse-grained model capable of reproducing large-amplitude conformational change effectively. The detail of results will be presented on the meeting.

Keywords: multidrug resistance, AcrB, membrane transport, coarse-grained, multiple-basin model

1 Murakami, S., et al. *Nature* **419**, 587-593 (2002).

2 Murakami, S., et al. *Nature* **443**, 173-179 (2006).

3 Okazaki, et al. *Proc Natl Acad Sci U S A* **103**, 11844-11849 (2006).

2P120 膜タンパク質系の分子動力学シミュレーションのための精密な構造モデリング法の開発 An accurate modeling of the protein-membrane complex structures

Takaharu Mori^{1,3}, Mizuki Morita^{2,3}, Shandar Ahmad^{2,3}, Kenji Mizuguchi^{2,3}, Yuji Sugita^{1,3,4}, (¹RIKEN Advanced Science Institute; ²National Institute of Biomedical Innovation; ³JST-BIRD; ⁴JST-CREST)

Recently, all-atom model molecular dynamics (MD) simulations have been widely used to understand the relationship between biological functions and structures of membrane proteins. In these simulations, careful modeling of the protein-membrane complex structures is essential, because the crystal structures of membrane proteins often lack two important information: 1) orientation of proteins with respect to the membrane plane and 2) structure of the surrounding lipid molecules. Although in the simulations these factors significantly affect the conformational stability of membrane proteins, no

standard protocols for structural modeling of the protein-membrane complex has been established. Here, we first developed a new method to predict an optimal orientation of protein using the Monte Carlo simulation with an implicit membrane model (GBMV), where the orientation of protein is optimized by minimizing the solvation free energy. By using this method, we successfully predicted the orientation of the SecY channel in the membrane, which was consistent with the results of the all-atom MD simulation (T. Mori et al., 2010). For the second problem, we developed a method to estimate surface area of bulk lipids around proteins, which is based on a Voronoi tessellation. By using this method, we successfully reproduced the bilayer structure, in consistent with the experimental data. We will also discuss the accuracy of our method by comparison with other database about membrane protein orientation (OPM and PDBTM) and coarse-grained MD simulations.

2P121 固体 NMR による膜に再構成した大腸菌由来 H⁺-ATP 合成酵素の構造解析 Structural analysis of the membrane-reconstituted subunit c-ring of *E. coli* H⁺-ATP synthase by solid-state NMR

Yasuto Todokoro^{1,2}, Masatoshi Kobayashi¹, Takeshi Sato¹, Toru Kawakami¹, Ikuko Yumen¹, Saburo Aimoto¹, Toshimichi Fujiwara¹, Hideo Akutsu¹, (¹IPR, Osaka Univ.; ²Nano. bio., Yokohama city univ.)

The subunit c-ring of H⁺-ATP synthase (F_oc-ring) plays an essential role in the proton translocation across a membrane driven by the electrochemical potential. To understand its structure and function, we have carried out solid-state NMR analysis under magic-angle sample spinning. The uniformly [¹³C, ¹⁵N]-labeled F_oc from *E. coli* (EF_oc) was reconstituted into lipid membranes as oligomers. Its high resolution two- and three-dimensional spectra were obtained, and the ¹³C and ¹⁵N signals were assigned. The obtained chemical shifts suggested that EF_oc takes on a hairpin-type helix-loop-helix structure in membranes as in an organic solution. The results on the magnetization transfer between the EF_oc and deuterated lipids was in good agreement with the cross-linking results previously reported by Fillingame and his colleagues. This agreement reveals that the reconstituted EF_oc oligomer takes on a ring structure similar to the intact one *in vivo*. On the other hand, analysis of the ¹³C nuclei distance of [3-¹³C]Ala24 and [4-¹³C]Asp61 in the F_oc-ring did not agree with the model structures proposed for the EF_oc-decamer and dodecamer. Interestingly, the carboxyl group of the essential Asp61 in the membrane-embedded EF_oc-ring turned out to be protonated as COOH even at neutral pH. The hydrophobic surface of the EF_oc-ring carries relatively short side chains in its central region, which may allow soft and smooth interactions with the hydrocarbon chains of lipids in the liquid-crystalline state.

2P122 新規コイルドコイルラベル法を用いた生細胞膜での受容体オリゴマー化解析法の開発 Stoichiometric analysis of receptor oligomerization in living cells using a novel coiled-coil labeling method

Yoshiaki Yano, Kaoru Omae, Katsumi Matsuzaki, (Graduate School of Pharmaceutical Sciences, Kyoto University)

Oligomerization of membrane receptors plays an important role in the regulation of receptor activity and trafficking. FRET and BRET techniques are often employed to detect receptor oligomerization in living cells by transfecting receptors genetically fused with fluorescent/luminescent proteins. However, the large size of the proteins often interferes the localization and function of target receptors. Furthermore, it is usually difficult to control the labeling ratio (donor/acceptor), which is important for quantitative analysis, when two fusion receptors with different colors are co-expressed. Posttranslational labeling methods using a short genetically encodable tag and a fluorescent probe that specifically binds to the tag enable a smaller size of label and easy control of labeling ratio in multicolor labeling. We recently developed a quick (< 1 min) and surface-specific tag-probe method using a high-affinity heterodimeric coiled-coil formation between the E3 tag (EIAALEK)3 attached to the target receptor and the K4 probe (KIAALKE)4 labeled with a fluorophore. Here we examine the oligomerization of the metabotropic glutamate receptor (mGluR) using this novel labeling technique. Stoichiometric analysis revealed that mGluRs transiently expressed in CHO cell membranes form dimers. The observed FRET signal did not depend on the expression level, indicating that the observed oligomerization was specific and concentration-independent. This method will be useful for analysis of oligomerization of various membrane proteins.

2P123 ミトコンドリア外膜タンパク Tom20-プレ配列複合体の分子動力学計算 Molecular dynamics simulation of outer mitochondrial membrane protein Tom20-presequence complex

Yasuaki Komuro¹, Takaharu Mori², Eiro Muneyuki¹, Takashi Saitoh³, Daisuke Kohda³, Yuji Sugita²,