located in β subunit which is thought to transition between open and closed conformations upon substrate binding and product release. It is widely accepted that cooperative open/close bending motions of the three β subunits are not only important for continuous ATP hydrolysis reaction but also essential for torque generation. However, it has been unknown how individual β subunit contributes in the torque generation.

As a clue for this issue, we measured the torque of hybrid F_1 that contains only one or two active β subunit(s) by using magnetic tweezers. In this system, torque can be measured through the angular deviation between magnetic trap center and magnetic bead attached to γ subunit during slow forced rotation of the bead. This method does not require spontaneous rotation for the torque measurement and enables us to measure the torque of the hybrid F_1 that is incapable of continuous catalysis.

Some of data obtained show that, during forced rotation in ATP hydrolysis direction, the hybrid F_1 generates positive torque at some angular regions and does work against magnetic trap potential only in the presence of ATP. This implies that the remaining one or two active β subunit(s) can generate positive torque with consumption of ATP.

2P185 暗視野観察とレーザートラップを用いたミオシン II カ発生 (素) 過程の計測
Observation of mechanical (elementary) process for myosin II by a dark-field microscopy combined with an optical trap

Mitsuhiro Iwaki¹, Atsuko Iwane², Toshio Yanagida², (¹ Graduate School of Medicine, Osaka University; ² Graduate School of Frontier Biosciences, Osaka University)

Myosin-II molecules are self-assembled and form a muscle. Individual myosin heads in muscle does not seem to interrupt each motion and fully exhibit their function. Myosin heads are connected to the backbone filament via S2-coiled-coil (≈ 60 nm) with non-linear elasticity. Motor domain undergoes Brownian motion (probably > 10 nm fluctuation), but small working stroke (< 10 nm) by the single myosin head is fully converted to macroscopic sliding motion of muscle. Thus, flexibility enabling myosin head's Brownian motion reduces molecular friction in muscle, but how small working stroke within the thermal noise can be fully converted to macroscopic motion is still unclear.

To answer the question, we should visualize the Brownian process and transition from pre-working state to post-working state. Detection for the process has succeeded using micro-needle method (Kitamura et al., 1999), but detail analysis at modulated conditions was hard due to the difficulty for the method.

Here we used dark-field microscopy combined with an optical trap and observed interaction between myosin II S-1 and actin. S1 was tagged with small optically-trapped particle (60 nm gold or 200 nm polystyrene bead), which enabled us to quantify non-processive steps and processive steps concerning a few myosin heads with rapid response time. We will discuss these results and trials of mechanical modulation to the myosin head at sub-milli second time scale to examine strain-dependent force generation like myosin-VI (Iwaki et al., 2009).

2P186 Single-molecule measurement of F_0F_1 rotation in living *E. coli*

Takuya Oosaka, Ryota Iino, Hiroyuki Noji, (The Institute of Scientific and Industrial Research, Osaka University)

In *E. coli*, there are two pathways of ATP synthesis, glycolysis and oxidative phosphorylation. F_0F_1 (ATP synthase) is rotary motor protein and responsible for the later pathway. In this study, we will measure the rotation rate of F_0F_1 and contribution to ATP supply in living *E. coli* under various culture conditions. Donor and acceptor fluorescent dyes are conjugated with the rotor and stator subunits of F_0F_1 and rotation rate in living *E. coli* will be measured by single-molecule Förster resonance energy transfer (FRET). There are three motives for this study. (1) F_0F_1 -knockout *E. coli* can grow in nutrient-rich medium, indicating that glycolysis is sufficient for ATP supply. (2) ATP concentration is considerably higher than that of ADP in the cell. This may counteract with ATP synthesis. (3) Proton motive force, especially $\Delta\mu_H$, across the cell membrane is not very large.

Firstly, SNAP-tag and CLIP-tag were introduced into the stator a-subunit and the rotor ϵ -subunit. Catalytic activity and specific labeling of purified F_0F_1 were confirmed. Furthermore, *E. coli* expressing F_0F_1 were reacted with fluorescent dye, and membrane localization of F_0F_1 were confirmed. Single-cell measurement showed the FRET level of ~ 0.5 , as expected from the structural model of F_0F_1 . The copy number of labeled F_0F_1 in *E. coli* were decreased by culturing the cell after labeling, and single-molecule F_0F_1 in *E. coli* was observed. Next, we will try single-molecule FRET measurement in living *E. coli*.

2P187 キネシンの機能性ループ L5 の修飾による ATPase 活性の制御
Control of kinesin ATPase activity by modification of functional loop L5

Kumiko Ishikawa, Keiko Tanaka, Shinsaku Maruta, (Division of bioinformatics, Graduate school of engineering, Soka University)

L5 is one of the unique loops located in the vicinity of ATP binding site of kinesin. In addition, it is believed that L5 is involved in the energy transduction. The novel rice plant specific kinesin K16 has several unique enzymatic characteristics comparing with conventional kinesin. The most interesting property is that the ADP-free K16 motor domain is very stable, contrast to conventional kinesin that is very labile in ADP-free state. The crystal structure of K16 revealed that the length of the loop L5 was much shorter than that of conventional kinesin. Therefore, the unique properties may be due to the short L5. We examined the influence of the change in length of L5 for K16. Microtubule dependent ATPase activity of K16 mutant that has a longest L5 of Eg5 was almost abolished. ATPase activity of K16 mutant that has no L5 was also abolished. On the other hand, K16 mutant that has L5 of conventional kinesin showed almost similar ATPase activity to K16 WT. The results suggested that the length of the L5 influences significantly in K16 ATPase activity. We have also prepared the two K16 mutants H101C and C214S, L5 of which was substituted by that of KIF5A. Mutants H101C and C214S that has a single reactive cysteine residue in the L5. Photochromic molecule of azobenzene derivative PAM was incorporated into the cysteine residue to induce conformational change of L5 by ultraviolet and visible light irradiation. The kinesin mutants modified by PAM alter the ATPase activity by ultra-violet and visible light irradiations reversibly.

2P188 フォトクロミック分子を導入した微小管によるキネシン ATPase の光制御
Photo-regulation of kinesin ATPase using microtubules modified with photochromic molecule

Koichiro Ishiyama, Takeshi Itaba, Hideki Shishido, shinsaku maruta, (Division of Bioinformatics, Graduate school of engineering, Soka University)

Azobenzene is a photochromic molecule that undergoes rapid and reversible isomerization between the *cis*- and *trans*-forms in response to ultraviolet (UV) and visible (VIS) light irradiation, respectively. Previously we have introduced the sulfhydryl-reactive azobenzene derivative 4-phenylazophenyl maleimide (PAM) into the functional region of kinesin and succeeded to regulate the ATPase activity of kinesin reversibly by photoirradiation. However, the regulation efficiency of the ATPase activity was not significant. It is well known that the contraction of skeletal muscle myosin is regulated by calcium binding to the rail protein actin. The regulating system achieves high efficient switching. Therefore, it is expected that introducing artificial switching system into the microtubules enables high efficient regulation of kinesin motor activity. In this study, we have done basic experiments on incorporation of photochromic molecule into microtubule. Quantitative analysis using DTNB revealed that 15 cysteine residues on the surface of alpha beta tubulin dimer. Interestingly the two of the cysteine residues were modified by fluorescent probe 5(6)-iodoacetamidomorpholinorhodamine at the microtubule state. In the preliminary experiment, approximately 10 cysteine residues were modified by PAM. Currently, we are determining the characterization of the tubuline dimer modified by PAM.

2P189 フォトクロミック分子を用いた低分子量 G 蛋白 Ras の光制御
Photocontrol of the small G protein Ras using photochromic molecule.

Hisashi Kobayashi, Hideki Shishido, Shinsaku Maruta, (Division of Bioinformatics, Graduate school of engineering, Soka University)

Small GTPase Ras is one of the G proteins. It was shown that the crystal structure of the GTPase domain strikingly resemble to that of ATP driven motor proteins myosin and kinesin. We have previously demonstrated that incorporation of photochromic molecules in the functional sites of kinesin enable the photo-regulation of their ATPase activities. Therefore, it is expected that Ras may be also regulated using photochromic molecules. In this study, basic experiments was performed in order to photoregulate its GTPase activity. A new fluorescent GTP analogue, NBD-GTP was synthesized in order to monitor the GTPase Ras. NBD-GTP showed initial burst enhancement of the fluorescent intensity by 120% when GDP free Ras was added and subsequently slow increase of fluorescence. The results may suggest that the initial burst of fluorescence reflects the binding of GTP to the free GTPase site of Ras, and the slow phase corresponds to the formation of Ras-GDP-Pi state. Quantitative analysis using DTNB cystein residue on the surface of Ras revealed that one of the intrinsic three cystein locate on the surface. However, SH group reactive photochromic molecule, 4-phenylazophenyl maleimide (PAM) was