coupled with FACS technology were employed to evaluate the extent of association and fusion of giant vesicles. Two fluorescent markers encapsulated in different vesicle populations were used as association markers; when these vesicles associate, the two independent markers should be observed simultaneously in a single detection event. The quenched fluorescent marker and the dequencher, which were also encapsulated in respective vesicle populations, were used as the fusion marker. When intermixing of the internal aqueous solutions occurs, the quenched marker is liberated by the dequencher and emits the third fluorescent signal. The population analysis of 100,000 giant vesicles revealed the stoichiometry of association and fusion of giant vesicles induced by exogenous addition of oppositely charged amphiphiles. Although a substantial fraction of vesicles associated together by adding small amount of the charged amphiphiles (5 % mole fraction to POPC), larger amount (25 %) was necessary to induce vesicle fusion. The present methodology also reveal a size dependency of association and fusion of giant vesicles, i.e., we show that giant vesicles of larger size tend to associate and fuse at higher frequency.

2P251 抗菌ペプチド・マガイニン 2 が脂質膜中に誘起するポア形成のキネティクスパスウェイ Kinetic Pathway of Antimicrobial Peptide Magainin 2-Induced Pore Formation in Lipid Membranes

Yukihiro Tamba, Hirotaka Ariyama, Victor Levadny, Jahangir Md. Alam, Masahito Yamazaki, (Shizuoka University)

The pore formation in lipid membranes induced by magainin 2 is considered to be the main cause for its bactericidal activity. To reveal the mechanism of the pore formation, it is important to elucidate the kinetic pathway of magainin 2-induced pore formation in lipid membranes. In this report, to examine the change in pore size over time during pore formation which can monitor its kinetic pathway, we investigated the rate of the leakage of various sized fluorescent probes through the magainin 2-induced pores in single giant unilamellar vesicles (GUVs) of 50% dioleoylphosphatidylglycerol(DOPG)/50%dioleoylphosphatidylcholine (DOPC) membrane using the single GUV method [1].

Magainin 2-induced leakage of Texas-Red dextran 10,000, Texas-Red dextran 3,000, and Alexa-Fluor trypsin inhibitor occurred in two stages; a transient rapid leakage in the initial stage followed by a stage of slow leakage. In contrast, magainin 2 induced a transient, but very small (10-20%) leakage of fluorescent probes of a larger size such as Texas-Red dextran 40,000 and FITC-BSA. These results indicate that magainin 2 molecules initially induce a large, transient pore in lipid membranes following which the radius of the pore decreases to a stable smaller size. We estimated the radius of these pores at the initial and the final stages of the leakage, which increased with magainin 2 concentration. Based on these data, we propose a hypothesis on the mechanism of magainin 2-induced pore formation.

1. Adv. Planar Lipid Bilayers and Liposomes, 7, 121-142, 2008

2P252 糖脂質分子膜の分子動力学シミュレーション Molecular dynamics study of gangliosides embedded in lipid membranes

Daisuke Takaiwa^{1,2}, Shinoda Wataru^{1,2}, Masuhiro Mikami^{1,2}, (¹Nanosystem Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Ibaraki, Japan: ²CREST-JST, 4-1-8 Honcho, Kawaguchi, Saitama 332-0012, Japan)

Gangliosides play an important role in molecular recognition. It is known that a finite amount of gangliosides is needed to effectively recognize biomolecules such as proteins, which suggests the importance of conformational change due to the interaction among carbohydrate headgroup chains [1] [2]. In this study, a series of molecular dynamics simulations of fully hydrated DPPC bilayers containing GM1 gangliosides at several different concentrations has been carried out to investigate the effect of GM1 to the structure and dynamics of membrane. Conformational changes of the oligosaccharide chain of GM1 gangliosides were found to be minor during the simulation time (production run time = 100 ns), which suggests the stiffness of the oligosaccharide chains. Increase in the deuterium order parameter of DPPC hydrocarbon chains is observed by adding the GM1, while the lateral diffusion coefficient of DPPC molecules is decreased. These observations suggest that GM1 increases the membrane rigidity and stability.

2P253 光照射による薬物放出制御システムの開発 Development of drug release system controlled by light irradiation

mariko hosoda¹, shin mizukami^{1,2}, kazuya kikuchi^{1,2}, (¹Graduate school of Engineering, Osaka Univ.: ²Immunology Frontier Research Center, Osaka Univ.)

Caged compounds are photoactivatable probes that are biologically or functionally inert before the uncaging. Utilization of caging/photouncaging techniques enables the spatiotemporal regulation of various biomolecules. Furthermore, these techniques are essentially similar to the controlled drug release systems investigated from the clinical perspectives.

We report a novel photocontrolled release system that can be applicable to photoactivation of various biomolecules. A photoresponsive drug carrier was used two parts: a drug carrier and a photoresponsive opener. We selected liposomes as the drug carriers and antimicrobial peptides (AMPs) as the opener. Usually, AMPs kill the bacteria by disrupting their membranes. Therefore, we assumed that this membrane-damaging property can be applied to construct a photocontrolled drug release system.

We designed a protected AMP derivative Bhmcoc-TL. Bhmcoc-TL consists of a short antimicrobial peptide temporin L (TL) and photoremovable 6-bromo-7-hydroxycoumarin-4-ylmethyloxycarbonyl (Bhmcoc) group. We demonstrated the membrane-damaging property of TL was controlled by the modification of the Lys residue with Bhmcoc group using liposomes included 5(6)-carboxyfluorescein as an indicator of membrane damage. When the liposomes Bhmcoc-TL were irradiated in the presence of Bhmcoc-TL, liposomes released the inclusions and fluorescence intensity was increased. In summary, photocontrolled release was achieved by the appropriate adjustment of the Bhmcoc-TL concentration and irradiation.

2P254 Melittin structure that is taken place during liposomal lysis.

Tomoyoshi Takahashi¹, Yasunori Yokoyama², Michio Homma¹, Kingo Takiguchi¹, (¹Division of Biological Science, Graduate School of Science, Nagoya University: ²Department of Computational Science and Engineering, Graduate School of Engineering, Nagoya University)

Melittin, major component of honey bee venom, is an amphipathic 26 amino acid peptide. For its simple structure and strong hemolytic toxicity, and solubilization, pore formation, membrane fusion that are induced in artificial membrane vesicles by the peptide, in numerous studies melittin has been used as a model peptide. Basis of the activities is membrane binding resulted from both electrostatic and hydrophobic interactions between the peptide and phospholipids, and subsequent assembly in membrane through neutralization of their charges. But the detail of mechanism is still unclear.

In our previous study, by addition of melittin to liposome made of 50% PC and 50% PG, we have found "phased shrinkage" of liposome we named, that is forming small particles on membrane surface and rapid shrinking. Increasing of brightness was observed in the liposomes that have shown the phased shrinkage. Such liposomes are named as "condensed liposome".

In this report, liposomal behavior under interaction with melittin was monitored using dark-field microscopy and compared with the results obtained by circular dichroism. We prepared liposomes composed of PC and PG at several molar ratios. When PC only or 10% PG liposome was used, membrane fluctuated. In this case, it is expected that melittin form random coil. In 30% and 50% PG, we observed solubilization, formation of the condensed liposome, increasing membrane surface area. The respectively estimated secondary structures of melittin were random coil, β -sheet like structure, and α -helix.

2P255 コレラ菌のアミノ酸走化性に関与する新規受容体の同定 Identification of novel receptors mediating taxis toward amino acids in *Vibrio cholerae*

Kimiko Yamamoto^{1,2}, Noriaki Sagoshi¹, So-ichiro Nishiyama^{3,4}, Ikuro Kawagishi^{1,3}, (¹Dept. Frontier Biosci., Grad. Sci. Eng., Hosei Univ.: ²Natl. Inst. Agro-Environ. Sci.: ³Dept. Frontier Biosci., Fac. Biosci., Appl. Chem., Hosei Univ.: ⁴Res. Cen. Micro-Nano Tech., Hosei Univ.)

Vibrio cholerae (causative agent of cholera) shows chemotaxis, which has been implicated in pathogenicity and survival in various environments. A classical biotype strain of *V. cholerae* O395N1 has 44 chemoreceptor homologs, which we name methyl-accepting chemotaxis protein-like proteins (MLPs). We previously found that Mlp24 and Mlp37 are major chemoreceptors mediating taxis toward amino acids in *V. cholerae*. However, the mutant lacking both *mlp24* and *mlp37* retains significant levels of response to some amino acids, suggesting the presence of yet other amino acid chemoreceptors. To identify such receptors, we examined chemotactic ability of the double mutant (Δ *mlp24* Δ *mlp37*) overproducing each of candidate MLPs by capillary assays. Cells overproducing Mlp2 or Mlp3 showed enhanced chemotactic responses toward L-serine and several other amino acids. Bioinformatic analyses predict that both Mlp2 and Mlp3 have shorter periplasmic regions than Mlp24 and Mlp37, each with two PAS-like domains in tandem, at least one of which is involved in amino acid binding. We are currently examining whether Mlp2 and Mlp3 actually mediate amino acid taxis.