measurement. β -LG is predominantly composed of β -barrel structure in the native state, and possesses an intermediate with non-native α -helical structure. To detect β -LG at the single-molecule level, we labeled fluorescence dye (Alexa 532) to the 51th cysteine of the E51C mutant. We checked that the fluorescence intensity of the labeled protein changes among the denatured state, the non-native intermediate and the native state. We observed the translational diffusion and the conformational changes of the labeled E51C by using the capillary cell system. We could roughly estimate the diffusion coefficient of the protein by tracking the centroid of the fluorescence spots. To improve the accuracy in the estimation, we are currently constructing a new detection system possessing a better positional accuracy for single molecule measurements.

3P075 Pressure dependence of protein conformations studied by generalized simulated tempering molecular dynamics simulations

Yoshiharu Mori, Yuko Okamoto, (Department of Physics, Nagoya University)

Structural change of proteins at high pressure has been studied experimentally and theoretically, which is known as pressure-induced denaturation of proteins. Understanding the molecular mechanism of the structural change at high pressure is important. Molecular simulations can be used to investigate such a mechanism.

We have recently developed a generalized-ensemble algorithm which is referred to as the isobaric-isothermal simulated tempering (PTST), which is an example of the generalized simulated tempering method. A PTST simulation performs a random walk in the two-dimensional pressure-temperature space, which allows one to obtain pressure dependence of molecular systems accurately and efficiently.

In the present study, we performed PTST molecular dynamics simulations of the systems of some proteins in explicit water molecules. We calculated several quantities (for example, the potential energy and volume of the systems and the structures of the proteins) at low and high pressure values and compared them.

We found that the potential energy and volume of the systems were widely distributed in the simulations as the results of a random walk in temperature and pressure. Therefore, the PTST simulations were successfully performed. We also found that the water molecules near the proteins played an important role in the structural change of the proteins at high pressure. We will discuss the protein structures and distributions of the water molecules near the proteins at the low and high pressure values.

3P076 Effect of Temperature on Pressure-Dissociation of Lysozyme 3E1250 Amyloid-Like Fibrils

Ryohei Kono^{1,3}, Akihito Kuroki², Yuta Kobayashi², Hideki Tachibana^{2,3}, (¹Sec Dept Int Med, Wakayama Med Univ; ²Dept Biotech Sci, Sch Biol-Oriented Sci Tech, Kinki Univ; ³High Pressure Prot Res Center, Inst Adv Tech, Kinki Univ)

Hen lysozyme disulfide-deficient variant OSS is intrinsically unfolded and forms amyloid-like fibrils, which are facily dissociated with high hydrostatic pressure. Analyses of the pressure-dissociation kinetics of OSS fibrils have shown a negative partial volume change on activation as well as a negative activation compressibility, indicating that the fibrillar state is rich in voids, which are partially hydrated in the transition state. Here, the effect of temperature on the pressure-dissociation of OSS fibril was investigated with high-pressure fluorescence measurements. The fibrils formed at 25 deg.C with an incubation period of 24 hr were subjected to high pressure (50 to 350 MP) at various temperatures (10 to 40 deg.C). The dissociation rate generally increased with increasing temperature. The absolute value of negative activation volume at 10 deg.C (at 0.1 MPa) was about half as much as that at 25 or 40 deg.C. The absolute value of negative activation compressibility at low temperature was also reduced to about half compared with those observed at high temperatures. These observations are consistent with the view that the fibrillar state has higher amount of voids and/or lower amount of hydrated water with the increase in temperature while the transition state does so to a less extent.

3P077 Refinement of Modeled Protein Structure by Replica Exchange Molecular Dynamics and Hydration Entropy based Scoring Function

Shiqiao Du¹, Yuichi Harano², Minoru Sakurai¹, Masahiro Kinoshita³, (¹Center for Biological Resource and Informatics, Tokyo Inst. Tech.; ²Institute for Protein Research, Osaka Univ.; ³Institute of Advanced Energy, Kyoto Univ.)

Modeling the native structure of a protein from its sequence information became more and more relevant to most issues of molecular biology. Anfinsen's Dogma tells us a useful strategy for this purpose: we can model the native structure as the global minimum conformation of the potential energy function in the same manner of optimization problem. The essence of the optimization

is mainly composed of two parts; sampling algorithm and scoring function. While information-based comparative modeling (CM) and fragment assembly (FA) are powerful sampling methods for rough modeling, physics-based molecular dynamics (MD) suites for refining the model structures at a fine resolution. Our scoring function "Fsolve" sheds light on the shape of a protein and how it affects water. It consists of hydration entropy (HE) and dehydration penalty (DP). HE can be calculated with morphological thermodynamics and angle-dependent integral equation theory. We have reported that Fsolve can discriminate native structure from those rough models with RMSD > 3.0 to native. In this study, we generated model structures by CM or FA and refined them by Replica Exchange MD. These models have quite good quality (RMSD < 3.0). We will report if Fsolve can select native structure from those decoys.

3P078 プリオンタンパク質への結合様式による抗プリオン化合物の分類と作用機構の解明 3E1305 Classification of anti-prion compounds based on the binding properties of prion proteins

Yuji Kamatari, Yosuke Hayano, Kei-ichi Yamaguchi, Junji Hosokawa-Muto, Kazuo Kuwata, (Center for Emerging Infectious Diseases, Gifu University)

To date, a variety of anti-prion compounds have been reported that are effective in *ex vivo* and *in vivo* experiments. However, the molecular mechanisms of most of these compounds are unknown. To understand those mechanisms, we have classified representative anti-prion compounds into several classes according to their binding properties for PrP^C. Surface plasmon resonance and NMR were used to determine the binding affinities and the binding sites of the compounds, respectively. Generally, compounds were classified into one of three classes. Compounds in the first class, including GN8 and GJP49, bound to the native structure and acted as "medicinal chaperones" to stabilize the native conformation and prohibit its pathogenic conversion. The second class of compounds, including Congo red, directly bound to PrP^C and promoted aggregation of the protein, which prevents further conformational change of prion. The third class of compounds, including pentosan polysulfate, did not bind to PrP^C but may interact with PrP^{Sc} or other relevant proteins. The proposed categorization of diverse anti-prion compounds would be useful for understanding the mechanisms of anti-prion compounds as well as facilitating further anti-prion drug discovery.

3P079 構造変化を伴うタンパク質 2 量体形成:結合・解離速度と初期構造依存性 Molecular Dynamics Simulation of Protein Dimerization: Association/Dissociation Rate and Hysteresis

Fumiko Takagi¹, Macoto Kikuchi², Nobuyasu Ito¹, (¹Department of Applied Physics, University of Tokyo; ²CMC, Osaka University)

Proteins in living cells often form protein complexes, and its conformation is often different in monomer and in a complex. They are in very crowded environment, and it is of interest that the effect of the confinement on protein complexes formation. We are studying protein homo-dimer formation accompanied with conformational change in a spherical cage using molecular dynamics simulation. We use a coarse-grained protein to perform large time scale simulation enough for protein binding or folding, and dual-Go model to describe the spontaneous structural change. In the past annual meeting, we reported the confinement effect on free energy profiles. Here, we investigated the confinement effect on kinetics, the cage size dependence of association/dissociation rate of dimer and structural change rate of monomers. We will also report that the initial structure dependence of the dimerization rate.

3P080 溶液添加剤によるタンパク質の熱失活抑制機構 Molecular mechanism of the solution additives on thermal inactivation of proteins

Tomita Shunsuke, Shiraki Kentaro, (Graduate School of Pure and Applied Sciences University of Tsukuba)

Protein inactivation poses a serious problem in various fields. One of the versatile approaches for suppression of protein inactivation is the addition of additives in the protein solution, though the mechanism remains elusive. To solve this issue, we investigated thermal inactivation and non-enzymatic chemical modification of RNase A and lysozyme in the presence of various solution additives, such as amino acids, amino acid derivatives, inorganic salts, and polyamines. The clear data were obtained that the chemical modifications on RNase A and lysozyme was suppressed in the presence of solution additives that prevented thermal inactivation of these enzymes. More importantly, the rate constants of heat-induced inactivation of the enzymes decreased with