WT [pXO1⁺pXO2⁺], pXO2-depleted [pXO1⁺], pXO1-depleted [pXO2⁺], and both plasmids-depleted *B.anthraxis*, were added to several cultured cells. As a result, we observed morphology change of HeLa and vero cell only treated by the culture supernatants of WT and pXO1⁺ strains. This morphology change of cultured cell might be attributed to pXO1. Of especially note is that fibrous form of vero cell was observed just for 5 hours. Now we tried to confirm the possibility of various cellular responses including apoptosis and actin-polymerization.

3P303 ミトコンドリア内膜タンパク質予測法の開発 Computational prediction of mitochondrial inner membrane proteins

Tsuji Toshiyuki, Nakai Kenta, (Human Genome Center, Institute of Medical Science, University of Tokyo)

Proteins are transferred to the correct subcellular location to properly perform their functions. Prediction of the protein subcellular localization is very important for exploring the function of proteins in the cell. Although, many subcellular localization prediction methods have been developed, these systems ignore the mitochondrial inner membrane proteins without N-terminal targeting signal. It is known that the inner membrane proteins have the internal transit noncleavable signal. However the detection of the internal signals is very difficult. In this study, we analyzed the propensities of amino acids and various physicochemical properties of five types of protein, which is localized at the mitochondrial inner membrane, the mitochondrial matrix, cytoplasm, the cell membrane or the extracellular space. These properties were used for developing a prediction method based on support vector machine (SVM). We analyzed the amino acid propensities around the cleavage site of the mitochondria targeting signal at N-terminal of mitochondrial matrix protein and made the position specific score matrix (PSSM). Using the PSSM, we searched the potential cleavage site of mitochondria inner membrane proteins. The distributions of physicochemical properties around the potential cleavage site were used as the input vector of SVM. The performance of our predictor outperforms previous predictors in the discrimination between the mitochondrial inner membrane proteins and plasma membrane proteins.

3P304 トランスクリプトーム形成の熱力学モデルと、マイクロアレイ分析への応用 A thermodynamic model of transcriptome formation and its application to microarray data analysis

Tomokazu Konishi, (Akita Prefectural University)

The genome supplies information on both the quality and quantity of the transcriptome. However, as it remains unknown how a cell determines transcript levels from the genome sequences, despite comprehensive knowledge of the cellular components involved, the quantity information held by the genome cannot as yet be derived from nucleotide sequences. The model presented here explains on a thermodynamic basis how the components decode the genome to form and maintain the transcriptome. The model describes the level of a transcript as a pseudo-equilibrium between velocities of synthesis and degradation, both of which are controlled by sequence-specific interactions between protein factors and nucleic acids. Each of the transcript levels can be described by a single equation expressing a function of the activity concentrations of the protein factors. Quantitative information in the genome can thus be transformed into constants determined from the nucleotide sequences. Using this model, the transcriptome can be traced back to the protein factors and the state of chromosome packaging. The total description of transcript levels allows the model to be verified through comparison of derived hypotheses with comprehensive measurements of the transcriptome. The hypotheses thus derived in the present study are well supported by microarray data, confirming the appropriateness of the model. Additionally, a characteristics of data distribution can be deduced by the model. This has derived a basis of parametric framework for data analysis.

3P305 Development of High Performance Predictor of Bacterial and Mitochondrial Outer Membrane β -barrel Proteins.

Kenichiro Imai^{1,2}, Naoya Fujita¹, Paul Horton¹, (¹Computational Biology Research Center, AIST; ²Research Fellow of the Japan Society for the Promotion of Science)

The outer membrane of mitochondria and chloroplasts as well as Gram-negative bacteria, their evolutionary cousins, harbor β -barrel outer membrane proteins (BOMPs). Bacterial BOMPs perform critical functions such as channels for ion transport, nutrient intake and enzymes. In pathogenic Gram-negative bacteria, BOMPs are often involved in their acquisition of drug resistance. Mitochondrial BOMPs also perform the import functions as exemplified by the formation of the pore of the translocase of the outer membrane, the component of the sorting and assembly machinery complex and the con-

tribution to membrane permeability in mitochondrial induced apoptosis. The accurate prediction of BOMPs is important for facilitating the development of novel drugs. In this study, we have developed prediction methods for bacterial and mitochondrial BOMPs which based on support vector machine method using not only amino acid composition but also features of C-terminal membrane integration signal, N-terminal signal peptide, periodic amphiphilicity and position weight matrices for the transmembrane β -strands. Measuring performance on a dataset of 1434 bacterial proteins with 30% identity cutoff (60 BOMP / 1374 Non-BOMP), we obtained a precision of 97.9%, recall of 71.7% and Mathew's correlation coefficient (MCC) of 0.83. For mitochondrial BOMPs, we obtained a precision of 91.4%, recall of 81.0% and MCC of 0.86 on a dataset of 2745 proteins with 30% identity cutoff (105 mitochondrial BOMP / 2640 Non-BOMP).

3P306 多塩基置換を考慮したコドン置換モデルにより評価したアミノ酸置換における選択的制約 Selective Constraints on Amino Acids Estimated by a Mechanistic Codon Substitution Model with Multiple Nucleotide Changes

Sanzo Miyazawa, (Graduate School of Engineering, Gunma University)

We develop a new codon-based model, which consists of mutations at the nucleotide level and selection at the amino acid level via a genetic code, and estimate selective constraints on amino acids from available empirical substitution matrices. The mutational process of individual codons is modeled as a general reversible Markov process, and codon replacements that require multiple nucleotide changes are assumed to occur with the same order of time as single nucleotide changes do. The present model with substitution rates that are assumed to obey a Γ distribution is fitted to each 1-PAM matrix of empirical amino acid substitution matrices (JTT, WAG, and LG) and an empirical codon substitution matrix (KHG). Maximum likelihood (ML) estimators of selective constraints on amino acids are calculated together with other parameters. Akaike information criterion (AIC) values indicate that the assumption of multiple nucleotide changes significantly better fits the model to the empirical substitution matrices. Also, one of interesting results is that the ML estimators of transition to transversion bias obtained from these empirical matrices are not so large as previously estimated. Selective constraints on amino acids are characteristic of each type of amino acid rather than each species. Thus, the present codon-based model with the ML estimators for the selective constraints and with adjustable mutation rates of nucleotides would be useful as a simple substitution model in the ML inference of molecular phylogenetic trees.

3P307 Detecting structural similarity of ligand interactions in the lipid metabolic system of NRs, lipid binding proteins, and metabolic enzymes

Clara Shionyu-Mitsuyama¹, Tsuyoshi Waku¹, Takuma Shiraki², Takuji Oyama¹, Tsuyoshi Shirai³, Kosuke Morikawa¹, (¹Institute for Protein Research, Osaka University; ²Department of Biochemistry, Tohoku University Graduate School of Medicine; ³Department of Bioscience, Nagahama Institute of Bio-science and Technology)

Nuclear receptors (NRs), intracellular lipid binding proteins (iLBPs), and metabolic enzymes are responsible for optimal metabolic homeostasis in higher organisms. The specific functional relationships were detected among the subfamilies of these proteins. In this study, the NR - iLBP - enzyme system, in which the interactions are mostly mediated by ligand molecules, was examined in terms of their ligand-binding structures to detect the similarity in the interactions between functionally related subfamilies. The complex structures from PDB were dissected into single amino acid motifs for ligand fragment binding, and the presence and evolutionary origin of the motifs were compared among the families. As a result, pairs of the functionally related NR and enzyme were found to share more motifs than expected, in agreement with the fact that the two families compete for the same ligand to coordinate metabolome-transcriptome system. This finding implies the co-evolution of the indirectly interacting protein system.

3P308 シグナル伝達におけるゆらぎの伝播が引き起こすシグナルの増幅と絶滅 Signal amplification and extinction caused by propagation of fluctuation in a signal transduction pathway

Masashi Kajita¹, Makio Tokunaga^{1,2}, (¹Graduate School of Bioscience & Biotechnology, Tokyo Institute of Technology; ²Research Center for Allergy & Immunology, RIKEN)

Dynamic features in signaling pathways are of particular interest to understanding the dynamic and variety functions of cells. Systems biology has contributed to clarifying properties of combined motifs, such as the MAP kinase