experiments involving ^{15}N spin relaxation parameters (R_1 , R_2 , NOE, R_2 relaxation dispersion), chemical shifts (^1H , ^{15}N , ^{13}C), hydration (Cleanex-NMR) and residual dipolar coupling demonstrates structure, dynamics and a population of N_2 state of the ubiquitin mutants.

1H1512 ヒューマン p 100 の C 末端ドメインと黄色ブドウ球菌由来核酸分解酵素間における構造エレメントの比較

Comparison of structural elements between Staphylococcal nuclease and the C-terminal domain of Human p100

Shou Ueyanagi, Hironari Kamikubo, Yoko Ogawa, Chikako Komeda, Yoichi Yamazaki, Mariko Yamaguchi, Mikio Kataoka (*Graduate School of Material Science, Nara Institute of Science and Technology*)

We revealed by systematic alanine insertion analysis that staphylococcal nuclease (SNase) is composed of several structure elements. Structure element is defined as the region where alanine insertion cannot be accepted. We consider that structure elements are building blocks a protein. We expect that structure elements appeared to be at the similar positions among structurally homologous proteins. To confirm this idea, the alanine insertion analysis is applied to a SND domain of a human transcriptional regulator protein, p100, whose tertiary structure is almost identical to that of SNase despite of low sequence identity (15%). Previously, we performed the alanine insertion analysis upon the $\alpha 1$ helix of SND, and found that a structure element is conserved in both $\alpha 1$ -helices of SNase and SND. In this work, to investigate the conservation of the other structure elements, the alanine insertion analysis was applied to the β -strand of SND adjacent to the $\alpha 1$ -helix. The corresponding β -strand of SNase involves structure elements. CD spectra of the insertion mutants of SND indicated that the alanine insertion into the regions between the 799th and the 807th residues and between the 815th and the 817th leads to the loss of the structure, while the insertion mutants in the other parts exhibit a native-like CD spectrum. The regions influenced by the insertion mutation are corresponding positions to the structure elements of SNase. This coincidence supports our idea that the structure elements are common building blocks of a protein.

1H1524 スタフィロコッカスクレアーゼの野生型と非天然構造を有する変異体に見られる局所構造の相違

Difference in local structure between staphylococcal nuclease wild type and its non-native mutant studied by tryptophan triplet quenching

Mariko Yamaguchi, Yoichi Yamazaki, Hironari Kamikubo, Mikio Kataoka (*Nara Institute of Science and Technology*)

Characterization of the denatured structure in physiological conditions is necessary for the deep understanding of the stabilization mechanism of native structure. Mutagenetic studies of staphylococcal nuclease (SNase) have found that 139A140, which is a mutant with alanine (Ala) inserted between the 139th and the 140th residues, does not take native structure owing to the lack of proper hydrophobic interactions. We compared the local structure of 139A140 with that of wild type (WT) in order to study how the hydrophobic interactions are modified by the Ala insertion. The local structure of WT and 139A140 were monitored by the quenching of triplet state (TS) of Trp140 by cysteine (Cys), which was substituted for Lys134. Signals from the TS showed two exponential decays for both WT and 139A140. The lifetime of the fast component was 1-2 μs and that of the slow component was 20-40 μs . The fast component is associated with the denatured states, in which the TS is quenched by Cys. The slow component, on the other hand, is associated with the native state, in which the TS is not quenched by Cys. Guanidine hydrochloride (GdnHCl) concentration dependence was not observed in the lifetimes of both the fast and the slow components for 139A140, while for WT the lifetime of the slow component shortened with increasing GdnHCl. The difference in the local structure around the Trp between WT and 139A140 is discussed from the dependence of the TS quenching on GdnHCl concentration and Trp-Cys distance.

1H1536 網羅的アラニン挿入変異解析法によるジヒドロ葉酸還元酵素の機能エレメントの分類

Classification of function element of dihydrofolate reductase by the systematic alanine insertion

Rumi Shiba¹, Hironari Kamikubo¹, Mika Umeyama¹, Sayaka Tsukasa¹, Yoichi Yamazaki¹, Mariko Yamaguchi¹, Masahiro Iwakura², Mikio Kataoka¹ (¹*Graduate School of Materials Science, Nara Institute of Science and Technology*, ²*Protein Design Research Group, Institute for Biological Resources and Functions, National Institute of Advanced Industrial Science and Technology*)

Decoding sequence information is equivalent to elucidating the design principles of proteins. For this purpose, we successfully applied systematic alanine (Ala) insertion analysis to dihydrofolate reductase (DHFR). We revealed that DHFR is composed of structure elements, function elements and linkers connecting these elements. Although some structure elements overlap with function elements, other structure elements are not function elements, namely, the alanine insertion mutants into the latter region lost their tertiary structure but possess their functions. To elucidate the mechanism that unstructured mutants can exhibit their activity, solution structure and enzymatic activity of the insertion mutants of the region from the 71st to the 86th residue were examined quantitatively. This region contains a structure element. The mutants with an Ala insertion into the structure element take non-native structures, while the other mutants take native structure. Fluorescence titration experiments were performed with the insertion mutants into the structure element in order to reveal the binding ability to the substrate of DHF. The fluorescence titration revealed all of the insertion mutants possess their binding ability of DHF. In addition, CD spectrum changes upon binding of DHF to the mutants are almost identical to that of WT, although the final CD spectrum is not necessarily close to native structure. These results suggest that DHF binding to the mutants of the structure element induces native-like substrate binding site.

1H1548 N 末端配列の相違による α ラクトアルブミン不安定化の構造的知見 Structural insights into the stability perturbations by N-terminal variations in human and goat α -lactalbumin

Koki Makabe^{1,2}, Takashi Nakamura¹, Kunihiro Kuwajima^{1,2} (¹*Okazaki Institute for Integrative Bioscience*, ²*Graduate University for Advanced Studies (Sokendai)*)

α -Lactalbumin is a major component of milk protein and it has been used as a model protein for folding study because it forms molten globule state under mild denaturation conditions at equilibrium, which is identical with a kinetic folding intermediate. It has been well known that the N-terminal portion of α -lactalbumin significantly contributes to the stability. The addition of an extra methionine residue at the N terminus for recombinant expression in bacteria results in a significant stability loss. This reduction of the stability hampers the mutational study of α -lactalbumin, such as the phi-value analysis in folding study, because the recombinantly expressed α -lactalbumin is largely destabilized with a single mutation and difficult to determine its free energy precisely. We have overcome this stability loss by mutating the first residue (Lys and Glu in human and goat α -lactalbumin, respectively) to methionine. Here, we show that the crystal structures of the N-terminal mutants of human and goat α -lactalbumin. The crystal structures of the mutants suggests the importance of a hydrogen bond between the N-terminal amide nitrogen atom and an O_γ atom of Thr38. Combining the structural information with the 2D-NMR results that show the chemical shifts perturbations caused by the N-terminal mutations, we will further discuss the effects of the N-terminal residue in their stability.

1H1612 Conformational fluctuation of c-Myb DNA-binding domain analyzed by temperature and pressure perturbation experiments

Satomi Inaba¹, Harumi Fukada², Sunilkumar N. Puthapurackal^{3,4}, Akihiro Maeno^{3,4}, Hiroshi Matsuo⁴, Hisayuki Morii⁵, Kazuyuki Akasaka⁴, Masayuki Oda¹ (¹*Grad. Sch. of Life and Environ. Sci., Kyoto Pref. Univ.*, ²*Grad. Sch. of Life and Environ. Sci., Osaka Pref. Univ.*, ³*Grad. Sch. of Biol. and Sci., Kinki Univ.*, ⁴*High-pressure Protein Res. Center, Kinki Univ.*, ⁵*Biomed. Res. Inst., Adv. Indust. Sci. and Tech.*)

The transcription factor protein, c-Myb, has three functional domains for DNA-binding, transcriptional activation, and negative regulation from amino terminus. The DNA-binding domain consists of three imperfect tandem repeats of 51 or 52 amino acid residues (designated R1, R2 and R3), and the last two repeats, R2 and R3, are sufficient for DNA-binding. In contrast to R3, R2 has a relatively large cavity in hydrophobic core, and Val103 and Val107 at first helix of R2 are key residues for forming its cavity. The previous studies have shown that the cavity-filling mutation in R2 results in increased thermal stability, but loss of specific DNA-binding. In this study, we focused on Val103, and analyzed its mutational effects on thermal and pressure stabilities by circular dichroism, differential scanning calorimetry, and high-pressure fluorescence measurements to determine the conformational fluctuation of c-Myb R2R3. The results showed that V103L mutant was more endurable to either temperature or pressure than wild-type. This should be correlated with the partial molar volume change determined in the high-pressure experiments. In the presentation, the