Koichiro M. Hirose¹, **Kenta J. Yoshida¹**, Kenichi G. N. Suzuki^{1,2}, Takahiro K. Fujiwara¹, Ankita Chadda¹, Akihiro Kusumi¹ (¹*Institute for Integrated Cell-Material Sciences and Institute for Frontier Medical Sciences, Kyoto University*, ²*PRESTO-JST*)

The signal of receptor engagement is propagated along the 2D plasma membrane, which would have to be regulated by both signal amplification and spatial spread by molecular diffusion. We examined these processes using IgE receptor (FcεRI) and its downstream signaling molecules as an important paradigm. About 30 s after FcεRI crosslinking by the multivalent antigen, FcεRI phosphorylation by Lyn was biochemically detected, but Lyn concentration at FcεRI clusters has not been detected by microscopy. Using simultaneous single-molecule imaging of Lyn tagged with Halo protein and FcεRI, we found that, within 30 s after multivalent antigen stimulation, single molecules of Lyn were frequently and transiently recruited to FcεRI clusters, with an exponential residency time of ≈ 120 ms, showing that Lyn is dynamically recruited but not accumulated. Lyn's downstream kinase Syk is then recruited to FcεRI clusters, and immobilized there for at least 4 s (the limit determined by photobleaching). Its downstream adapter transmembrane protein LAT is also recruited to FcεRI clusters one molecule after another, but transiently (≈ 120 ms), suggesting that the signal amplification in this pathway is performed via the LAT phosphorylation process by Syk. LAT diffusion is as fast as that of phospholipid, suggesting that LAT is responsible for spatially spreading the signal along the plasma membrane. These data are at total variance with the traditional view in which these molecules form large complexes centered around the FcεRI clusters.

2K1536 Src-family キナーゼ活性を制御するラフト機構：足場タンパク質 Cbp のリクルートの1分子検出

Raft mechanism for regulating Src-family kinases: detection of single-molecule recruitment of a scaffolding transmembrane protein Cbp

Yuki Shirai¹, Kenichi Suzuki^{1,2}, Hideaki Yoshimura¹, Koichiro Hirose¹, Takahiro Fujiwara¹, Akihiro Kusumi¹ (¹*Institute for Integrated Cell-Material Sciences and Institute for Frontier Medical Sciences, Kyoto University*, ²*PRESTO-JST, Kyoto University*)

Our group previously showed that a Src-family kinase protein, Lyn, is recruited to the rafts induced by the clustering of GPI-anchored receptor CD59 (Suzuki et al. 2007a, b; J. Cell Biol.). Simultaneous single-molecule tracking of Lyn and CD59-cluster rafts revealed that Lyn is recruited to CD59-cluster rafts very often, but transiently with a lifetime of ~ 0.2 s. Furthermore, our data suggested that Lyn activated at the raft induced the binding of the raft to the membrane skeletal actin filaments, leading to its Stimulation-induced Temporary Arrest of Lateral diffusion or STALL, at which PLCgamma is recruited, probably triggering IP3-calcium responses. This shows that the CD59-cluster rafts undergoing STALL is the key (although transient) site for relaying the extracellular CD59 signal to intracellular responses. One of the key molecules for the negative feedback of Lyn is the Csk-binding-protein (Cbp), a transmembrane scaffolding protein, which could bind to both Lyn and Csk. Here, we examined the possibility that Cbp is recruited to the CD59-cluster rafts to induce its STALL. Simultaneous single molecule tracking of Cbp and CD59-cluster rafts revealed the recruitment of Cbp (0.13-s lifetime). Furthermore, Cbp recruitment/residence event at the CD59 cluster occurs two-times more often during the STALL period (including the periods 0.27 s before and after the STALL) than the non-STALL period, suggesting that although Cbp is involved in inducing STALL, the recruitment of another unidentified molecule is likely required to induce STALL.

2K1548 接着分子によるイノシトールリン脂質代謝系自己組織化パターンの制御

Regulation of self-organization in chemotactic signaling system by an adhesion molecule, talin

Shinichi Yamazaki¹, Satomi Matsuoka^{1,2}, Masatsune Tsujioka¹, Hiroaki Takagi³, Masahiro Ueda^{1,2} (¹*Graduate school of Frontier Biosciences, Osaka University*, ²*Riken, QBiC*, ³*Nara Medical University, Department of Physics*)

Cellular motility is a basic function underlying important physiological phenomena such as embryogenesis, wound healing, and immune response. Moving cells show clear anterior-posterior polarity in which pseudopodia are frequently extended at the leading edge, while are suppressed at the lateral and trailing edge. In addition, adhesion to the substratum is enhanced at the anterior region of the cells, while is reduced at the posterior region. In order to migrate

effectively, intracellular molecular networks that undertake actin cytoskeleton, adhesion and signal transduction should work together in a coordinated fashion. Phosphatidylinositol (PtdIns) lipids are key signaling mediators for cellular migration and chemotaxis. We have previously reported that the signaling system is self-organized to generate traveling waves of PtdIns 3,4,5-trisphosphate and PTEN (PIP₃/PTEN wave) on the plasma membranes of Dictyostelium cells. Self-organized PIP₃/PTEN wave serves as a spontaneous signal for random cell movement in the absence of external guidance cues. Here we report that an adhesion molecule, talin is involved in proper self-organization of PIP₃/PTEN wave. An extraordinary PIP₃/PTEN wave was observed in the mutant cell lacking talin gene, suggesting an existence of regulatory pathway from adhesion system to PtdIns lipids signaling system. Possible coordination between actin cytoskeleton and adhesion system by self-organization in PtdIns lipids signaling system will be discussed.

2K1612 Effects of cyclophilin D on energy metabolism of cells

Hikaru Osaki, Takuma Masuguchi, Hitomi Nakazato, Yoshihiro Matsunomoto, Chisako Fujita, Yoshihiro Ohta (*Dept Life Sci & biotech, grad. sch. of engineering, Tokyo university of agriculture and technology*)

Cyclophilin D (cypD) is a peptidyl prolyl cis-trans isomerase (PPIase) localized in the mitochondria matrix. This protein is known to influence apoptosis, necrosis and autophagic cell death. These kinds of cell death are reported to depend on the energy state of the cell. The aim of this study is to examine the effects of CypD on the energy metabolism on cells. For this purpose, we established three kinds of C6 glioma cell lines, cells over expressing wild-type CypD(WT), cells with PPIase deficient mutants of CypD (R97A), control cells transfected with empty vector(V). In R97A mutants, CypD activity was not observed probably because of dominant negative effect. To examine the energy metabolism in these cells, we measured oxygen consumption rate, the amount of lactic acid released to the extracellular medium and intracellular ATP concentration. Concerning to oxygen consumption, we measured the effects of oligomycin. Oligomycin reduced the respiration rate in all three cells, the reduction rate in WT cells was smaller than those in the other cells. These results suggest that CypD may stimulate uncoupling of oxidative phosphorylation in mitochondria. Together with the results of lactic acid release and intracellular ATP concentration, we will discuss the effect of CypD on the energy metabolism of cell.

2K1624 マスト細胞の開口放出におけるアクティブゾーンタンパク質と細胞接着の関与

Involvement of active-zone molecules and cell adhesion in exocytosis of mast cells

Hidehiro Nomura, Satoshi Tadokoro, Masahiko Tanaka, **Naohide Hirashima** (*Grad. Sch. Pharmaceut. Sci., Naogyo City Univ.*)

The active zone is a specialized region of the presynaptic plasma membrane where the neurotransmitter release occurs by exocytosis. At synapses, pre- and post-synaptic cells make contact each other and the active-zone serves to enhance the efficiency of neurotransmitter release, allowing fast signal transmission from pre- to post-synaptic cells. Mast cells also release inflammatory mediators by exocytosis resulting in induction of allergic responses. In this study, we investigated the involvement of active-zone proteins and cell adhesion in exocytotic release from mast cells. We found the expression of active-zone proteins such as ELKS and liprin- α . Knockdown experiments revealed that these active-zone proteins regulated mast cell exocytosis positively. We also investigated the effects of cell adhesion on exocytosis using polystyrene beads. Contact of polystyrene beads with mast cells enhanced exocytotic release. To elucidate the mechanism we observed the intracellular localization of secretory granules, active-zone proteins and SNARE (soluble N-ethyl maleimide-sensitive factor attachment protein receptor) proteins. We found that not only secretory granules but also active-zone proteins and SNARE proteins were accumulated to the cell adhesion sites with beads. These results suggest that exocytotic release is facilitated by cell adhesion by recruiting molecules to the adhesion sites.

2K1636 走化性運動中の粘菌細胞におけるクラスリン依存性エンドサイトーシス

Clathrin-mediated endocytosis in chemotaxing Dictyostelium cells

Kunito Yoshida, Christien Merrifield (*MRC-LMB, Cambridge*)