we prepared domain deletion mutants of FlgT and analyzed their effects on the basal-body ring formation. The results suggest that FlgT-N contributes mainly to construct the H-ring structure, and FlgT-M may mediate the T-ring association on the LP-ring. FlgT-C is not essential, but stabilizes the H-ring structure. Based on these results, we will discuss the assembly mechanism of the basal-body rings and the stator units of the sodium-driven flagellar motor.

3N1334 菌体細胞膜に埋まったビブリオ菌のナトリウムイオン駆動型べん毛モーター構造の電子顕微鏡による可視化

Visualization of *Vibrio* Na⁺-driven flagellar motor structure embedded in cell membrane by electron microscopy

Masafumi Koike¹, Atsuhiko Narita¹, Michio Homma², Yuichiro Maeda¹ (¹Structural Biology Research Center, Nagoya Univ., ²Dept. Biological Sci., Grad. Sch. Sci., Nagoya Univ.)

The flagellum is one of the biggest architecture in the bacterial organ. At the base of each flagellum, the ion-driven rotary motor is embedded in the membrane. *Escherichia coli* and *Salmonella* have H⁺-driven motor, and *Vibrio alginolyticus* has Na⁺-driven type. The flagellar motor is composed of basal body, C ring and stator. Basal body is the rotor part of the motor. *Vibrio* basal body has unique ring structures important for the function named T and H ring. C ring composed of FliG, FliM and FliN, is mounted on the cytoplasmic side of the basal body, and essential for torque generation by interact with stator. FliG participates most directly in the torque generation among them. Since, however, C-ring structure has isolated and identified just only from *Salmonella*, the C-ring structure of Na⁺-driven flagellar motor is still unknown. Previously, we isolated FliG associate with the basal body from the Na⁺-driven polar flagellum of *V. alginolyticus*. However, FliM and FliN dissociated during the C-ring preparation. Recent fluorescent analyses show the turnover of FliM and FliN in *E. coli* cells. It suggests that C ring is the fragile structure in bacterial cells. And in spite of the continuous structural studies of flagellar motor, structures of the stator proteins mounted on the motor have not been observed by electron microscopy. To visualize C ring and stator with basal body, currently, we attempt to observe flagellar motor embedded in *Vibrio* cell membrane by electron microscopy.

3N1346 溶液 NMR を用いたビブリオ菌べん毛の固定子タンパク質 FliG の動的構造変化の解析

Structural changes investigated by solution NMR in *Vibrio* flagellar rotor protein FliG

Mizuki Gohara¹, Rei Abe-Yoshizumi⁴, Shiori Kobayashi¹, Yoshikazu Hattori^{2,3}, Chojiro Kojima^{2,3}, Michio Homma¹ (¹Division of Biological Science, Graduate School of Science, Nagoya University, ²Institute for Protein Research, Osaka University, ³Graduate School of Biological Sciences, Nara Institute of Science and Technology, ⁴Department of Frontier Materials, Nagoya Institute of Technology)

In the bacterial flagellar motor, torque is generated by the interactions between rotor and stator. In the rotor, a C ring component FliG is mainly participated in torque generation. Recently, the crystal structure of whole FliG in *Aquifex aeolicus* was published. FliG consists of three (N, M, and C) domains that contain several α -helices. The paper suggested that dynamic conformational changes occur during rotational switching and torque generation. To investigate structural changes in FliG derived from *Vibrio* Na⁺-driven motor, we performed solution NMR. First we overproduced and purified full-length FliG, and measured its ¹H-¹⁵N HSQC spectrum to search an optimal condition for the measurement. We found that in pH 6.5 signal peaks were separated better than pH 8.0. Next, we made N-terminally truncated FliG variants, G122-FliG (MC-FliG) and G214-FliG (C-FliG), and measured ¹H-¹⁵N HSQC in pH 6.5 using NMR at 800 MHz. From these constructs, we could get much better peaks, which are possible to assign. Furthermore, we carried out DSC (Differential Scanning Calorimetry) measurements to examine their thermodynamic properties. Results suggested that FliG protein has domain structures, and G122-FliG might form a multimeric structure in room temperature. Currently we are trying to assign the peaks of FliG, and conformational dynamics will be examined using the FliG deletion derivatives.

3N1358 べん毛フックタンパク質 FlgE 変異体におけるフックの長さ制御

Hook-length control in *flgE* mutants

Kaoru Uchida, Kohei Dono, Shin-ichi Aizawa (Prefectural University of Hiroshima)

In bacterial flagella, a flagellar protein FliK controls the length of the hook, which connects the filament to the flagellar basal body (BB) embedded into the membrane. The hook is composed of a single kind of hook protein (FlgE). A *flgE* mutant SJW2219 is temperature sensitive; cells produce normal flagella and actively swim when grown at 30C, whereas they became non-flagellated and thus non-motile at 37C. At 30C, most cells produced wild-type flagella. We found by electron microscopy that mutant cells grown at 37C produced intermediate structures of the flagellar hook-basal body (HBB) as many as wild type. About half of the population produced a few intact flagella/ cell, in which hook length was not different from that of the wild type. In addition, SDS-PAGE analysis showed that a considerable amount of mutant FlgE was secreted into the culture media. When we introduced modified *fliK* constructs into the mutant, the recombinant produced short and long hooks according to the size of FliK expressed from the constructs. Therefore, we conclude that FliK controls hook length in this *flgE* mutant, even though the polymerization rate of FlgE might be slower than that of wild type.

3N1422 べん毛モータのトルク発生器における帯電残基の性質

Characterization of the electrically-charged residues in the torque-generating unit of the flagellar motor

Satoshi Inaba, Shin-Ichi Aizawa (Prefectural University of Hiroshima)

The flagellar motor is composed with three kinds of proteins: FliG, FliM, and FliN, in *Salmonella enterica* serovar Typhimurium. It has been postulated that five conserved charged residues (K264, R281, D288, D289, R297) in the FliG C-terminal domain are directly involved in torque generation. We substituted these five residues with Thr individually and observed the behavior of the mutants. The D288T & D289T double-mutant strain was Mot⁻, while the rest were Mot⁺. To examine the functional role of the R281 residue that is the highly conserved one, we substituted this residue with all 20 amino acids and observed behavior of the constructs. Four substitutes (R281D, R281E, R281G, and R281P) were Mot⁻, while the rest could swim at various speeds on agar plates. Accordingly, their swimming speeds widely ranged from zero to wild-type speed: wild-type R > YQTSKCAH > V > WMFILN > DEGP = Mot⁻. As examined by the tethered cell method, the rotation speed of a single motor of the variants continuously dispersed in a wide range as follows; from wild-type speed R > T > S > Q > K > Y > C > A > W > V > (M, H, L, F, I) to zero speed (D, E, P, G), indicating that torque generated in the motor may fluctuate under different physiological conditions in the micro-environment of the site R281.

3N1434 Ascaris 精子をもちいたアメーバ運動装置の in vitro 再構成

Simultaneous reconstitution of MSP-based protrusion and retraction in the amoeboid sperm of Ascaris

Katsuya Shimabukuro^{1,2}, Naoki Noda³, Masasuke Yoshida^{1,4}, Murray Stewart⁵, Thomas Roberts¹ (¹JST ICORP Yoshida ATP synthesis regulation, ²Florida State University, ³MBL, ⁴Kyoto Sangyo University, ⁵MRC)

Crawling movement in eukaryotic cells requires coordination of the force for leading edge protrusion with that for cell body retraction. In *Ascaris* sperm, protrusion and retraction are powered by a motility system comprised of major sperm protein (MSP) in place of the actin apparatus typically associated with cell migration. By adjusting the concentration of cell-free sperm extracts, we reconstituted protrusion and retraction simultaneously in MSP filament meshworks, called fibers, that assemble behind plasma membrane-derived vesicles. As polymerization of filaments at the growing end of the fiber pushed the vesicle ahead (protrusion), filament rearrangement and disassembly pulled the rear of the fiber forward (retraction). Because nematode sperm do not use motor proteins to power retraction, we used these fibers to explore the mechanism of retraction by focusing on the cytoskeletal dynamics that occur during the transition from protrusion to retraction. Using a combination of fluorescent speckle microscopy, electron tomography, and advanced polarization microscopy, we found that although fibers progressively lost filaments during retraction the average length of the filaments did not change. The surviving filaments were able to move and reorient so that their packing density was conserved. To do so, fewer filaments packed into a smaller volume and this rearrangement appears to provide the driving force for retraction.

3N1446 ショウジョウバエ形態形成におけるアポトーシスの力学的寄与

Active Mechanical Role of Apoptosis in Tissue Dynamics during *Drosophila* Morphogenesis

Yusuke Toyama^{1,2,3} (¹Dep. Bio. Sci., Nat. Univ. Singapore, ²Machanobiology Inst., Singapore, ³Temasek Life Sciences Lab.)

Understanding tissue dynamics during embryo development requires knowledge of active mechanical forces exerted by cells and passive response by the neighboring cells. Apoptosis, or programmed cell death, is the process that remodels tissue by eliminating cells that are no longer biologically necessary. During the apoptotic process, the dying cell is squeezed out from the original tissue by mechanical forces produced within the cell and within the neighboring non-apoptotic cells. Those neighboring cells are also deformed during the cell elimination, indicating that the mechanical stress in the surrounding tissue is increased. Combining *in vivo* imaging, laser microsurgery, quantitative image analysis, and theoretical modeling, we deduced the tension of the tissue and found a correlation between cell death and mechanical force. In this presentation, we will present two discrete developmental phenomena during *Drosophila* morphogenesis as the examples of tissue dynamics where apoptosis could play an active mechanical role.

3N1458 血管平滑筋ミオシン I による F アクチン束化に対するリン脂質、カルシウム及びガリモジュリンの作用

F-actin bundling by smooth muscle myosin I and the regulatory roles of phospholipids, calcium ion and calmodulin

Kai Sunada, Syougo Tokuda, Masaru Yosida, Tatsuya Kwai, Yoh Okamoto (Muran Institute of Technology, Division of Applied Science)

It has been reported that actin cytoskeletal dynamics are occurring during the activation of smooth muscle contraction.(S.J.Gunst and W.Zhang(2008)) We have isolated and characterized a single head myosin from vascular smooth muscle (Hasegawa Y. et al.(1996,1998)). We have found the myosin I induced F-actin bundling at physiological ionic condition.(Okamoto, Y. et al. (2010). The F-actin bundling has been inhibited by phosphatidyl glycerol, IP₃ and PIP₂ but not with phosphatidyl choline. These observations are consistent with our previous results of the binding abilities to the myosin I, presumably to the short tail (Hasegawa Y.et al (1996). We have done further characterization of F-actin bundling activity of the myosin I with respect to the contributions of calcium ion, calmodulin and a structure of actin. The bundling has been inhibited with increase of calcium ion concentration. We are screening the essential site on actin for the bundling by myosin I. A chemical modification of actin also caused a remarkable inhibition of the bundling with myosin I. The F-actin bundling with smooth muscle myosin I appears to be a process depending upon the lipid binding to the short tail and also calcium ion binding to the calmodulin at the junction between the head and tail of myosin I.

3N1522 蛍光顕微鏡によるコフィリンのアクチンフィラメントへの協同的結合の観察

Cooperative binding of cofilin to the actin filament examined by fluorescence microscopy

Kimihide Hayakawa^{1,3}, Hitoshi Tatsumi², Kazuki Nishida², Masahiro Sokabe^{1,2,3} (¹Cell Mechanosensing Project, SORST, JST, ²Dept. of Physiol., Nagoya Univ. Graduate School of Medicine, ³Res. Center for Innovative Nanobiodevice, Nagoya Univ.)

ADF/cofilin proteins bind to F-actin and modulate the structure and dynamics of the F-actin. Detailed examinations by electron microscopy (EM) have revealed two types of filaments in the same EM-micrograph, bare filaments and cofilin-decorated filaments under a low cofilin to actin ratio condition, suggesting that cofilin binds to the F-actin in a cooperative manner. However, bare filaments and cofilin-decorated filaments in solution have not been imaged in detail. We examined the fluorescent cofilin-decorated F-actin in solution by fluorescence microscopy. The specimen was prepared based on a newly developed method as follows; rhodamine labeled actin (250 μM) was polymerized in F-buffer and mixed with an equal volume of F-buffer containing iodoacetamide-fluorescein labeled cofilin (125 μM), and incubated for 30 sec. The cofilin to actin ratio was 1 to 2. Actin filaments (0.5 μm to 2 μm) were observed with a fluorescence microscope immediately after (~5 sec) fixation by acetone (95 %). Both bare filaments and cofilin-decorated filaments were detected. In addition to these two types, we found intermittently decorated filaments, along which one or two clusters of cofilin were observed. The size of cofilin cluster was ranged from 0.5 μm to 1 μm in majority of the observations. By contrast, we did not detect filaments decorated with cofilin sparsely and uniformly over the filament. These results consistently suggest that cofilin binding to F-actin is cooperative in solution, which conforms to the results in the past EM studies.

3N1534 電子線クライオトモグラフィー法による神経細胞 NG108-15 の超微細構造観察

The 3D ultrastructure of the intact nerve cells, NG108-15 observed by electron cryo-tomography

Shinji Aramaki¹, Kazuhiro Aoyama², Kouta Mayanagi³, Takuo Yasunaga¹ (¹Dept. of Bioscience and Bioinformatics, Faculty of Computer Science and Systems Engineering, Kyushu Inst. of Tech., ²FEL Company Japan Ltd., Application Lab., ³Medical Inst. of Bioregulation, Kyushu Univ.)

Actin filaments, microtubules, and intermediate filaments, called cytoskeleton, play important roles in cells such as cell adhesion, cell motility, and intracellular transportation. Recently, live cell imaging techniques by light microscopy revealed many dynamic mechanisms of cell motility. However, these methods cannot explain the relationship between molecular structures and function of cells at the molecular levels. Thus we need to elucidate ultrastructure of cells by electron-cryo microscopy. In this study we focus on nerve cell elongation system by cytoskeleton.

First, we succeeded in culturing cell on a carbon holey grid for an electron microscope in order to observe cells. Next, for reconstructing 3D ultrastructure we acquired tilt-series of elongation regions of cells by cryo- TEM (Transmission Electron Microscopy) and STEM (Scanning Transmission Electron Microscopy): After the cells on a grid were rapidly frozen, the electron cryo-microscopy supplied observation of intact cells. Their fourier transform suggested that filament should include actin in the dendrite like structure.

We are going to develop transgenic cells which have two kinds of tags for light microscopy and electron microscopy such as GFP (Green Fluorescent Protein) and methallothionein, respectively, to combine electron microscopy and optical microscopy, refereed CLEM (Correlative Light and Electron Microscope). These revolutionary methods will provide us new insights.

3N1546 リアルタイムビデオナノ計測とコンピュータ制御されたレーザー照射により明らかになる、大腸菌の走化性誘引物質への応答時間

Real-time video nanometry and controlled laser irradiation reveals a cellular response time of *E. coli* cell for chemoattractant

Takashi Sagawa¹, Yu Kikuchi¹, Hajime Fukuoka^{1,2}, Yuichi Inoue^{1,2}, Hiroto Takahashi², Takahiro Muraoka², Kazushi Kinbara², Akihiko Ishijima^{1,2} (¹Grad. Sch. Life Sci., Tohoku Univ., ²IMRAM, Tohoku Univ.)

Chemotaxis is an essential process of signal transduction in bacterial cells. The rotational direction of the bacterial flagellar motor is regulated by chemotactic system. We reported last year that serine photo-released from caged-serine by laser irradiation induced continuous CCW rotation of motors as a response to chemotactic signal. Using the system, we tried to estimate cellular response time, which is the time required for the switching from CW to CCW after photolysis. However, the measured time included diffusion of serine from laser irradiation area to cell in addition to cellular response time.

In this study, to eliminate the time for diffusion of serine, laser was irradiated on the cell directly and the laser power is regulated to decrease effect of the laser irradiation for cell. Also we developed a measurement system to measure the rotational direction in real time (every 0.8 ms) and to be able to irradiate laser beam after CW rotation within 60 ms after the switching from CCW to CW rotation by using this system. The measured response time was about 220 ms. This would mean that about 200ms are needed from the recognition of the serine by the receptor to the response of the cell.

We will measure the response time by multiple motors on a cell to understand process of the propagation of intracellular signal by the recognition of attractant signal. We would like to discuss about this result in annual meeting.

3N1558 細胞内シグナル伝達における CheZ の役割についての考察

Consideration of the role of CheZ in the intracellular signaling of *Escherichia coli* cell

Hajime Fukuoka, Yuichi Inoue, Hiroto Takahashi, Akihiko Ishijima (IMRAM, Tohoku Univ.)

E. coli cell senses extracellular stimuli by receptors to transduce to intracellular signal. Phosphorylated CheY (CheY-P) is an intracellular signal protein and regulates the switching of rotational direction of flagellar motors from counterclockwise (CCW) to clockwise (CW). We recently showed the coordinated switching with delay time of two different motors on a cell, therefore we proposed both a transient increase and decrease in the CheY-P concentration are propagated from the receptor patch to motors with a delay time. However we do not understand how such signal propagation is generated. Especially, the decrease of signal is important because motors need to response coordinately to each signal generated at intervals of sub-seconds order. In this