chromatic adaptation. Unlike the most microbial rhodopsins, the cytoplasmic (CP) half of ASR is unusually hydrophilic due to many polar residues and water molecules. Basically, ASR lacks electrogenic proton transport. However, unusual proton transfer was observed: the Schiff base proton is transferred to Asp217 in the CP domain upon the formation of M photo-intermediate (Shi *et al.*, *J. Mol. Biol.* 2006). Thus, ASR has a potential to move proton in the direction opposite to that of proton-pumping rhodopsins.

In this study, we examined the proton transfer reactions of ASR by using indium-tin oxide electrode, which enables the time-dependent measurements of local pH changes by flash-activated ASR. At the neutral pH, ASR induces the transient alkalization of the local medium, indicating that photolyzed ASR accepts proton from the medium and then releases it at the same side. On the other hand, above pH 8, the sequence of the proton movement is reversed: proton release occurs first, followed by uptake. This suggests that a certain residue acts as proton releasing residue in the alkaline condition. These proton transfers were greatly modified by the mutations of Asp217 and Glu36 locating at the CP domain. Thus, the proton transfers mainly occur through this hydrophilic channel. Based on this analysis, we will discuss the proton transfer reactions of ASR.

2PT178 ウシロドプシンの構造に関する理論研究

A theoretical study on the structure of bovine rhodopsins

Motoshi Kamiya, Shigehiko Hayashi (*Grad. Sch. Sci., Kyoto Univ.*)

Rhodopsin is a photoreceptor in the retina. It consists of the protein moiety opsin and an all-trans-retinal which is derived from Vitamin A. The photoisomerization of the retinal induces the structural change of the protein and eventually activates the associated G-protein, transducin. The purpose of this study is to elucidate the structures of early intermediates during the photoreactions such as lumirhodopsin. Computationally, the structure around Schiff base of the retinal is difficult to be described with the classical forcefield, although accurate description of the structure around the retinal is critically important for the calculations of vibrational frequencies and the excitation energy. Therefore, QM/MM calculations must be involved to obtain reasonable structure. On the other hand, structural rearrangements/relaxations at finite temperature (not only fluctuations) of the system, which are usually not involved in QM/MM methods, shall not be ignored for the structural relaxation which takes more than nanoseconds. In order to fulfill these two requisites, in this study, we employed QM/MM-RWFE-SCF method [Kosugi and Hayashi, *JCTC*, 2011]. The structures and vibrational frequencies of the Schiff base in rhodopsin, bathorhodopsin, and lumirhodopsin will be discussed.

2PT179 グロエオバクターロドプシンの三量体形成に関する研究

Homo-trimeric assembly of a cyanobacterial ion-pump *Gloeobacter* rhodopsin

Takashi Tsukamoto¹, Takashi Kikukawa^{1,2}, Masakatsu Kamiya^{1,2}, Tomoyasu Aizawa^{1,2}, Keiichi Kawano^{1,2}, Kwang-Hwan Jung³, Naoki Kamo^{2,4}, Makoto Demura^{1,2} (¹*Grad. Sch. Life Sci., Hokkaido Univ.*, ²*Fac. Adv. Life Sci., Hokkaido Univ.*, ³*Dept. Life Sci. & Inst. Biol. Interfaces, Sogang Univ.*, ⁴*College Pharm. Sci., Matsuyama Univ.*)

Rhodopsin-like molecules in microorganisms (microbial rhodopsins) play roles as an ion-pump or a photosensor. The four microbial rhodopsins from halophilic archaea have been investigated well so far; bacteriorhodopsin (BR), halorhodopsin (HR), sensory rhodopsin I (SRI) and sensory rhodopsin II (SRII, also called phoborhodopsin). These proteins form intrinsic oligomers depending on their function. The phototaxis sensors SRI and SRII form a tetramer which is composed of two units of receptor-transducer hetero-dimer. On the other hand, the ion-pumps BR and HR form a homo-trimer. This is a unique structural property of them and recent studies have revealed several factors for the trimer formation and stabilization (Tsukamoto *et al.*, *Biophys. J.* 2012). Other than BR and HR, however, little is known about the trimer formation of the ion-pumping rhodopsins.

We have started to investigate the oligomeric structure of *Gloeobacter* rhodopsin (GR), which is an ion-pumping protein discovered from freshwater cyanobacterium, by means of size-exclusion chromatography (SEC) and circular dichroism (CD) spectroscopy. The analysis of SEC gave us a conclusion that GR formed a trimer as well as BR and HR. However, the CD spectrum of GR ((+) 500 nm/(-) 567 nm, crossover point; 540 nm) indicated that the trimeric state of GR was different from that of BR and HR. Interestingly, the trimeric state of GR altered in a pH-dependent manner. Based on these analyses, we will discuss the homo-trimeric structure of GR.

2PT180 パルミトイル修飾とロドプシンのラフト親和性の関係

Relationship between palmitoylations and raftophilicity of rhodopsin

Keiji Seno¹, Fumio Hayashi² (¹*Hamamatsu Univ. Sch. Med.*, ²*Grad. Sch. Sci., Kobe Univ.*)

As we reported in the last meeting of BPSJ, we have found that rhodopsin forms dimer in the complex of retinal G protein transducin (G_t). The rhodopsin dimer favors *l_o* lipids and distributes in the detergent-resistant membrane (DRM) prepared from vertebrate (bovine and frog) rod photoreceptor outer segment membranes. Our single molecule observation of rhodopsin and transducin revealed that the increased affinity to raft of rhodopsin upon dimerization hinders the diffusion of rhodopsin. Here, we studied the contribution of tandem palmitoyls at the C terminal end of helix-8 of rhodopsin to the raftophilicity of rhodopsin dimer. We examined the effect of depalmitoylation of rhodopsin by reducing agent (20 mM dithiothreitol) on the distribution of rhodopsin-transducin complex or IgG-crosslinked (artificially dimerized) rhodopsin to the DRM of frog rod photoreceptor outer segments. Depalmitoylation was confirmed by MALDI-TOF MS. Rhodopsin-transducin complex in DRM fraction was reduced to almost half by depalmitoylation. When we artificially dimerized rhodopsin by anti-rhodopsin monoclonal antibody, significant portion of anti-rhodopsin IgG, which means rhodopsin dimer, was detected in DRM fractions, however, when we used depalmitoylated rhodopsin, much less IgG was detected in DRM fractions. These data support our idea that the palmitoyl modifications on rhodopsin are indispensable for raftophilicity of rhodopsin dimer.

2PT181 海洋性細菌におけるプロテオロドプシンの生理的役割および分子特性

Physiological role and molecular properties of proteorhodopsin from marine bacterium

Rei Abe-Yoshizumi¹, Keiichi Inoue¹, Susumu Yoshizawa², Kazuhiro Kogure², Hideki Kandori¹ (¹*Dept. of Frontier Materials, Nagoya Inst. Tech.*, ²*AORI, Univ. Tokyo*)

Nature created two kinds of light-driven ion pumps, bacteriorhodopsin (BR) and halorhodopsin (HR), functioning as outward proton and inward chloride pumps, respectively [1]. Then, thousands of proteorhodopsin (PR) have been discovered and are widely distributed among marine prokaryotes [2]. But its physiological role was unclear, because very few strains carrying the PR gene have been cultured. Recently, we demonstrated light-driven proton pump activity of PR in native *Flavobacteria* cells [3].

Here we studied the molecular mechanism and the physiological role of PR from *Krokinobacter* sp. in *Flavobacterium* that codes two microbial rhodopsin genes in their genome. We first examined their pump activity in native cells. It was found that PR is expressed in stationary phase, suggesting that PR-mediated electrochemical proton gradient is used to ATP synthesis when nutrition is depleted. We then studied the spectral and photochemical properties of recombinant PR from *Krokinobacter* (KroPR). KroPR shows λ_{max} at 525 nm (pH 7.0), being similar to GPR. On the other hand, pK_a of the counterion (4.3) is lower than that of GPR (6.9), although the amino acid identity between KroPR and GPR is 41 % and His75 (GPR) is conserved. Flash photolysis measurement of KroPR is in progress and the molecular mechanism of KroPR will be discussed.

[1] Kandori, *Mol. Sci.* 5, A0043 (2011).

[2] Beja *et al.* *Science* 289, 1902 (2000).

[3] Yoshizawa *et al.* *Environ. Microbiol.* 14, 1240 (2012).

2PT182 時間分解赤外分光法によるファラオニスハロロドプシンの機能発現過程での水の水素結合ダイナミクス

Water hydrogen-bonding dynamics in the function of *pharaonis* halorhodopsin studied by time-resolved FTIR spectroscopy

Kuniyo Fujiwara^{1,2}, Tetsunari Kimura^{1,2,3}, Takashi Kikukawa⁴, Makoto Demura⁴, Hideki Kandori⁵, Yuji Furutani^{1,2,6} (¹*SOKEINAI*, ²*IMS*, ³*JST CREST*, ⁴*Grad. Sch. Life Sci., Hokkaido Univ.*, ⁵*Grad. Sch. Eng., NITech*, ⁶*JST PRESTO*)

pharaonis Halorhodopsin (pHR) is an inward chloride ion pump protein. pHR has a retinal as the chromophore, which is bound through the protonated Schiff base and all-*trans* to 13-*cis* photoisomerization triggers conformational changes of the protein moiety for pumping a chloride ion. The ion transportation is performed through the sequential formation of some intermediates which are named K, L₁, L₂, N and O. The structural detail of each intermediate state is still

unclear, and especially, it is open question how water molecules involve in the translocation of a chlorine ion inside protein during the photocyclic reaction. To analyze the hydrogen-bonding dynamics of water and protein, we performed the time-resolved Fourier transform infrared (TR-FTIR) spectroscopy in the whole mid-infrared region. The measurement was performed under the condition of hydration with H₂O or H₂¹⁸O, which allow us to assign the water O-H stretching bands. As a result, we observed a water band at 3626 cm⁻¹ which exhibited down shift to 3605 cm⁻¹ upon the formation of the N and O states. We also analyzed the dependency on the chloride concentration to discriminate whether the water change occurs in the N or O intermediates, which are related to release and uptake of chloride ion. Anion dependency (Br⁻, NO₃⁻ and SO₄²⁻) was investigated as well. By comparing these experimental data, the relationship between the water hydrogen-bonding dynamics and the conformational changes of protein will be discussed.

2PT183 全反射赤外分光法によるキメラチャンネルロドプシンの構造変化解析
Structural changes in the chimeric Channelrhodopsins revealed by ATR-FTIR spectroscopy

Asumi Inaguma^{1,2}, Hisao Tsukamoto¹, Tetsunari Kimura^{1,3}, Toru Ishizuka^{3,4}, Hiromu Yawo^{3,4}, Yuji Furutani^{1,2} (¹*Inst. Mol. Sci.*, ²*JST PREST*, ³*JST CREST*, ⁴*Grad. Sch. Life Sci., Tohoku Univ.*)

Channelrhodopsin 1 and 2 (ChR1 and ChR2) can function as light-gated cation channels which were discovered from *Chlamydomonas reinhardtii*. Recently, ChR2 and its derivatives have been utilized as a useful tool to control the neural circuit by light. ChR1/2 chimera such as ChR-wide receiver (ChR-WR) and ChR-fast receiver (ChR-FR) have improved characters as the electrophysiological tool. ChR-WR is sensitive to the light in the wider range of wavelength compare to the wild type. ChR-FR exhibits faster ON/OFF kinetics in the photocurrent. However, the molecular mechanisms behind these useful properties have not been understood well. Nowadays, the crystal structure of ChR has been reported, but the structural changes connecting the closed and open conformation have not been answered well. In this research, we applied attenuated total reflection Fourier transform infrared (ATR-FTIR) spectroscopy to reveal the structural changes of the chimeric ChRs during the formation of the open/photostationary state and return to the closed state. The light-induced difference FTIR spectra of ChR-WR and ChR-FR were very similar to each other, suggesting that the structural changes are mostly same. Interestingly, the recovery from the photostationary state of ChR-FR was faster than ChR-WR. We will discuss the relationship between the structural changes and the photocurrent properties of these chimera ChRs.

2PT184 Crystal Structure of deltarhodopsin from Haloterrigena Thermotolerans

Jin Zhang, Katsuhide Mizuno, Mizuno Kouyama (*Nagoya University*)

Deltarhodopsin, a new member of the microbial rhodopsin family, functions as a light-driven proton pump. In this study, we investigated the crystal structure of deltarhodopsin (dR3) from *Haloterrigena thermotolerans*, a thermo-tolerant archaeon isolated from a solar saltern in Puerto Rico. For preparation of a large quantity of the target protein, the gene encoding deltaopsin-3 was transformed into a bR-deficient host strain of *Halobacterium salinarum*. When the membrane fusion method was applied, the purified sample of dR3 was crystallized into trigonal crystals belonging to space group R32. In this crystal form, dR3 forms a trimeric assembly as observed for bR and other archaeal rhodopsins. It is demonstrated that the inner part of the protein (the retinal binding pocket and the proton release and uptake pathways) is similar to that of bR and other microbial rhodopsins. One of major differences is observed in the protein-protein contact regions. Interestingly, the carotenoid bacterioruberin binds to the crevice between the AB helices of one subunit and the DE helices of a neighboring subunit. Another peculiar property of dR3 is a highly crowded distribution of positively charged residues on the cytoplasmic surface. Structural comparison of dR3 with other archaeal rhodopsins allows us to recognize the structural motif that is relevant to the proton pumping activity, while also offering insight into the formation mechanism of the trimeric structure and the recognition mechanism of specific lipid components.

2PT185 The chloride ion uptake mechanism in pharaonis halorhodopsin as studied by the energy representation method

Hiroyuki Tamura¹, Shuntaro Chiba¹, Shun Sakuraba², Nobuyuki Matsubayashi², Minoru Sakurai¹ (¹*Center for Biol. Res. & Inform. Tokyo Tech*, ²*ICR, Kyoto Univ*)

Halorhodopsin (hR) is a light-driven chloride ion pump exist in halophilic archaea. hR uses the light energy to move chloride ions from extracellular side to cytoplasmic side. The process is thought to be divided into the two steps: (i) a chloride ion passively enters the binding site near retinal Schiff base linkage in the protein from the extracellular medium with a high NaCl concentration (~4M), and (ii) the chloride ion is transported into the cytoplasmic side after the photoisomerization of retinal. The purpose of this work is to clarify the mechanism of the first step by calculating the free energy of a chloride ion in the following two cases: it is located in the extracellular side (bulk) and at the binding site. The free energies were calculated using the method of energy representation combined with molecular dynamics simulations. First, we calculated the free energy of a chloride ion in the aqueous solution of NaCl. As a result, the free energy decreased from -88.6 to -122.0 kcal/mol as the NaCl concentration was increased from 0M to 4M. Next, we calculated the free energy of a chloride ion at the binding site: the result was -120.3 kcal/mol, which is equal to that in the aqueous solution of 3.4 M NaCl. These results indicate that a chloride ion in bulk passively enters the binding site if the NaCl concentration in bulk is lower than 3.4 M. The largest contributor to the stabilization of a chloride ion at the binding site is the chromophore (-49.3 kcal/mol) and the next is the nearby water molecule (-20.5 kcal/mol).

2PT186 ATP ショットと caged-ATP の光分解により開始されるホタル生体発光の時間発展及びその pH 依存性
Time Dependent Characteristics of Firefly Bioluminescence Initiated by Two Methods with Usual ATP Injection and Photolysis of Caged-ATP

Yuki Yanagisawa¹, Takeshi Kageyama¹, Masatoshi Tanaka¹, Shin-ya Ohno¹, Naohisa Wada² (¹*Faculty of Engineering, Yokohama National University*, ²*Department of Life Sciences, Toyo University*)

The time-resolved spectrum and the rise time course of firefly bioluminescence initiated by ATP injection into the HEPES buffer containing luciferin, luciferase, and Mg²⁺ were first measured with time resolution of 10 ms, and compared with those of the bioluminescence triggered by photolysis of caged-ATP. The time course is significantly dependent on pH; both time constants of rise and decay increase with decreasing pH from 7.8 to 6.8. The observed time course is constituted by two components; the diffusion-controlled reaction follows the homogenization of reactants. The usual ATP injection may prevent ATP⁴⁻ from falling into the catalytic center of Luc. Contrarily, in the case of the bioluminescence triggered by photolysis of caged-ATP, the intrinsic reaction dominates the time course not correlative with pH [1]. The time-resolved spectra can be decomposed into two Gaussians peaked at 2.0 eV and 2.2 eV. The intensity of 2.2 eV component decreases with decreasing pH, being consistent with previous report [2]. Spectral shapes do not change from the initiation of the enzymatic reaction up to 200 s for both methods of ATP addition. After 200 s, the 2.0 eV component decays slower than the 2.2 eV component in the case of the photolysis of caged-ATP, which may be caused by the product of photolysis. Interestingly, above results are not influenced regardless of the concentration ratio [Ln]/[Luc].

[1] T. Kageyama, et al., Photochem. Photobiol. 87 (2011) 653-658.

[2] Y. Ando, et al., Nature Photonics 2 (2008) 44-47.

2PT187 水溶液中のホタルルシフェリン蛍光スペクトルの理論的研究
Theoretical Analysis of Fluorescence Spectra of Firefly Luciferin in Aqueous Solutions

Miyabi Hiyama¹, Hidefumi Akiyama², Kenta Yamada³, Nobuaki Koga¹ (¹*Grad. Sch. Info. Nagoya Univ.*, ²*ISSP Univ. Tokyo*, ³*Grad. Sch. Nanobio. Yokohama City Univ.*)

To understand the mechanism of firefly bioluminescence, researchers have investigated the spectroscopic properties of firefly luciferin as an analog of oxyluciferin [1-6]. Despite the previous experimental and theoretical studies, some peaks in the fluorescence spectra of firefly luciferin still remain unassigned. To elucidate the origin of fluorescence spectra of firefly luciferin, the excited states of firefly luciferin and its conjugate acids and bases were investigated by performing the density functional theory calculations (DFT). Time-dependent DFT calculations were performed for the excited states of these molecules. The solvent effect in aqueous solution was taken into account by the polarizable continuum model. We obtained the energies and oscillator strengths of excitation of firefly luciferin and its conjugate acids and bases [7]. Although the wavelengths of energies in theoretical fluorescence spectra are approximately 40 nm lower than those in the experimental data, the relative